

# Landscape Genetics

## How Landscapes Affect Ecological Processes

Janine Bolliger, Gwenaëlle Le Lay,  
Rolf Holderegger

*Landscape genetics evaluates the effect of landscape structure and composition on the movement, dispersal and gene flow of animals and plants. It is applied, for instance, to test whether de-fragmentation measures such as green bridges or landscape corridors effectively enhance the movement of individuals in fragmented landscapes.*



Schweizerische Akademische  
Gesellschaft für Umweltforschung  
und Ökologie

Société Académique Suisse pour  
la Recherche sur l'Environnement  
et l'Ecologie

Swiss Academic Society for  
Environmental Research  
and Ecology

**Landscape Genetics – How Landscapes Affect Ecological Processes** | GAIA 19/3 (2010): 238–240

**Keywords:** connectivity, conservation management, dispersal, fragmentation, genetic techniques, landscape ecology, spatial analysis

### Combining Population Genetics and Landscape Ecology

Landscape genetics seeks to assess how landscape structure and composition affect movement, dispersal and gene flow of animals and plants. Landscape genetics has recently been promoted in review articles (e.g., Holderegger and Wagner 2008) and at international conferences (e.g., Balkenhol et al. 2009). The research field amalgamates population genetics and landscape ecology by combining the theory and methods of the former with the GIS-applications, spatial analyses and modelling of the latter (Manel et al. 2003). While landscape ecology stresses the dynamics of patterns and processes in changing landscapes by characterizing landscape structure (e.g., suitable versus unsuitable habitats, barriers versus corridors, etc.), population genetics uses various genetic techniques to indirectly measure how animals and plants

(and therefore their genes) move and disperse across landscapes. By merging the two fields, landscape genetics provides a powerful tool to identify the effects of landscape properties on movement, dispersal and gene flow. Given the appeal of landscape genetics, both in science and application, ETH Zurich will offer a one-week winter school on landscape genetics next February (box 1) and SAGUF will introduce this new field to its members at its annual meeting in 2011 (box 2).

### An Innovative Research Field

The combination of genetic and landscape or environment data with modelling approaches in landscape genetics offers a setting to test emerging questions of species behavior under land-use and climate change. A prominent example of landscape genetic research and its application in conservation management relates to the effectiveness of de-fragmentation measures. The basic question is whether implemented structural connectivity measures lead to functional connectivity; the latter being the effective exchange of individuals or genes among populations. For instance, cantonal authorities in Switzerland already use landscape genetic approaches to assess the degree of population fragmentation due to highways, major roads, fenced railways or

settlements in wildlife corridors of national importance. This knowledge of the present situation will also enable the future assessment of the effectiveness of mitigation measures such as green bridges or underpasses once they have been implemented. Similarly, landscape genetic methods are used to study the effectiveness of connectivity measures taken on agricultural land under national or international agri-environmental schemes. In doing so, landscape genetics develops into a valuable planning and monitoring tool in practical management. What we have learned from landscape genetics so far shows that especially animals but also plants more often spread over larger distances than had been suggested hitherto (Holderegger and Wagner 2008).

### Approaches and Challenges

Currently, landscape genetics is using three approaches to detect the effects of landscape structure and composition on movement, dispersal and gene flow, namely 1. approaches based on genetic discontinuities (*Where can distinct gene pools be found in a landscape?*), 2. approaches based on genetic distance (*How different are two populations or individuals in genetic terms?*) and 3. approaches assessing current movement and gene flow.

**Contact authors:** PD Dr. Janine Bolliger | SAGUF Board Member | WSL Swiss Federal Research Institute | Birmensdorf | Switzerland | E-Mail: janine.bolliger@wsl.ch

**Contact SAGUF:** SAGUF-Geschäftsstelle | Dr. Claudia Zingerli | ETH Zürich D-UWIS/CHN | 8092 Zurich | Switzerland | E-Mail: saguf@env.ethz.ch | <http://saguf.snatweb.ch>

1. Approaches based on ***genetic discontinuities*** investigate whether the boundaries between genetically distinct populations (i.e., populations that are not in contact with each other) spatially coincide with particular landscape structures (e.g., roads, rivers or settlements) on topographic or land use/land cover maps, which potentially act as barriers to movement and dispersal. Various clustering methods are used to assign individuals to genetic pools. An empirical example is provided by Riley et al. (2006). They showed that coyotes and bobcats form genetically isolated groups on different sides of a Californian highway and that the highway therefore acts as a barrier to gene flow.
2. ***Genetic distance*** provides a measurement of how often populations or individuals come into mating contact. In other words, genetic distance is a measure of the abundance of movement, dispersal and gene flow between two localities. Approaches based on genetic distance use GIS data on landscape elements such as forests, roads, open fields or meadows. Most often, a resistance value is given to each landscape element, reflecting the permeability of the element to the movement of a given species. This procedure allows calculating a minimum cumulative landscape resistance between two localities. This least cost distance describes a potential movement path minimizing the costs of movement through a particular landscape. The correlation between genetic distance and different least cost distance models can then be evaluated and allows identifying those landscape elements that mainly influence movement, dispersal and gene flow. Using such an approach, and regarding roe deer dispersal in a fragmented landscape, Coulon et al. (2004) showed that roe deer strongly follows woodland patches and avoids crossing open land without shelter.
3. The last approach estimates the ***contemporary movement of individuals or current gene flow***. Assignment tests determine contemporary migrants among populations based on their genetic com-

**BOX 1:****Winter School in Landscape Genetics at ETH Zurich**

February 14 – February 19, 2011

WSL Swiss Federal Research Institute, Birmensdorf, Switzerland

This course will be held by *Rolf Holderegger, Janine Bolliger* and *Gwenaelle Le Lay*. It will introduce students to major concepts and methods of landscape genetics. Special emphasis is given to current state-of-the-art methods and to hands-on exercises. The course will deal with types of genetic and landscape ecological data for landscape genetic analysis, landscape resistance and gene flow (least cost paths, Mantel tests, regression with permutation), overlays and genetic boundaries (clustering, barrier detection, kriging), contemporary gene flow (paternity analysis, assignment tests), identification of genetic markers of adaptive relevance (i.e., landscape genomics) and methods for model evaluation. The benefits and limitations of landscape genetics will also be highlighted. Basic knowledge in population genetics and GIS is required.

**MORE INFORMATION:**

- Rolf Holderegger, E-Mail: [rolf.holderegger@wsl.ch](mailto:rolf.holderegger@wsl.ch)
- Janine Bolliger, E-Mail: [janine.bolliger@wsl.ch](mailto:janine.bolliger@wsl.ch)

position. In other words, given the genetic composition of a given individual, to which population of origin does it most probably belong? The genetic composition of most individuals matches that of the population where they were sampled – these are residents. However, some individuals will be assigned to a population different from where they were sampled – these are migrants from other populations. Using such an assignment test, Angelone and Holderegger (2009) showed that functional connectivity at the landscape scale in the valley of the Swiss river Reuss could be achieved through the construction of new ponds; the ponds increased the movement of tree frogs, thus acting as stepping stones between existing populations. Another method to identify contemporary gene flow,

called parentage analysis, has mainly been used in plants. Kamm et al. (2009) found that even widely scattered individuals of a rare forest tree species in northern Switzerland form part of an extensive network connected by regular pollen exchange over distances of more than ten kilometres. These trees, although spatially isolated, were functionally well connected.

**Outlook**

Many of the above mentioned methods and approaches still suffer from conceptual and methodological shortcomings (Balkenhol et al. 2009). Future progress in landscape genetics depends on the development of appropriate analytical tools allowing a better assessment of the relationships between genetic and landscape data. Hence, close collaboration between land-

**BOX 2:****SAGUF-Jahrestagung 2010****Erfahrungsaustausch zu inter- und transdisziplinärer Praxis**

25. November 2010, 9 bis 16 Uhr, Altes Spital, Solothurn, Schweiz

Die diesjährige SAGUF-Jahrestagung findet statt als Workshop zum Thema ***Erfahrungsaustausch zu inter- und transdisziplinärer Praxis*** und wird von der SAGUF und dem Nationalen Forschungsprogramm *Nachhaltige Wassernutzung* (NFP 61) gemeinsam organisiert. Die Teilnahme am Workshop ist beschränkt auf SAGUF-Mitglieder und Forschende des NFP 61.

**Anmeldung bis 31. Oktober 2010 an:** SAGUF Geschäftsstelle, Dr. Claudia Zingerli, ETH Zentrum CHN, 8092 Zürich, Schweiz, E-Mail: [saguf@env.ethz.ch](mailto:saguf@env.ethz.ch)

**Kontakt:** Dr. Patricia Fry, Wissensmanagement Umwelt GmbH, Umsetzungsbeauftragte NFP 61, Leiterin AG Wissensaustausch SAGUF, Idaplatz 3, 8003 Zürich, Schweiz, Tel.: +41 44 4613327, E-Mail: [contact@patriciafry.ch](mailto:contact@patriciafry.ch)

**Vorschau:** Thema der SAGUF-Jahrestagung 2011: ***Landschaftsgenetik***

scape ecologists and population geneticists is required, since only statistically robust and easy-to-use approaches will allow landscape genetics to expand its full power with respect to application in management practice (Segelbacher et al. 2010). One particular advantage of landscape genetics is that large spatial areas (i.e., whole landscapes) can be monitored over extended time periods. Therefore, landscape genetics can provide the whole landscape view of functional fragmentation or connectivity that is urgently needed in conservation and landscape management.

future research needs in landscape genetics: Where to from here? *Landscape Ecology* 24: 455–463.

Coulon, A. et al. 2004. Landscape connectivity influences gene flow in a roe deer population inhabiting a fragmented landscape: An individual-based approach. *Molecular Ecology* 13: 2841–2850.

Holderegger, R., H. H. Wagner. 2008. Landscape genetics. *BioScience* 58: 199–207.

Kamm, U., P. Rotach, F. Gugerli, M. Siroky, P. Edwards, R. Holderegger. 2009. Frequent long-distance gene flow in a rare temperate forest tree (*Sorbus domestica*) on the landscape scale. *Heredity* 103: 476–482.

Manel, S., M. Schwartz, G. Luikart, P. Taberlet. 2003. Landscape genetics: Combining landscape

## Landscape genetics offers a setting to test emerging questions of species behavior under land-use and climate change.

### References

- Angelone, S., R. Holderegger. 2009. Population genetics suggests effectiveness of habitat connectivity measures for the European tree frog in Switzerland. *Journal of Applied Ecology* 46: 879–887.
- Balkenhol, N., F. Gugerli, S. A. Cushman, L. P. Waits, R. Holderegger, H. H. Wagner. 2009. Identifying ecology and population genetics. *Trends in Ecology and Evolution* 18: 189–197.
- Riley, S. et al. 2006. A southern Californian freeway is a physical and social barrier to gene flow in carnivores. *Molecular Ecology* 15: 1733–1741.
- Segelbacher, G. et al. 2010. Applications of landscape genetics in conservation biology: Concepts and challenges. *Conservation Genetics* 11: 375–385.



© Daniel Keller

Landscape genetics is well suited to complement small-scale ecological studies on the dispersal of marked animals – here a marked damselfly – on larger spatial scales.

## GAIA ECOLOGICAL PERSPECTIVES FOR SCIENCE AND SOCIETY

© 2010 Verein Gaia | Konstanz, St. Gallen, Zürich

### HAUPTHERAUSGEBERIN

Verenा Winiwarter | Wien  
(verantwortlich im Sinne des Presserechts)

### REDAKTION

Dr. Susanna Bucher/Dr. Almut Jödicke | ETH Zentrum | Postfach CAB 42 | 8092 Zürich | Schweiz | Tel.: +41 52 2222805 | E-Mail: redgai@env.ethz.ch

Dr. Ulrike Sehy | oekom verlag | Hagenbuchrain 13 | 8047 Zürich | Schweiz | Tel.: +41 43 3110785 | E-Mail: sehy@oekom.ch

Tobias Mickler | oekom verlag | Waltherstr. 29 | 80337 München | Deutschland | Tel.: +49 89 54418445 | E-Mail: mickler@oekom.de

### REDAKTIONELLE MITARBEIT

Dr. Maiken Winter

### GRAFISCHE GESTALTUNG UND SATZ

Heike Tiller | München | E-Mail: h.tiller@freenet.de

### VERLAG

oekom verlag – Gesellschaft für ökologische Kommunikation mbH | Waltherstr. 29 | 80337 München | Deutschland | [www.oekom.de](http://www.oekom.de) | Gesellschafter und Anteile: Jacob Radloff, Feldafing, 77 Prozent, und Christoph von Braun, München, 23 Prozent

### ANZEIGEN

Stefanie Ott (verantwortlich) | oekom GmbH | Tel.: +49 89 54418425 | E-Mail: anzeigen@oekom.de

### DRUCK

Kessler Druck + Medien | 86399 Bobingen | Deutschland | [www.kesseldruck.de](http://www.kesseldruck.de)

Die Zeitschrift und alle ihre Inhalte sind urheberrechtlich geschützt. Jede Verwertung, die nicht ausdrücklich vom Urheberrechtsgesetz zugelassen ist, bedarf der Zustimmung des Vereins Gaia. Namentlich gekennzeichnete Artikel müssen nicht die Meinung der Herausgeber/Redaktion wiedergeben. Unverlangt eingesandte Manuskripte, für die keine Haftung übernommen wird, gelten als Veröffentlichungsvorschlag zu den Bedingungen des Verlags. Es werden nur unveröffentlichte Originalarbeiten angenommen. Die Verfasser(innen) erklären sich mit einer nicht sinnentstellenden redaktionellen Bearbeitung einverstanden.

**ERSCHEINUNGSWEISE** Viermal im Jahr.

### BEZUGSBEDINGUNGEN

Probeabonnement (2 Ausgaben inkl. Versand, nur in Deutschland): 19,90 EUR | Jahresabonnement: 95,80 EUR für Privatpersonen; 167,90 EUR für Unternehmen, Institutionen, Bibliotheken; 64,90 EUR für Studierende, Auszubildende, Schüler(innen) (jährlicher Nachweis erforderlich) | Einzelheft 26.– EUR. Alle Preise inkl. MwSt., zzgl. Porto- und Versandkosten. Abbestellungen sechs Wochen vor Ablauf des Abonnementjahrs. Zahlungen im Voraus.

### BESTELLUNG, ABOVERWALTUNG, VERTRIEB

Rhenus Medien Logistik GmbH & Co. KG | Auslieferung oekom verlag | 86899 Landsberg am Lech | Deutschland | Tel.: +49 8191 97000608 | Fax: +49 8191 97000405 | E-Mail: oekom@de.rhenus.com [www.oekom.de/gaia/abonnement](http://www.oekom.de/gaia/abonnement)

KONTO für Deutschland: Postbank München | BLZ 70010080 | Konto 358 744 803  
für die Schweiz: PostFinance SWISS POST | BLZ 30001101 | Konto 40-344357-9

oekom ist der erste klimaneutrale Verlag Deutschlands.

**Climate Partner**

ISSN 0940-5550

Printed in Germany.  
Gedruckt auf FSC-zertifiziertem Papier.

