Fishing in the water: effect of sampled water volume on environmental DNA-based detection of macroinvertebrates

Elvira Mächler^{1,2*}, Kristy Deiner^{1,3}, Fabienne Spahn⁴ & Florian Altermatt^{1,2}

¹ Eawag, Swiss Federal Institute of Aquatic Science and Technology, 8600 Dübendorf, Switzerland

² Department of Evolutionary Biology and Environmental Studies, University of Zurich, Winterthurerstr. 190, 8057 Zürich, Switzerland

³ University of Notre Dame, Department of Biological Sciences, 180 Galvin Life Sciences, Notre Dame, IN 46556 USA

⁴ ETH Zurich, Department of Biology, 8093 Zürich, Switzerland

* corresponding author: elvira.maechler@eawag.ch, phone: +41 58 765 6734, fax: +41 58 765 5802

Keywords: PCR replication, eDNA, river system, detection rate, *Ancylus fluviatilis*, *Baetis buceratus*, *Gammarus pulex*.

This document is the accepted manuscript version of the following article:
Mächler, E., Deiner, K., Spahn, F., & Altermatt, F. (2016). Fishing in the water: effect of sampled water volume on environmental DNA-based detection of macroinvertebrates. *Environmental Science and Technology*, 50(1), 305-312.
<https://doi.org/10.1021/acs.est.5b04188>

Abstract

Accurate detection of organisms is crucial for effective management of threatened and invasive species because false detections directly affect implementation of management actions. The use of environmental DNA (eDNA) as a species detection tool is in a rapid development stage, however, concerns about accurate detections using eDNA have been raised. We evaluated the effect of sampled water volume (0.25 to 2 L) on the detection rate for three macroinvertebrate species. Additionally, we tested, depending on the sampled water volume, what amount of total extracted DNA should be screened in order to reduce uncertainty in detections. We found that all three species were detected in all volumes of water. Surprisingly, however, only one species had a positive relationship between an increased sample volume and an increase in the detection rate. We conclude that the optimal sample volume may depend on the species-habitat combination and should be tested for the system where management actions are warranted. Nevertheless, we minimally recommend sampling water volumes of 1 L and screening at least 14 μ L of extracted eDNA for each sample to reduce uncertainty in detections when studying macroinvertebrates in rivers and using our molecular workflow.

Introduction

By shedding hairs, cells, gametes or feces, all organisms leave traces of their occurrence in the environment in the form of so-called environmental DNA (eDNA). Recent reviews illustrate that many eukaryotes, including plants and animals, are readily and non-invasively detected from traces of their DNA found in water, soil and air.^{1,2} The utility of this non-invasive molecular method for species detection has large implications for environmental management actions³ and subsequent policy and stakeholder decisions e.g.,⁴.

The method of detecting macro-eukaryotic species from traces of their DNA in the environment is, however, in a rapid development phase.^{1,3} Experimental evidence of the power, but also the limitations of the tool is greatly needed in order to effectively track organisms in their environment from eDNA. Of particular concern are false negative detections, for example see^{5,6,7}. False negative detection, or process type II error, in terms of environmental DNA, means that there is no DNA detected although the species is present at the sampled location (please note that false negative detections can also occur during “classic sampling”, see for example⁸). False negative detections may be of particular concern for invasive species and species threatened with extinction because not detecting the species can have severe consequences for management decisions. It is therefore paramount that we gain a better understanding of the causes of false negative detections with eDNA such that it can be established as a viable and defensible method for species detection.^{2,9}

Reasonable work has already focused on the causes of false negative detections and the possible means by which false negatives can be reduced. Causes of false negatives are due to the inherent problem of the detection limits of the molecular technology used to capture, extract and amplify the DNA found in the environment¹⁰⁻¹³ but, in the case of meta-barcoding approaches,

also due to limitations of reference databases. The latter are not discussed here further, as we focus on a targeted approach, but see for example^{14,15}. Far less attention, however, has been paid on estimating how sampling bias affects the detection rate for a species' eDNA and subsequently resulting in false negative detections. Any biological sample suffers from a sampling bias based on the probability of detection and is governed by the method(s) used for detection.¹⁶ Environmental DNA detection is no exception and has mainly two steps which can cause a sampling bias either in the field and/or in the lab.⁵ The first sampling bias can happen when a particular amount of the environmental sample is collected, such as a set volume of air, soil or water. The second sampling bias can happen when amplifying the targeted species' DNA from a small fraction of the total purified DNA contained in the sample. Both of these steps involve subsampling the potential pool of DNA that is tested and can result in a false negative detection simply due to sampling error.

Focusing on freshwater, there are two possible environments to collect DNA from, namely from the sediment or from the water column. Here we focus on estimating detection rates from water, because DNA extracted from water has been commonly used in many macro-organisms' detection protocols (Table 1). Water-based samples are thought to reflect the contemporary, regional community of macroinvertebrates, while sediment-based samples are more likely reflecting local communities, possibly integrating over time. It was not the goal of our study to compare these two methods, as the methods differ and are covering different, complementing aspects. For a comparison and discussion of sediment-based samples, see for example^{5,17}.

In a water-based approach, a first sampling bias can be introduced when different sets of water volumes are used. Likely, the volume of water sampled in different studies (for an

overview on volumes used in previous studies using filtration, see Table 1) has been a choice of practicality based on the specific field and molecular protocol used to capture and concentrate the eDNA (i.e., the logistical aspects of sampling) and is not necessarily a reflection of the optimal amount needed to reduce uncertainty in the detection rate. For example, when eDNA is captured from freshwater through precipitation, usually 15 mL of water is used due to the limitation of centrifuge size needed for the next step to process the water in most standard molecular laboratories. Filtration from freshwater as a capture method is more flexible with respect to volume and previous studies have thus used volumes ranging from 100 mL¹⁸ up to 10 L¹⁹, with an average of approximately 2 L (Table 1). Thus, when left to interpretation and method choice, one could justify to sample and filter 100 mL to 10 L, but it is unclear if and how the volume of water sampled affects the detection rate for a species.

A second sampling bias can occur at the polymerase chain reaction (PCR) stage, where a wide range of total volume of extracted eDNA screened for targeted DNA has been used (Table 1). This is further confounded by the fact that varying molecular protocols have been used for purification of DNA from freshwater, such that the total DNA screened in addition to the total eDNA recovered from the sampled water is also a possible confounding factor.¹⁰ Surprisingly, we have no clear evidence pointing to an optimal amount of extracted eDNA needed to reduce false negative detections (Table 1). For example, Goldberg *et al.*¹⁹ screened 1 μ L (but extracted DNA from 10 L), while Wilcox *et al.*¹² screened 128 μ L of their DNA extraction (Table 1). It is known, that the PCR has an inherent stochastic component, which plays a major role when DNA concentrations are low. As this is typical for extracted DNA from environmental samples, the stochastic component of PCR needs to be considered when performing PCR on eDNA. Overall, it is not completely clear how much volume of the total extracted DNA should be screened to have a precise estimate of the effect on the detection rate for a species.

In this study we tested the effect of sampling at these two stages and how it affects false negative detections. Specifically, we sampled DNA from the environment in different volumes of water and then analyzed different volumes of DNA extracted from a given amount of water. We sampled independent volumes of water ranging from 250 mL to 2000 mL and tested for the detection of three macroinvertebrate species belonging to the orders of Mollusca, Ephemeroptera and Amphipoda at a location in a river where all three species are known to be present. We compared the detection rate with respect to the volume of sampled water and the volume of extracted DNA that was screened. We conclude with recommendations for optimal volumes of water to sample and how much volume of extracted DNA to screen in order to reduce false negative detections. Our recommendations refer to a similar set of species and study systems (i.e., macroinvertebrates in rivers). For other species or habitats they may be used as first guiding values.

Material and Methods

Field sampling

Our study site was located at the river Glatt (47° 26' 35.21" N, 8° 33' 03.94" E). It is a natural river belonging to the headwaters of the river Rhine catchment in Switzerland (for pictures of the study river, see²⁰). We sampled water on September 17th and 30th in 2014. These sampling dates reflect classic sampling time points for macroinvertebrate (commonly done either in spring and/or fall). We sampled at two time points to avoid spurious effects due to a given day's hydrological regime, but close enough to each other to avoid changes in communities looked at. On each day we sampled two replicates of each of the following volumes: 250 mL, 500 mL, 1000 mL, 1500 mL and 2000 mL. This range of volumes was chosen based on previous work (see Table 1) to be suitable for our habitat, that is, freshwater streams and macroinvertebrates, and is

reflecting most of previous volumes considered. We sampled each volume independently in one or two individual 1-L sterile octagonal polyethylene terephthalate bottles (VWR International, Radnor, Pennsylvania, USA) that were previously decontaminated with 10% household bleach, rinsed with Milli-Q® (Merck Millipore, EMD Millipore Co., Billerica, Massachusetts, USA) water and exposed to ultraviolet C light (UVC) and sealed in a DNA clean lab in order to remove all possible contaminants of DNA. We collected surface water from the edge of the river and filtered each volume on site. For each of the sampled volumes we sequentially filtered batches of 250 mL of water onto a single 25 mm 0.70-µm glass fiber filter (GF/F, Whatman International Ltd., Maidstone, UK). The total number of filters used for each volume class ranged from one to eight filters. The filters were housed in a 25 mm filter case (Swinnex, EMD Millipore Co., Billerica, Massachusetts, USA) that was attached to a disposable 50-mL syringe. For each volume class we used the same filter housing and the same syringe and changed only the filter as necessary to process the total volume. After filtration, we transferred the filters into individual 1.5 mL tubes containing tissue lysis buffer (100 mM Tris-HCL pH 8.0, 5 mM EDTA pH 8.0, 0.2% SDS, 200 mM NaCl₂) using tweezers that were decontaminated with 10% household beach between volume replicates and rinsed with ethanol. The tubes were immediately stored on ice. The time between the first and last sample was approximately 2.5 h during field filtration. Samples were immediately transported to the laboratory with a maximum travel time of 30 minutes. Additionally, we created two negative filtration controls, which consisted of decontaminated Milli-Q® water treated with UVC light and sealed in a DNA free laboratory. We brought this water to the field and filtered 2000 mL before environmental samples were taken on each sampling day, resulting in 4 negative filtration controls. Upon return to the laboratory we immediately began the extraction as described below.

DNA extraction and species eDNA amplification

In the laboratory we performed a modified cell lysis phenol-chloroform-isoamyl extraction on each single filter as this has been shown as an effective extraction method for eDNA from glass fiber filters.¹⁰ We added for each set of extraction a negative control (further called negative extraction control). All eDNA extractions of each target volume were subsequently pooled, such that eDNA was resuspended across all filtration volumes in a total volume of 100 μ L. For example, the two filters used for the 500 mL volume were each resuspended in 50 μ L and then pooled to equal the total extracted DNA volume of 100 μ L. All pooled DNA extractions were cleaned with the OneStepTM PCR Inhibitor Removal Kit (Zymo Research, Irvine, California, USA) according to the provided protocol. To counteract for volume loss during clean-up we added 50 μ L of AE buffer (this is the elution buffer from DNeasy Blood & Tissue kit delivered from Qiagen GmbH, Hilden Germany) to each cleaned DNA extraction to allow for testing of multiple species with the same extraction. Cleaning eDNA with this additional step has been shown to be effective for removal of PCR inhibition of riverine samples of environmental DNA¹³ and we did not want PCR inhibition to additionally confound the detection rate. All pooled and cleaned extractions were quantified by using the Qubit (1.0) fluorometer following the recommended protocol for the high sensitivity (HS) assay for dsDNA (Life Technologies, Carlsbad, CA, USA) and can be consulted in Figure S1.

We conducted standard PCR, because the primers used in this study were designed for standard PCR for a previous study,²¹ and thus allows comparability. Additionally we wanted to use a method that seems commonly available to federal offices. We then performed eight PCR replicates, where each replicate screened 2 μ L with concentrations ranging from 0.153 to 3.73 μ g/mL of extracted DNA, equaling 16 μ L screened for each of the following three target species: *Ancylus fluviatilis* (Mollusca), *Baetis buceratus* (Ephemeroptera) and *Gammarus pulex* (Amphipoda). Based on previous studies, we know that these species belong to the regional

species pool at this site from long-term monitoring data (1995 to 2012) provided by the Canton of Zurich and our own sampling.²²⁻²⁴ All negative filtration controls and all negative extraction controls were tested individually in eight PCR replicates for the presence of each species. We used primer probes previously designed and tested for eDNA detection of these species at the study site.²¹ For each PCR run we added a negative PCR control by adding molecular grade DNA free water (Sigma-Aldrich, Co. LLC. St. Lewis, MO, USA) as template and a positive PCR control (using tissue extracted DNA from the target species as PCR template). The PCR components and thermocycling temperature were performed exactly as described in Mächler *et al.*²¹ and can be found in the supplementary information of this paper (S1). All PCR products were visualized by electrophoresis on a 1.4% agarose gel stained with peqGreen (VWR International, Radnor, Pennsylvania, USA).

From each volume of water we confirmed at least one positive PCR reaction by using Sanger sequencing. We cleaned the PCR product with Exo I Nuclease (EXO I) and Shrimp Alkaline Phosphatase (SAP) (Thermo Fischer Scientific, Waltham, Maryland, USA) as described in Mächler *et al.*²¹. Sequencing was performed in both directions with BigDye® Terminator (version 3.1) system on an ABI 3730xl. The software Sequencher® version 4.9 (Gene Codes, Ann Arbor, Michigan, USA) was used to align, edit and compare our sequences with previous eDNA sequences obtained from this site in a previous year and tissue derived sequences.²¹ We used the same criteria for a positive detection (band present on a gel and sequence confirmation for each experimental volume) and rigorous laboratory precautions as described in Mächler *et al.*²¹ by creating, in addition to our negative filtration controls, negative controls for extraction and PCR. In total we screened four negative filtration, two negative extraction and twelve negative PCR controls for potential contamination.

Analysis

We analyzed the detection rate of each individual species with generalized linear models (GLMM). Volume of water was used as predictor variable and detection rate as a binary response variable consisting of the number of positive and negative detections out of the eight PCR replications. We used the two replicates per volume on a single day as a random effect, nested within the sampling day for *G. pulex* and *B. buceratus*. For *A. fluviatilis*, we had only one sampling day, due to a contamination on the first day.

We tested how the uncertainty in detection rates changed as a result of increasing the amount of extracted DNA screened for each species, using a resampling approach (bootstrap approach) of the individual assessments. We sampled 10,000 outcomes in detection rates from our data when screening 2 to 16 μ L of DNA. As a measure of uncertainty, we subsequently calculated the median range (absolute difference between minimum and maximum) in detection rates over all outcomes for a given sample volume and species (data from the two sampling days were pooled). When uncertainty is one, detection rates can be any value between 0 and 1, that is, the estimate is uninformative; when uncertainty is zero, then the detection rate is the same for all outcomes and is maximally informative. All statistical analysis were done in R version 2.15.3 (R Development Core Team 2014)²⁵, and the package “lme4”²⁶.

Results

All three species were detected at the sampling site by the use of eDNA in all volumes and on both sampling dates (Fig. 1). We found a positive, significant relationship between sampling

volume and detection rate for *G. pulex* ($p < 0.05$, Table 2). For the other two species there was no significant relationship (*A. fluviatilis* $p = 0.78$, *B. buceratus* $p = 0.72$, Table 2).

We showed for all three species that the uncertainty in the detection rate decreases when increasing the volume of extracted eDNA screened (Fig. 2). The uncertainty decreased differently between the three species, however, all species reached zero uncertainty (i.e., detection rate never changes between outcomes) when screening at least 14 μL of extracted DNA.

We detected a contamination (a confirmed amplicon of the targeted species) in one negative filtration control for *A. fluviatilis* on the first sampling day. From the total 16 μL of extracted DNA, the negative filtration control screened showed a positive amplicon in three PCR replicates (i.e., 6 μL of screened DNA). Attempts to measure DNA concentration from this negative filtration control failed as DNA concentrations were too low (limit of detection reported for Qubit high sensitivity assays $< 0.001 \mu\text{g/mL}$). All other negative controls (filtration, extraction and PCR) were blank. The three positive amplifications of this negative filtration control happened within replicates in the same PCR set up, thus we are confident that the contamination happened during the preparation of the PCR reactions, and was not a contamination from the field.

Unfortunately, due to the testing of the extractions for so many replicates and species, we ran out of the extracted DNA before being able to repeat this PCR. In order to be most conservative, we excluded the data for this species from the first day from all analyses.

Discussion

Comparing the effect of sampled water volume and an eDNA based detection across three macroinvertebrate species, we surprisingly found that only one species (*Gammarus pulex*) had a positive relationship with an increased detection rate when more water was sampled. The detection rate of the other two species did not correlate with sampled water volume. Based on the

results of a previous study using the same primers²¹, we calculated the detection probability (i.e., the detection with eDNA divided by the proven presence with traditional monitoring method) in river systems using the same primers. This detection probability was 0.83 for *A. fluviatilis*, 1 for *B. buceratus* and 0.71 for *G. pulex*. Thus, as the latter species seems to have in general a lower detectability, it may be the one most affected by the total amount of water volume sampled. Indeed, we found a dependence of water volume and detection rate for this species at the lower volumes, but the effect leveled off above 1 L, and then saturated at the species' overall detection probability (i.e., detection rate at 1 L = 0.72 ± 0.18 and at 2 L = 0.75 ± 0.15). Surprisingly we did not find a positive relationship between water volume and detection rate for all three species. We speculate that such a relationship exists also for the other two species. We think that we did not reach the lower limit of the detection where this relationship exists for *A. fluviatilis* and *B. buceratus*, while for *G. pulex* we were within the water volume range where the saturation occurs. Our results thus indicate that detection rates may vary by species and volume, which should be considered when designing targeted eDNA detection tool.

All species' DNA was detected in at least one PCR replicate at the smallest volume of 250 mL. The detection rate had a lower uncertainty at higher volumes of DNA extraction screened compared to lower volumes (Fig. 2) and indicates that screening more of the DNA extraction allows for a smaller uncertainty in detection rate for a species with the used protocols across any volume. The volume of extraction we screened is similar to other studies using filtration as DNA capturing approach²⁷⁻²⁹, and these studies sampled similar volumes (0.2 to 1 L) of water (Table 1). Even when we sampled two liters of water, we found in none of the three species studied a positive signal in 100% of the PCR replications. PCR is a stochastic process and, subsequently, PCR replication (e.g., the volume of screened eDNA) cannot be neglected even when sampling larger volumes. In Fig. 2 we illustrate the importance of screening enough volume of extracted

DNA to decrease the uncertainty in the outcome of the detection. Although this is intuitive, it is an important aspect to keep in mind when creating a protocol for species detection with eDNA. We recommend testing volume dependence for each species that should be detected with eDNA, so that eDNA protocols are optimized with respect to sampling volume and screened volume of eDNA extractions. When resources are limited, we suggest maximizing the volume of extracted eDNA which is screened and not necessarily to maximize the water volume sampled. PCR replication is more cost effective than the filtration and extraction of larger volumes of water due to two reasons: first, at our study site we were not able to filter more than 250 mL on one single glass fiber filter because of free floating particles that clog the filters, even though we conducted our work in a river which is known to have an overall low level of particle and sediment load. For filtering volumes of two liters, we needed up to eight filters. The usage of multiple filters for the same sample is a time consuming and costly step, as each filter costs about 1 USD. Second, the extraction will be even more time intense, as each of the filters needs to be extracted separately due to limitations of tube sizes and limited ability to handle large volumes (greater than 2 L) for most standard molecular genetic laboratory centrifuges. One may need to validate these results across various river systems, in order to adjust methods for different eDNA quality and degradation, primer sensitivity and sediment loads. However, the conditions in the river Glatt are likely valid for rivers and streams in human-modified temperate landscapes with a mixed land-use of urban areas, forests and agricultural land-use.

We especially want to highlight the relevance of negative controls. We performed negative controls during three steps in the handling process: for filtration, extraction and PCR. Contaminations can occur, especially while filtering in the field, but through a good study design it is possible to track down the source of contaminations. We suggest that it is relevant to not only screen a certain volume of extracted DNA but also to screen an adequate volume for the

negative filtration controls. In our study we had contamination for one species in one filtration control for the first day. As all the positive amplicon showed up in one set of PCR it is very likely that that our contamination happened during the preparation of the PCR. However, as we cannot rule out field contamination, we were stringent and have excluded all detections of this species from the first day. The implication of excluding dubious data is especially important for environmental DNA work where low amounts of DNA are handled and small contaminations can lead to false positives. Given our findings regarding the effect of volume (either from the environment or from the extraction) on uncertainty in detection rates we especially encourage eDNA researchers to report the number and volume of negative controls that are screened to ensure accountability. When this tool becomes used in controversial cases it will be paramount to uphold the same practices and standards we apply to samples also to negative controls for proper inference.

Optimization of single species detection through eDNA may depend on the specific species and environmental settings; however, some critical considerations and guidelines can still be inferred from our results. Overall we find that there may be different factors contributing to the successful detection of species using eDNA. First, we conclude based on first principles that at a given concentration of eDNA molecules in the environment, there must be a lower volume threshold at which detection becomes less likely, while the detection rate saturates at higher volume due to other factors (e.g., primer performance¹² and competition of target DNA vs. non-target DNA during PCR²¹). Our data, however, suggest that this level is below 250 mL of water, at least for the three species tested here and under this molecular workflow. While increasing sampling volume may indeed be beneficial in reducing false negative detections (i.e., reducing uncertainty in detection rate in our study) for some species, increasing the volume of extracted DNA screened and primer performance may become more important due to a lower detection

rate for smaller environmental sampling volumes. Second, when comparing methods, one not only needs to quantify detection thresholds and rates of false negatives for the eDNA method, but also the method it is compared with. While the issue of both false positives and false negative detections has been raised for eDNA-approaches^{2,9}, it is often ignored for traditional methods, where a perfect sampling is often implicitly assumed (but see^{8,30,31} on extensive discussion on this topic). Based on our results we can give a rough recommendation that sampling at least 1 L and screening a minimum of 14 µL of total extracted DNA should reduce false negative detections particularly for macroinvertebrates in freshwater and potentially for other macro-species surveillance. The recommendations should be taken with some precautions as results might be changing even within similar species and environments. We caution researchers to carefully plan sampling designs with regard to volume and encourage pilot testing these parameters in order to maximize potential detection rates for other systems.

Acknowledgments

We thank Pravin Ganesanandamoorthy and Leila Rubin for help during the laboratory work. We thank Emanuel A. Fronhofer and three anonymous reviewers for discussion and helpful comments. Genetic data were analyzed in collaboration with the Genetic Diversity Center (GDC), ETH Zurich. Funding is from the Swiss National Science Foundation Grant PP00P3_150698 (to F.A.) and Eawag Discretionary Funds (to F.A. and K.D.).

References

- 340 (1) Lawson Handley, L. How will the ‘molecular revolution’ contribute to biological
341 recording? *Biological Journal of the Linnean Society*. **2015**, *115*, (3), 750-766.
- 342 (2) Thomsen, P. F.; Willerslev, E. Environmental DNA—an emerging tool in conservation for
343 monitoring past and present biodiversity. *Biological Conservation*. **2015**, *183*, 4-18.
- 344 (3) Goldberg, C. S.; Strickler, K. M.; Pilliod, D. S. Moving environmental DNA methods
345 from concept to practice for monitoring aquatic macroorganisms. *Biological Conservation*. **2015**,
346 *183*, 1-3.
- 347 (4) Kelly, R. P.; Port, J. A.; Yamahara, K. M.; Martone, R. G.; Lowell, N.; Thomsen, P. F.;
348 Mach, M. E.; Bennett, M.; Prahler, E.; Caldwell, M. R. Harnessing DNA to improve
349 environmental management. *Science*. **2014**, *344*, (6191), 1455-1456.
- 350 (5) Roussel, J. M.; Paillisson, J. M.; Tréguier, A.; Petit, E. The downside of eDNA as a
351 survey tool in water bodies. *Journal of Applied Ecology*. **2015**, *52*, (4), 823-826.
- 352 (6) Rees, H. C.; Maddison, B. C.; Middleditch, D. J.; Patmore, J. R.; Gough, K. C. The
353 detection of aquatic animal species using environmental DNA—a review of eDNA as a survey
354 tool in ecology. *Journal of Applied Ecology*. **2014**, *51*, (5), 1450-1459.
- 355 (7) Jerde, C. L.; Mahon, A. R. Improving confidence in environmental DNA species
356 detection. *Molecular Ecology Resources*. **2015**, *15*, (3), 461-463.
- 357 (8) Haase, P.; Murray-Bligh, J.; Lohse, S.; Pauls, S.; Sundermann, A.; Gunn, R.; Clarke, R.,
358 Assessing the impact of errors in sorting and identifying macroinvertebrate samples. In *The*
359 *Ecological Status of European Rivers: Evaluation and Intercalibration of Assessment Methods*,
360 Springer: 2006; pp 505-521.
- 361 (9) Schmidt, B. R.; Kéry, M.; Ursenbacher, S.; Hyman, O. J.; Collins, J. P. Site occupancy
362 models in the analysis of environmental DNA presence/absence surveys: A case study of an
363 emerging amphibian pathogen. *Methods in Ecology and Evolution*. **2013**, *4*, (7), 646-653.

- 364 (10) Deiner, K.; Walser, J.-C.; Mächler, E.; Altermatt, F. Choice of capture and extraction
365 methods affect detection of freshwater biodiversity from environmental DNA. *Biological*
366 *Conservation*. **2015**, *183*, 53-63.
- 367 (11) Eichmiller, J. J.; Miller, L. M.; Sorensen, P. W. Optimizing techniques to capture and
368 extract environmental DNA for detection and quantification of fish. *Molecular ecology*
369 *resources*. **2015**.
- 370 (12) Wilcox, T.; McKelvey, K.; Young, M.; Jane, S.; Lowe, W.; Willson, R. C. Robust
371 Detection of Rare Species Using Environmental DNA: The Importance of Primer Specificity.
372 *PLoS One*. **2013**, *8*, (3), e59520.
- 373 (13) McKee, A. M.; Spear, S. F.; Pierson, T. W. The effect of dilution and the use of a post-
374 extraction nucleic acid purification column on the accuracy, precision, and inhibition of
375 environmental DNA samples. *Biological Conservation*. **2015**, *183*, 70-76.
- 376 (14) Schnell, I. B.; Bohmann, K.; Gilbert, M. T. P. Tag jumps illuminated—reducing sequence-
377 to-sample misidentifications in metabarcoding studies. *Molecular Ecology Resources*. **2015**, *15*,
378 (6), 1289-303.
- 379 (15) Esling, P.; Lejzerowicz, F.; Pawlowski, J. Accurate multiplexing and filtering for high-
380 throughput amplicon-sequencing. *Nucleic Acids Research*. **2015**, *43*, (5), gkv107.
- 381 (16) Gotelli, N. J. A primer of ecology. Sunderland, Massachusetts. In Sinauer Associates, Inc:
382 2001.
- 383 (17) Turner, C. R.; Uy, K. L.; Everhart, R. C. Fish environmental DNA is more concentrated in
384 aquatic sediments than surface water. *Biological Conservation*. **2015**, *183*, 93-102.
- 385 (18) Schill, W. B.; Mathes, M. V. Real-time PCR detection and quantification of nine potential
386 sources of fecal contamination by analysis of mitochondrial cytochrome b targets. *Environmental*
387 *science & technology*. **2008**, *42*, (14), 5229-5234.

- 388 (19) Goldberg, C. S.; Pilliod, D. S.; Arkle, R. S.; Waits, L. P. Molecular detection of
389 vertebrates in stream water: a demonstration using Rocky Mountain tailed frogs and Idaho giant
390 salamanders. *PLoS One*. **2011**, 6, (7), e22746.
- 391 (20) Deiner, K.; Altermatt, F. Transport distance of invertebrate environmental DNA in a
392 natural river. *PloS one*. **2014**, 9, (2), e88786.
- 393 (21) Mächler, E.; Deiner, K.; Steinmann, P.; Altermatt, F. Utility of environmental DNA for
394 monitoring rare and indicator macroinvertebrate species. *Freshwater Science*. **2014**, 33, (4),
395 1174-1183.
- 396 (22) AWEL *Zürcher Gewässer 2012: Entwicklung - Zustand - Ausblick*.; AWEL Amt für
397 Abfall, Wasser, Energie und Luft, Zürich.: 2012.
- 398 (23) Altermatt, F.; Seymour, M.; Martinez, N. River network properties shape α -diversity and
399 community similarity patterns of aquatic insect communities across major drainage basins.
400 *Journal of Biogeography*. **2013**, 40, (12), 2249-2260.
- 401 (24) Altermatt, F.; Alther, R.; Fišer, C.; Jokela, J.; Konec, M.; Kury, D.; Mächler, E.; Stucki,
402 P.; Westram, A. M. Diversity and Distribution of Freshwater Amphipod Species in Switzerland
403 (Crustacea: Amphipoda). *PLoS One*. **2014**, 9, (10), e110328.
- 404 (25) Team, R. C. R: A language and environment for statistical computing. R Foundation for
405 Statistical Computing, Vienna, Austria, 2012. In ISBN 3-900051-07-0: 2014.
- 406 (26) Bates, D.; Mächler, M.; Bolker, B.; Walker, S. Fitting linear mixed-effects models using
407 lme4. *arXiv preprint arXiv:1406.5823*. **2014**.
- 408 (27) Eichmiller, J. J.; Bajer, P. G.; Sorensen, P. W. The relationship between the distribution of
409 common carp and their environmental DNA in a small lake. *PLoS One*. **2014**, 9, (11), e112611.

- 410 (28) Spear, S. F.; Groves, J. D.; Williams, L. A.; Waits, L. P. Using environmental DNA
411 methods to improve detectability in a hellbender (*Cryptobranchus alleganiensis*) monitoring
412 program. *Biological Conservation*. **2015**, *183*, 38-45.
- 413 (29) Takahara, T.; Minamoto, T.; Doi, H. Using environmental DNA to estimate the
414 distribution of an invasive fish species in ponds. *PLoS One*. **2013**, *8*, (2), e56584.
- 415 (30) Kery, M. Inferring the absence of a species: a case study of snakes. *The Journal of*
416 *wildlife management*. **2002**, *66*, (2), 330-338.
- 417 (31) Kery, M.; Plattner, M. Species richness estimation and determinants of species
418 detectability in butterfly monitoring programmes. *Ecological Entomology*. **2007**, *32*, (1), 53-61.
- 419 (32) Kortbaoui, R.; Locas, A.; Imbeau, M.; Payment, P.; Villemur, R. Universal mitochondrial
420 PCR combined with species-specific dot-blot assay as a source-tracking method of human,
421 bovine, chicken, ovine, and porcine in fecal-contaminated surface water. *Water research*. **2009**,
422 *43*, (7), 2002-2010.
- 423 (33) Jerde, C. L.; Mahon, A. R.; Chadderton, W. L.; Lodge, D. M. "Sight-unseen" detection of
424 rare aquatic species using environmental DNA. *Conservation Letters*. **2011**, *4*, (2), 150-157.
- 425 (34) Minamoto, T.; Yamanaka, H.; Takahara, T.; Honjo, M. N.; Kawabata, Z. i. Surveillance
426 of fish species composition using environmental DNA. *Limnology*. **2012**, *13*, (2), 193-197.
- 427 (35) Olson, Z. H.; Briggler, J. T.; Williams, R. N. An eDNA approach to detect eastern
428 hellbenders (*Cryptobranchus a. alleganiensis*) using samples of water. *Wildlife Research*. **2012**,
429 *39*, (7), 629-636.
- 430 (36) Takahara, T.; Minamoto, T.; Yamanaka, H.; Doi, H.; Kawabata, Z. i. Estimation of fish
431 biomass using environmental DNA. *PloS one*. **2012**, *7*, (4), e35868.

432 (37) Tambalo, D. D.; Boa, T.; Liljebjelke, K.; Yost, C. K. Evaluation of two quantitative PCR
433 assays using Bacteroidales and mitochondrial DNA markers for tracking dog fecal contamination
434 in waterbodies. *Journal of microbiological methods*. **2012**, *91*, (3), 459-467.

435 (38) Thomsen, P. F.; Kielgast, J.; Iversen, L. L.; Møller, P. R.; Rasmussen, M.; Willerslev, E.
436 Detection of a diverse marine fish fauna using environmental DNA from seawater samples. *PLoS*
437 *One*. **2012**, *7*, (8), e41732.

438 (39) Goldberg, C. S.; Sepulveda, A.; Ray, A.; Baumgardt, J.; Waits, L. P. Environmental DNA
439 as a new method for early detection of New Zealand mudsnails (*Potamopyrgus antipodarum*).
440 *Freshwater Science*. **2013**, *32*, (3), 792-800.

441 (40) Jerde, C. L.; Chadderton, W. L.; Mahon, A. R.; Renshaw, M. A.; Corush, J.; Budny, M.
442 L.; Mysorekar, S.; Lodge, D. M. Detection of Asian carp DNA as part of a Great Lakes basin-
443 wide surveillance program. *Canadian Journal of Fisheries and Aquatic Sciences*. **2013**, *70*,
444 (999), 1-5.

445 (41) Mahon, A. R.; Jerde, C. L.; Galaska, M.; Bergner, J. L.; Chadderton, W. L.; Lodge, D.
446 M.; Hunter, M. E.; Nico, L. G. Validation of eDNA surveillance sensitivity for detection of Asian
447 carps in controlled and field experiments. *PLoS One*. **2013**, *8*, (3), e58316.

448 (42) Pilliod, D. S.; Goldberg, C. S.; Arkle, R. S.; Waits, L. P. Estimating Occupancy and
449 Abundance of Stream Amphibians Using Environmental DNA from Filtered Water Samples.
450 *Canadian Journal of Fisheries and Aquatic Sciences*. **2013**, *70*, (ja), 1123-1130.

451 (43) Vuong, N.-M.; Villemur, R.; Payment, P.; Brousseau, R.; Topp, E.; Masson, L. Fecal
452 source tracking in water using a mitochondrial DNA microarray. *Water research*. **2013**, *47*, (1),
453 16-30.

- 454 (44) Jane, S. F.; Wilcox, T. M.; McKelvey, K. S.; Young, M. K.; Schwartz, M. K.; Lowe, W.
455 H.; Letcher, B. H.; Whiteley, A. R. Distance, flow and PCR inhibition: eDNA dynamics in two
456 headwater streams. *Molecular Ecology Resources*. **2015**, *15*, (1), 216-227.
- 457 (45) Keskin, E. Detection of invasive freshwater fish species using environmental DNA
458 survey. *Biochemical Systematics and Ecology*. **2014**, *56*, 68-74.
- 459 (46) Pilliod, D. S.; Goldberg, C. S.; Arkle, R. S.; Waits, L. P. Factors influencing detection of
460 eDNA from a stream-dwelling amphibian. *Molecular Ecology Resources*. **2014**, *14*, (1), 109-116.
- 461 (47) Amberg, J. J.; McCalla, S. G.; Monroe, E.; Lance, R.; Baerwaldt, K.; Gaikowski, M. P.
462 Improving efficiency and reliability of environmental DNA analysis for silver carp. *Journal of*
463 *Great Lakes Research*. **2015**, *41*, (2), 367-373.
- 464 (48) Fukumoto, S.; Ushimaru, A.; Minamoto, T. A basin-scale application of environmental
465 DNA assessment for rare endemic species and closely related exotic species in rivers: a case
466 study of giant salamanders in Japan. *Journal of Applied Ecology*. **2015**, *52*, (2), 358-365.
- 467 (49) Hunter, M. E.; Oyler-McCance, S. J.; Dorazio, R. M.; Fike, J. A.; Smith, B. J.; Hunter, C.
468 T.; Reed, R. N.; Hart, K. M. Environmental DNA (eDNA) sampling improves occurrence and
469 detection estimates of invasive Burmese pythons. *PLoS One*. **2015**, *10*, (4), e0121655.
- 470 (50) Janosik, A. M.; Johnston, C. E. Environmental DNA as an effective tool for detection of
471 imperiled fishes. *Environmental Biology of Fishes*. **2015**, *98*, (8), 1-5.
- 472 (51) Laramie, M. B.; Pilliod, D. S.; Goldberg, C. S. Characterizing the distribution of an
473 endangered salmonid using environmental DNA analysis. *Biological Conservation*. **2015**, *183*,
474 29-37.

Figures

Fig. 1: Detection rate, as proportion of positive amplifications across eight PCR replicates for each volume replication, relative to water volume sampled. Error bars stand for standard errors across four samples (*B. buceratus* and *G. pulex*) and two samples (*A. fluviatilis*) respectively.

482

483 **Fig. 2:** Uncertainty in the detection rate relative to screened volume of extracted environmental
 484 DNA. When uncertainty is one, detection rates and thus the outcome vary between 0 and 1; when
 485 uncertainty is zero, detection rate is completely consistent between outcomes. The uncertainty of
 486 detection rate is decreasing with increasing volume of screened DNA for all three species and
 487 reaches zero at 14 μL of screened volume of eDNA.

488 **Tables**

489 **Table 1:** Overview on published methods and quantities used for filtering eDNA from macrofauna in natural aquatic systems as well as
 490 used PCR conditions. If dilution series were done the given volume for PCR template is the maximum. Samples per site are only reported
 491 if they are taken at the same sampling site, several samples that were taken within the same sampling system were not reported.

Authors	Year	Citation	Filter material	Filter pore size (µm)	Sampled water volume (mL)	Type of PCR	Template volume used for PCR (µL)	Total PCR volume (µL)	Number of PCR replications	Samples per site (e.g. volume replication)	Total PCR replication per site	Volume of extraction screened
Schill & Mathes	2008	¹⁸	polyethylensulfone	0.22	100-1000	real time	2.5	12.5	6	1	6	15
Kortbaoui et al.	2009	³²	nitrocellulose	0.22 or 0.45	150-300	singleplex	NA	50	1	NA	NA	NA
						nested	2 ⁺	50	1	NA	NA	NA
Goldberg et al.	2011	¹⁹	cellulose nitrate*	0.45	5000-10000	standard	1	10	1	1 (10 L), 2 (5L)	1-2	1-2
			cellulose nitrate*	0.45	5000	multiplex	1	7	6		6	6
Jerde et al.	2011	³³	glass fiber	1.5	2000	standard	NA	25	8	NA	NA	NA
Minamoto et al.	2012	³⁴	polycarbon	3	2000	standard	4	25	1	1	1	4
Olson et al.	2012	³⁵	glass fiber	1.5	8000	standard	1	10	10	1	10	10
Takahara et al.	2012	³⁶	cellulose acetate	3	2000	quantitative	5	20	3	1	3	9
Tambalo et al.	2012	³⁷	NA	0.45	500	quantitative	2-4	25	2	1	2	4-8
Thomsen et al.	2012	³⁸	nylon	0.45	500	standard	2	25	8	3	24	48
Goldberg et al.	2013	³⁹	cellulose nitrate	0.45	4000	quantitative	2.5	10	3-9	3	9-27	22.5-67.5
Jerde et al.	2013	⁴⁰	glass fiber	1.5	2000	standard	NA	25	8	NA	NA	NA
Mahon et al.	2013	⁴¹	glass fiber	1.5	2000	standard	NA	25	8	NA	NA	NA
Pilliod et al.	2013	⁴²	cellulose nitrate*	0.45	1000	quantitative	2	10	3-6	1	3-6	6-12

Authors	Year	Citation	Filter material	Filter pore size (µm)	Sampled water volume (mL)	Type of PCR	Template volume used for PCR (µL)	Total PCR volume (µL)	Number of PCR replications	Samples per site (e.g. volume replication)	Total PCR replication per site	Volume of extraction screened
Schmidt et al.	2013	⁹	polyethylensulfone	NA	600	quantitative	NA	NA	2	1	2	NA
Takahara et al.	2013	²⁹	cellulose acetate	3	1000	real time	2	20	8	1	8	16
Vuong et al.	2013	⁴³	NA	0.45	300	standard	NA	25	NA	NA	NA	NA
						nested	1 ⁺	50	NA	NA	NA	NA
						quantitative	5	20	3	NA	NA	NA
Wilcox et al.	2013	¹²	glass fiber	1.5	6000	quantitative	4	20	26-32	1	26-32	104-128
Eichmiller et al.	2014	²⁷	glass fiber	1.5	200	quantitative	5	25	3	1	3	15
Jane et al.	2014	⁴⁴	glass fiber	1.5	6000	quantitative	4	20	3	1	3	12
Keskin et al.	2014	⁴⁵	polyethylensulfone	0.22	2000	standard	5	25	3-6	3	9-18	45-90
Mächler et al.	2014	²¹	glass fiber	0.7	900	standard	2	15	8	1	8	16
Pilliod et al.	2014	⁴⁶	cellulose nitrate*	0.45	2000	quantitative	2	10	3	1	3-6	6-12
Amberg et al.	2015	⁴⁷	glass fiber	1.5	2000	standard	NA	25	8	NA	NA	NA
			glass fiber	1.5	2000	quantitative	1	25	8	NA	NA	8
Fukumoto et al.	2015	⁴⁸	glass fiber	0.7	4000	real time	2	20	4	1	4	8
Hunter et al.	2015	⁴⁹	cellulose nitrate	0.45	250-1000	quantitative	NA	20.4	3	1 - 3	9 - 27	NA
Janosik & Johnston	2015	⁵⁰	glass fiber	1.5	2000	standard	1	25	3	NA	NA	3
Laramie et al.	2015	⁵¹	cellulose nitrate	0.45	1000	quantitative	3	15	3	3	9	27
McKee et al.	2015	¹³	cellulose nitrate	0.45	250-1000	quantitative	3.75	15	3	1	3	11.25
Spear et al.	2015	²⁸	cellulose nitrate*	0.45	1000	quantitative	3	15	3-6	1	1	9-18

492 * Used only half filter to extract DNA

493 + Template of first PCR product

494 NA Not available from publication

495 **Table 2:** GLMM results on the effect of volume for the detection of each species.

	Coefficient	Standard error	Z-value	p-value
A) <i>Ancylus fluviatilis</i>				
Intercept	0.0973	0.4309	0.23	0.82
Volume	0.0001	0.0004	0.28	0.78
B) <i>Baetis buceratus</i>				
Intercept	1.5214	0.7362	2.07	0.04
Volume	-0.0001	0.0003	-0.36	0.72
C) <i>Gammarus pulex</i>				
Intercept	-1.1762	1.1871	-0.99	0.32
Volume	0.0016	0.0004	4.34	< 0.01

496

250 mL 500 mL 1000 mL 1500 mL 2000 mL