

# Supplementary Material- Crop productivity changes in 1.5°C and 2°C worlds under climate sensitivity uncertainty

Carl-Friedrich Schleussner<sup>1,2</sup>, Delphine Deryng<sup>1,3</sup>, Christoph Müller<sup>2</sup>, Joshua Elliott<sup>3,4</sup>, Fahad Saeed<sup>1</sup>, Christian Folberth<sup>5</sup>, Wenfeng Liu<sup>6</sup>, Xuhui Wang<sup>7,8</sup>, Thomas A. M. Pugh<sup>9,10</sup>, Wim Thiery<sup>11,12</sup>, Sonia I. Seneviratne<sup>11</sup>, Joeri Rogelj<sup>5</sup>

<sup>1</sup> Climate Analytics, 10969 Berlin, Germany

<sup>2</sup> Potsdam Institute for Climate Impact Research, 14473 Potsdam, Germany

<sup>3</sup> Columbia University Center for Climate Systems Research, New York, NY 10025, USA

<sup>4</sup> University of Chicago and ANL Computation Institute, Chicago, IL 60637, USA

<sup>5</sup> International Institute for Applied Systems Analysis, Ecosystem Services and Management Program, 2361 Laxenburg, Austria

<sup>6</sup> Eawag, Swiss Federal Institute of Aquatic Science and Technology, 8600 Duebendorf, Switzerland

<sup>7</sup> Laboratoire des Sciences du Climat et de l'Environnement. CEA CNRS UVSQ Orme des Merisiers

<sup>8</sup> Sino-French Institute of Earth System Sciences, Peking University, 100871 Beijing, China

<sup>9</sup> School of Geography, Earth & Environmental Sciences and Birmingham Institute of Forest Research, University of Birmingham, B15 2TT, UK

<sup>10</sup> Karlsruhe Institute of Technology, IMK-IFU, 82467 Garmisch-Partenkirchen, Germany

<sup>11</sup> ETH Zurich, 8092 Zürich, Switzerland

<sup>12</sup> Vrije Universiteit Brussel, 1050 Brussels, Belgium

Table S1 | Changes in model setups from the detailed descriptions provided in (Müller et al 2017).

GGCM	Changes in GGCM setup	References
<b>LPJmL</b>	-	
<b>LPJ-Guess</b>	Includes nitrogen dynamics and limitations	(Olin <i>et al</i> 2015b, 2015a)
<b>GEPIEC</b>	Use of static (annually re-initialized) soil profile rather than dynamic tracking of soil processes Soil data were pre-conditioned in a dynamic spin-up run Spin-up years changed from 30 to 8.	-
<b>PEPIC</b>	Spin-up years changed from 20 to 8.	(Liu <i>et al</i> 2016b, 2016a)
<b>ORCHIDEE-crop</b>	-	
<b>Pegasus</b>	-	
<b>CLM-crop-ETH</b>	Crop area abundance prescribed according to land cover maps from the Land Use Harmonisation Project Phase II (LUH2, Lawrence et al., 2016), rather than simulating 'all crops everywhere'. Winter wheat and spring wheat were simulated separately and aggregated into a single wheat field by (i) identifying the pixels with the highest winter, respectively spring wheat yield in the CAM4 All-Hist simulation (rainfed and irrigated crops separately), and (ii) subsequently applying this mask to all simulation output.	(Lawrence <i>et al</i> 2011, 2016)

Table S2: Overview of GGCM data availability. Not all GGCMs have modelled all GCMs based on the bias corrected HAPPI model intercomparison climate input or all crops. The ensemble members from the HAPPI multi-ensemble simulations as well as the years per ensemble member (historical period or future warming) are given. The short time period leads to anomalies for annual harvest for some GGCMs, which leads to exclusion of the first or last year of each period. The resulting total number of model years per warming level is also given. The column CO<sub>2</sub> experiment indicates the models that provided data for all CO<sub>2</sub> levels investigated.

	GCM input				Crop				CO <sub>2</sub> Exp	Ensemble members	Years per run	Years per warming level and GCM
	MIROC5	NorESM1	CAM4	ECHAM6	Wheat	Maize	Soy	Rice				
<b>CLM-crop-ETH</b>	x	x	x		x	x	x		x	1-5	1-10	50
<b>LPJmL</b>	x	x	x	x	x	x	x	x	x	1-5	1-10	50
<b>LPJ-Guess</b>	x	x	x		x	x			x	1-5	2-10	45
<b>GEPIC</b>	no rice, wheat,	x	x	x	x	x	x	x	x	1-5	1-9	45
<b>PEPIC</b>		x	x		x	x		x	x	1-5	1-10	50
<b>ORCHIDEE-crop</b>	x	x			x	x		x		1-5	1-9	45
<b>Pegasus</b>	x	x		x	x	x	x		x	1-5	2-10	45

Table S3 |Global mean temperature differences between the recent past and the 1.5°C and 2°C future periods in the ensemble of HAPPI GCMs.

	Warming relative to the 2006-2015 period [°C]				Ensemble Average
	MIROC5	ECHAM6	NorESM1	CAM4	
Plus 1.5°C	0.68	0.64	0.71	0.66	<b>0.67</b>
Plus 2°C	1.14	1.11	1.12	1.13	<b>1.12</b>
Plus 2°C-1.5°C	0.46	0.47	0.41	0.47	<b>0.45</b>




Figure S1: Probabilities of staying below certain warming levels in the transient climate response following IPCC AR5 estimates for the TCR distribution (see Methods).




Figure S2: As Figure 1, but resolved for different GCMs. See table S1 for data availability of GCM-GGCM pairs.

Table S4: Changes in crop productivity between the 1.5°C and 2°C periods relative to 2006-2015 (in percent). Crop production is aggregated over SREX regions, the Tropics (between 30°S/°N) and global. Regions with less than 0.1% contribution to global crop production according to the MIRCA2000 dataset are masked out. Numbers in brackets give the 66% (likely) range over the GGCM-GCM ensemble.

	Wheat		Maize		Soybean		Rice	
	423ppm	486ppm	423ppm	486ppm	423ppm	486ppm	423ppm	486ppm
Global	-2.1 [-2.7,-0.1]	-1.9 [-2.4,0.3]	-3.3 [-4.7,0.6]	-3.1 [-4.4,0.6]	-3.7 [-5.7,-0.8]	-2.9 [-5.6,-0.9]	-3.8 [-4.1,-3.1]	-3.5 [-4.1,-3.2]
Tropics	-5.3 [-8.6,-2.9]	-4.3 [-6.6,-2.6]	-2.0 [-2.7,0.2]	-1.8 [-2.6,0.9]	-3.3 [-4.7,-2.3]	-3.4 [-5.1,-1.0]	-3.1 [-4.5,-3.1]	-3.1 [-4.4,-3.0]
AMZ	-	-	-1.3 [-2.3,2.1]	-1.1 [-2.1,2.8]	-2.6 [-4.2,-1.7]	-2.6 [-4.4,-1.9]	-3.1 [-3.5,-1.6]	-3.1 [-3.4,-1.5]
CAM	-4.9 [-6.4,-2.9]	-4.3 [-6.2,-2.2]	-3.1 [-4.9,0.8]	-2.1 [-3.8,0.9]	-3.9 [-6.1,-2.3]	-2.8 [-5.0,-1.6]	-4.0 [-5.3,-2.9]	-4.0 [-5.0,-2.9]
CAS	-1.8 [-6.3,-0.5]	-1.6 [-4.4,-0.4]	-2.5 [-4.5,0.6]	-2.0 [-3.2,2.9]	-	-	-3.5 [-6.3,-2.4]	-3.7 [-6.0,-2.5]
CEU	-2.1 [-3.8,-1.1]	-1.8 [-3.7,-0.8]	-3.0 [-6.5,1.2]	-3.0 [-6.2,1.5]	-2.7 [-7.5,0.4]	-2.1 [-5.8,3.0]	-5.3 [-5.9,-4.6]	-5.5 [-6.0,-3.0]
CNA	-2.4 [-7.7,1.7]	-1.5 [-6.2,1.9]	-3.7 [-8.3,1.7]	-3.7 [-8.6,2.0]	-5.4 [-10.2,-1.1]	-5.9 [-10.0,3.1]	-4.0 [-5.8,-3.4]	-4.3 [-5.9,-2.4]
EEF	-3.5 [-5.0,-1.8]	-3.4 [-4.4,-0.1]	-1.2 [-3.4,0.3]	-1.2 [-3.0,1.6]	-1.4 [-8.3,2.1]	-1.1 [-7.2,19.1]	-1.9 [-4.0,-0.8]	-1.9 [-4.3,-0.7]
EAS	0.7 [-0.3,3.1]	1.7 [-0.2,3.6]	-2.2 [-4.4,1.4]	-2.0 [-4.4,0.9]	-2.9 [-4.3,1.9]	-1.7 [-4.2,2.0]	-3.1 [-4.3,-2.8]	-3.2 [-4.0,-2.8]
ENA	-1.8 [-3.4,-0.3]	-1.1 [-3.1,-0.0]	-3.1 [-6.1,1.4]	-3.1 [-4.7,1.7]	-4.1 [-7.0,-0.7]	-3.8 [-6.6,0.2]	-	-
MED	-2.4 [-3.6,-0.7]	-2.1 [-3.3,0.1]	-2.2 [-5.9,0.3]	-2.2 [-6.0,0.6]	-6.4 [-9.5,-1.7]	-5.1 [-9.6,2.4]	-5.5 [-7.2,-4.1]	-4.3 [-7.0,-3.5]
NAS	-0.4 [-2.4,1.8]	-0.3 [-2.5,3.2]	1.6 [-2.1,8.3]	2.1 [-2.1,7.3]	1.6 [-2.4,9.6]	4.1 [-1.6,11.2]	-	-
NAU	-4.2 [-9.8,0.9]	-1.6 [-7.0,2.3]	-	-	-	-	-	-
NEB	-	-	-1.8 [-4.2,0.1]	-1.5 [-3.0,1.2]	-3.6 [-6.1,-1.8]	-3.5 [-6.1,-1.7]	-2.8 [-3.4,-1.3]	-2.9 [-3.6,-1.3]
NEU	-2.6 [-4.4,0.1]	-2.5 [-5.8,0.3]	-2.2 [-6.8,3.2]	-2.0 [-6.2,3.2]	-	-	-	-
SAF	-3.6 [-9.0,-1.2]	-3.3 [-8.4,-0.9]	-1.0 [-4.8,1.4]	-0.8 [-4.5,1.7]	-2.2 [-7.9,-0.1]	-2.3 [-6.6,0.1]	-1.9 [-4.5,-1.0]	-1.9 [-4.8,-0.9]
SAH	-3.1 [-5.6,1.1]	-1.3 [-5.1,1.9]	-4.4 [-6.4,-2.6]	-4.0 [-4.6,-1.6]	-	-	-5.3 [-8.5,-4.7]	-5.6 [-8.6,-4.1]
SAS	-3.3 [-8.2,-1.3]	-2.9 [-7.4,-0.9]	-2.3 [-3.2,0.9]	-1.6 [-3.1,2.4]	-2.8 [-4.2,-1.0]	-2.2 [-4.4,-0.6]	-4.2 [-5.0,-3.5]	-4.3 [-5.1,-3.5]
SAU	-0.4 [-2.9,2.3]	0.5 [-1.6,3.3]	-	-	-	-	-	-
SEA	-	-	-2.0 [-2.9,0.2]	-1.5 [-2.3,1.4]	-1.8 [-6.1,-0.2]	-1.8 [-2.6,1.8]	-2.9 [-4.3,-2.4]	-2.9 [-4.0,-2.4]
SSA	-3.2 [-6.9,-0.9]	-2.7 [-6.1,-0.2]	-1.1 [-3.7,0.6]	-0.7 [-2.7,1.4]	-2.9 [-9.3,-1.2]	-2.4 [-6.4,-1.2]	-2.1 [-4.0,-1.7]	-2.1 [-4.1,-1.6]
TIB	-2.3 [-4.8,2.7]	-2.0 [-4.4,2.8]	-1.6 [-3.8,-0.3]	-1.1 [-3.7,1.8]	-2.3 [-4.7,2.5]	-1.7 [-4.2,3.1]	-3.6 [-3.9,-2.7]	-3.8 [-4.1,-1.5]
WAF	-	-	-1.7 [-4.9,1.6]	-1.3 [-4.0,3.1]	-2.9 [-7.9,-0.3]	-1.6 [-3.9,0.0]	-1.6 [-3.9,-1.3]	-1.6 [-3.4,-1.3]
WAS	-0.1 [-3.3,1.2]	0.4 [-1.8,1.5]	-2.3 [-4.6,2.9]	-1.5 [-4.3,3.1]	-3.5 [-4.8,-0.4]	-1.2 [-4.5,2.7]	-5.8 [-6.9,-5.0]	-5.8 [-7.0,-4.5]
WNA	-4.3 [-10.8,-0.7]	-4.5 [-9.7,-0.2]	-2.6 [-4.6,0.7]	-2.2 [-4.5,-0.6]	-	-	-5.7 [-6.5,-5.4]	-6.0 [-6.7,-4.2]
WSA	-1.5 [-3.8,-0.7]	-1.1 [-2.9,2.3]	0.7 [-0.8,3.4]	0.8 [-0.0,4.3]	-	-	-1.3 [-2.3,0.1]	-1.2 [-2.5,0.2]

Table S5: Share of production areas (number of grid cells relative to global total based on MIRCA2000) for different the SREX regions and the Tropics (between 23.5°S/°N). Regions with less than 0.1% contribution to global crop production according to the MIRCA2000 dataset are masked out.

	Wheat	Maize	Soybean	Rice
Tropics	28%	44%	43%	58%
AMZ	-	7%	6%	11%
CAM	2%	3%	3%	4%
CAS	2%	1%	-	2%
CEU	5%	5%	4%	1%
CNA	5%	4%	6%	2%
EAF	4%	5%	5%	6%
EAS	9%	8%	10%	10%
ENA	2%	3%	2%	-
MED	4%	3%	4%	4%
NAS	9%	5%	5%	-
NAU	1%	-	1%	-
NEB	-	2%	2%	4%
NEU	1%	5%	-	-
SAF	5%	5%	5%	5%
SAH	1%	1%	-	1%
SAS	5%	4%	7%	7%
SAU	1%	-	-	-
SEA	-	4%	6%	7%
SSA	4%	4%	6%	5%
TIB	3%	1%	1%	1%
WAF	-	7%	5%	8%
WAS	4%	3%	4%	3%
WNA	5%	4%	-	1%
WSA	1%	1%	-	1%

## References Supplementary Material

- Lawrence D M, Oleson K W, Flanner M G, Thornton P E, Swenson S C, Lawrence P J, Zeng X, Yang Z-L, Levis S, Sakaguchi K and others 2011 Parameterization improvements and functional and structural advances in version 4 of the Community Land Model *J. Adv. Model. Earth Syst.* **3**
- Lawrence, D. M., Hurtt, G. C., Calvin, K. V., Jones, A. D., Jones, C. D., Lawrence, P. J., & Seneviratne, S. I. 2016. The Land Use Model Intercomparison Project (LUMIP) contribution to CMIP6: rationale and experimental design. *Geosci. Model Dev.*, 9(9), 2973.
- Liu W, Yang H, Folberth C, Wang X, Luo Q and Schulín R 2016a Global investigation of impacts of PET methods on simulating crop-water relations for maize *Agric. For. Meteorol.* **221** 164–75
- Liu W, Yang H, Liu J, Azevedo L B, Wang X, Xu Z, Abbaspour K C and Schulín R 2016b Global assessment of nitrogen losses and trade-offs with yields from major crop cultivations *Sci. Total Environ.* **572** 526–37
- Müller C, Elliott J, Chrysanthopoulos J, Arneth A, Balkovic J, Ciais P, Deryng D, Folberth C, Glotter M, Hoek S, Iizumi T, Izaurralde R C, Jones C, Khabarov N, Lawrence P, Liu W, Olin S, Pugh T A M, Ray D K, Reddy A, Rosenzweig C, Ruane A C, Sakurai G, Schmid E, Skalsky R, Song C X, Wang X, de Wit A and Yang H 2017 Global gridded crop model evaluation: benchmarking, skills, deficiencies and implications *Geosci. Model Dev.* **10** 1403–22
- Olin S, Lindeskog M, Pugh T A M, Schurgers G, Wärlind D, Mishurov M, Zaehle S, Stocker B D, Smith B and Arneth A 2015a Soil carbon management in large-scale Earth system modelling: Implications for crop yields and nitrogen leaching *Earth Syst. Dyn.* **6** 745–68
- Olin S, Schurgers G, Lindeskog M, Wärlind D, Smith B, Bodin P, Holmér J and Arneth A 2015b Modelling the response of yields and tissue C : N to changes in atmospheric CO<sub>2</sub> and N management in the main wheat regions of western Europe *Biogeosciences* **12** 2489–515