This document is the accepted manuscript version of the following article:
Ferrari, B. J. D., Vignati, D. A. L., Roullet, J. L., Coquery, M., Szalinska, E., Bobrowski, A., ... Dominik, J. (2019). Chromium bioavailability in aquatic systems impacted by tannery wastewaters. Part 2: new insights from laboratory and in situ testing with *Chironomus riparius* Meigen (Diptera, Chironomidae). Science of the Total Environment, 653, 1–9. <https://doi.org/10.1016/j.scitotenv.2018.10.258>

This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

**Chromium bioavailability in aquatic systems impacted by tannery wastewaters. Part 2:
new insights from laboratory and *in situ* testing with *Chironomus riparius* Meigen
(Diptera, Chironomidae)**

Published in Science of the Total Environment, doi:

<https://doi.org/10.1016/j.scitotenv.2018.10.258>

Ferrari B.J.D.^{a,b}, Vignati D.A.L.^{a,c}, Roullet J.-L.^d, Coquery M.^d, Szalinska E.^e, Bobrowski A.^f,
Czaplicka A.^g, Dominik, J.^{a,h}

a) Department F.-A. Forel for Environmental and Aquatic Sciences, University of Geneva,
Uni Carl Vogt, 66 boulevard Carl-Vogt CH-1211 Geneva, Switzerland

b) Swiss Centre for Applied Ecotoxicology Eawag-EPFL (Centre Ecotox), EPFL-ENAC-
IIE-GE, Station 2, 1015 Lausanne, Switzerland

c) Université de Lorraine, CNRS, LIEC, F-57000 Metz, France

d) Irstea, UR RiverLy, centre de Lyon-Villeurbanne, F-69625 Villeurbanne, France

e) Department of Environment Protection, Faculty of Geology, Geophysics and
Environmental Protection, AGH University of Science and Technology, 30 A.
Mickiewicza Av. , 30-059 Krakow, Poland.

f) Department of Building Materials Technology Faculty of Materials Science and Ceramics,
AGH University of Science and Technology, 30 A. Mickiewicza Av. , 30-059 Krakow,
Poland

g) Department of Water Supply, Sewerage and Environmental Monitoring, Cracow
University of Technology, 24 Warszawska ul., 31-155 Krakow, Poland

h) Institute of Marine Science - National Research Council (ISMAR-CNR) Arsenale - Tesa
104, Castello 2737/F, 30122 Venice, Italy

Abstract

Chromium is widely used as a tanning agent and can become a contaminant of concern in aquatic ecosystems receiving discharges from industrial or artisanal tanning activities. In a companion study, we showed that Cr discharged by tanneries was bioavailable to indigenous chironomids with accumulation via sediment ingestion likely to represent the predominant exposure route. However, Cr accumulation by chironomids did not directly reflect the degree of sediment contamination and the potential adverse effects of Cr accumulation on chironomids were not evaluated. In the present study, chironomids were exposed to homogenised, field-collected sediments in the laboratory and to intact sediments in situ using a customised caging system. Chromium concentrations were assessed in sediments, exposed larvae of laboratory-reared *Chironomus riparius* and overlying waters of in situ cages. Experimental results of Cr bioaccumulation were compared with expected Cr body burden in chironomids calculated using biodynamic modelling. Our data provided strong support to the hypothesis that Cr bioaccumulation in the field is specifically controlled by the deposition of contaminated suspended particulate matter (SPM) containing a pool of Cr readily bioavailable to surface deposit feeders. Considering freshly deposited SPM as an additional route of exposure for surface deposit feeders leads to a good agreement between the modelling and experimental results. Additionally, a Cr body burden of about $77 \mu\text{g g}^{-1} \text{ d.w}$ was identified as a tentative threshold above which effects on the growth of *C. riparius* may appear. While both laboratory and in situ experiments provide evidence for the availability of Cr in aquatic system impacted by tannery wastewaters, standard laboratory exposure conditions may miss additional exposure routes in the field and underestimate possible adverse effects on benthic organisms.

Keywords. Chironomids; Bioaccumulation; Suspended matter; Body residue; Sediment; Reservoir

1. Introduction

Bioaccumulation and toxicity of trace elements strongly depend on their speciation and on the actual exposure conditions experienced by target organisms; e.g. continuous vs. discontinuous inputs, contaminant uptake via different exposure routes (water vs. food), biological traits (e.g. feeding habit, respiration, growth, reproduction) of the species of interest (Fairbrother et al., 2007). Standardized laboratory experiments cannot account for all these factors although their importance is well-recognized and their predictive ability can be improved by a careful characterization of the actual exposure conditions in the selected laboratory settings (Simpson and Batley, 2007). Similarly, pulse and chase experiments using radiolabelled elements allow the study of the relative importance of different exposure routes, notably water exposure vs. food ingestion, in a variety of organisms (Baumann and Fisher, 2011; Wang et al., 1997), but cannot reproduce the complex array of environmental factors that influences bioavailability and toxicity in situ. To address these problems, the scientific community has engaged in actively developing tools to narrow the gap between laboratory and field studies. At present, microcosm and in situ techniques are sufficiently mature and versatile to be of use as supporting studies in tiered risk assessment procedures (Burton Jr et al., 2012; Crane et al., 2007; Ferrari and Faburé, 2017; Ferrari et al., 2014). An appropriate integration between laboratory and in situ approaches can help to better understand the bioavailability and toxicity of specific contaminants in ecosystems impacted by complex mixtures of contaminants such as tannery effluents (Vignati et al., 2007).

Adverse biological effects on aquatic organism in tannery contaminated systems are well documented (Khwaja et al., 2001; Koukal et al., 2004; Leghouchi et al., 2009). However, tannery effluents have a very complex composition and specifically ascribing the observed effects to Cr, typically used as a tanning agent in its trivalent form, remains difficult. Understanding of Cr bioavailability and toxicity in the field is further complicated by the

possible simultaneous presence of trivalent and hexavalent Cr forms that are characterized by different environmental mobility and toxicity (Vignati et al., this issue). On the other hand, Michailova et al. (2011) showed a genotoxic response in chironomids growing in a tannery-impacted system (the Dunajec river, southern Poland), where Cr and Cd were the only trace elements of concern. Although genotoxic and cancerogenic effects are mostly attributed to Cr(VI) exposure ((De Flora et al., 1990), genotoxic effects of Cr(III) organic compounds were also observed, along with changes in DNA and protein levels, in the yeast *Saccharomyces cerevisiae* (Chatterjee and Luo, 2010). The number of studies documenting a high toxicity of Cr(III) in standard laboratory settings is also increasing (Aharchaou et al., 2018; Bencheikh-Latmani et al., 2007; Kováčik et al., 2015; Ponti et al., 2014; Vignati et al., 2010), suggesting that Cr(III) is less harmless than admitted by the current scientific consensus. Significant polytene chromosome aberrations were found in *Chironomus riparius* exposed to Cr(III) contaminated sediments in laboratory conditions (Michailova et al., 2001).

In part 1 of the present study (Vignati et al., this issue), we showed that Cr discharged by tanneries is bioavailable to chironomids, but that the accumulation patterns do not simply reflect the contamination in abiotic matrices. By combining laboratory tests and in situ studies with caged, transplanted larvae of the model species *Chironomus riparius* Meigen (Diptera, Chironomidae), we specifically tested: a) if bioaccumulation in the field can be controlled by the ingestion of freshly-deposited contaminated SPM containing a pool of Cr readily bioavailable to surface deposit feeders such as chironomids and, b) at which level Cr accumulation may have adverse effects on the growth of *C. riparius*.

2. Material and Methods

2.1 Model organisms and study area

The non-biting midge larvae *Chironomus riparius* were used as model species for all experiments. This species is commonly used for ecotoxicological purposes, can be reared in laboratory conditions and has easily measurable and well modelled life-cycle traits. *Chironomus riparius* has been considered as a model organism in several internationally validated guidelines for assessing the toxicity of chemicals and for evaluating the quality of natural sediments (see Ferrari et al. 2017 and references therein).

Larvae used in the tests originated from a culture maintained in our laboratory. The culture was started from egg masses kindly provided by Dr. J. Garric (Irstea, formerly Cemagref, Lyon, France). The methods for maintaining the culture and preparing the organisms for experiments are detailed in Ferrari et al. (2014). For this study, two kinds of experiments were conducted, either by starting with young second instar larvae (7-day laboratory tests) or by starting with young fourth instar larvae (4-day laboratory and in situ tests). The larval stages were determined by checking the length of the larvae and by controlling on few specimens the head capsule width. Each set of experiment was designed to allow assessment of bioaccumulation of Cr, but the toxicity of the whole sediment on chironomids (growth and survival) was recorded at the same time. The chronological steps for implementing the tests are depicted in Figure S1.

In situ exposure and sample collection for the concomitant laboratory experiments were carried out in the Czorsztyn reservoir (Southern Poland, Fig. 1) which receives the tannery contaminated wastewaters of the Dunajec river (about 300 small tanneries in the Nowy Targ region) and of other smaller creeks. The in situ exposure/sampling site was located in Maniowy Bay (Fig. 1) which is locally fed by a small creek collecting the effluent from a rural wastewater treatment plant (Fig. 1). Tanneries are also present in the region of Maniowy, although only some of them confer their wastes to the wastewater treatment plant. More

details on Cr contamination in the Czorsztyn reservoir can be found elsewhere (Dominik et al., 2007; Szalinska et al., 2010) and in part 1 of this study (Vignati et al., this issue).

Figure 1. Study area (Czorsztyn reservoir; southern Poland) and location of the Maniowy Bay area where in situ exposures performed and samples collected for laboratory experiments. The Nowy Targ region, where the many artisanal tanneries are active, is also indicated.

2.2 In situ bioaccumulation

In situ exposure of *C. riparius* was carried out with a customized Sediment Water Interface Study System (S-WISS1.1) suitable for exposing surface deposit feeders to intact sediments for periods up to 4 days and allowing for food delivery to caged organisms during this period. Construction details and functioning were described in detail in Ferrari et al. (2014). Briefly, each exposure chamber consisted in a transparent polyvinyl chloride tube surmounted by a customized corer-head to deliver the chironomids, ensure water exchange between the

exposure chambers and the surrounding waters and suspensions (150 µm Nitex mesh) and deliver the required food doses.

Twelve S-WISS1.1 were deployed at Maniowy (water depth approx. 3 m) by scuba divers. In addition, twelve control cages consisting in 1-L high-density polyethylene (HDPE), large-mouth bottles containing clean sand and partly covered with 150 µm Nitex mesh (Ferrari et al., 2014) to ensure water exchange were deployed in the vicinity of the S-WISS1.1 devices. Food delivery in the control cages was also ensured by adopting the design described in Ferrari et al. (2014). At t=0, 20 individuals of *C. riparius* (4th instar larvae) were introduced into each exposure system (i.e. S-WISS1.1 and control cages). Food doses (1 mg Tetramin® per individual) were delivered immediately after deployment and then every 24 hours until the end of the exposure period (4 days). At t=24, 48, 72, and 96 hours, three exposure chambers and three control cages were randomly recovered.

Upon arrival at the laboratory, each S-WISS1.1 was opened and an aliquot of the overlying water was sampled, filtered at 1.2 µm, acidified to 1% v/v with concentrated (65 %) Suprapur HNO₃ and stored in a polypropylene (PP) container at 4°C pending Cr analysis. After siphoning-off the remaining water, an aliquot of the top layer of sediments (0–0.5 cm) was gently scraped with a plastic spatula, passed through a 1 mm sieve to remove debris, and possibly chironomids, and stored in a PP container at –22 °C pending Cr analysis.

The sediment from the surface layer was chosen because it represents the most relevant exposure zone for surface deposit-feeders such as *C. riparius* (Rasmussen, 1984). The whole sediment core was then sieved (500 µm) to recover surviving chironomids. No indigenous chironomids were found in the cores. Recovered organisms were rinsed with site water and placed for 5 min in trays filled with ultrapure water (MilliQ water, Millipore), followed by 10 min in 1 mM EDTA, and additional 5 min in ultrapure water. After gently blotting with paper towels, each larva was measured using a digital image analysis system and organisms were

pooled and conditioned for Cr analysis (see Ferrari et al., 2014 for more details). Individual length (n=20) and initial Cr body residue of unexposed larvae (2 pools of 30 individuals) were determined at the start of each experiment.

2.3 Laboratory bioaccumulation studies

Laboratory experiments were set up using both field-collected sediments from Maniowy and clean sand. Maniowy sediments were collected with an Ekman grab during the deployment of the in situ test systems and stored at 4 °C. Upon return to the laboratory, the overlying water was decanted and sediment were sieved at 1mm and homogenised on the day before the start of the tests. Sediments (clean sand for controls) were transferred into 1-L HDPE bottles analogous to those used for the control cages in the field experiment and covered with filtered (1.2 µm) Lake Geneva water at a minimum ratio of 1:4. The following day, 20 individuals of *C. riparius* were introduced into each exposure bottle and fed daily as the chironomids exposed in situ. Young second instar larvae were used for performing a 7-day bioaccumulation tests, while young fourth instar larvae were used for a 4-day tests.

All exposures were performed in triplicate at 21 °C under a 16:8-h light-dark photoperiod and light intensity of 500 lux. Air was continuously bubbled into each beaker during the test taking care to avoid sediment resuspension. The water level was controlled daily in each beaker and adjusted with deionised water if necessary. Larvae were fed daily ad libitum with 0.6 mg of commercial food (Tetramin®) per individual (Péry et al., 2002).

For the 4-day bioaccumulation tests, organisms were retrieved and handled for length measurement (i.e. each individual) and Cr content (i.e. pool of 20 individuals) as in situ exposed ones at day 1, 2, 3 and 4. For the 7-day exposure tests, organisms were retrieved and handled in the same way only at the end of the test. For each exposure condition, the surface layer sediments (or sand) were sampled for total Cr analysis.

2.4 Chemical analyses

Chromium concentration in sediments and chironomids were assayed using the same methodologies described in Vignati et al. (this issue). Total Cr concentrations in overlying waters (filtered at 1.2 µm) in S-WISS1.1 cages were determined by ICP-MS, while Cr(VI) levels were determined by catalytic adsorptive stripping voltammetry (Bobrowski et al., 2012). Accumulation by chironomids was corrected for gut sediment content according to (Hare et al., 1989) assuming that ingested sediments represented 6% of the total weight of individual organisms (see Ferrari et al., 2014). For quality control procedures, reference materials LKSD-4 (Lynch, 1990) and WQB3 (NRC, Canada) were used for sediments, TORT-2 (NRC, Canada) for biological material and SLRS3 (NRC, Canada) and 1643e (NIST) for waters. Reference material 1643e was analyzed following a 10-fold dilution.

2.5 Modelling of bioaccumulation

According to the biodynamic bioaccumulation modelling — DYMBAM (Baumann and Fisher, 2011; Wang et al., 1997), accumulation of a contaminant by living organism can be described by the following formula:

$$C_{biota} = \frac{k_u \cdot C_w}{k_{ew} + g} + \frac{AE \cdot IR \cdot C_s}{k_{es} + g} \quad (1)$$

where C_{biota} is the concentration accumulated by organisms (in µg g⁻¹), k_u is the uptake rate constant from the dissolved phase (in L g⁻¹ d⁻¹), C_w is the concentration in the water phase (filtered water, in µg L⁻¹), k_{ew} and k_{es} are the efflux rate constants following uptake from the dissolved phase and from food (sediments in the case of deposit feeders; both in d⁻¹), AE is the

assimilation efficiency (in % total content) from food (i.e., sediment), IR is the food ingestion rate (in $\text{g g}^{-1} \text{d}^{-1}$), C_s is the concentration in sediments (in $\mu\text{g g}^{-1}$) and g is the growth rate constant (in d^{-1}).

Although model parameters were not available for *C. riparius*, first-order estimates of expected Cr accumulation were obtained based on the following considerations. Accumulation from the aqueous phase was neglected considering that sediments should be the dominant route of contaminant uptake for chironomids (Dabrin et al., 2012; De Jonge et al., 2010; Martin et al., 2008). An ingestion rate of $0.325 \text{ g g}^{-1} \text{d}^{-1}$ estimated for *C. riparius* at 10°C (Bervoets et al., 2003) was adopted and corrected for the field/laboratory temperature of this study (21°C) as described in the supporting information. A value of 0.1%, determined using the biomimetic approach described in the companion paper (Vignati et al., this issue), was used for the AE of Cr from Maniowy sediments. Estimation of the organisms' growth rate constant (g) was performed by fitting an exponential curve to the plot of weight of individual chironomids' larvae vs. time (Fig. S2). Finally, the k_{es} constant was neglected considering that literature values for Cr egestion by various organisms (Baumann and Fisher, 2011; Roditi and Fisher, 1999; Wang et al., 1997) did not exceed 0.019 d^{-1} and thus were 4 to 8-fold lower than the growth rates calculated from our experimental data (Fig. S2). The conclusions of the present study would not have been altered by considering k_{es} in model calculations (see section 4.1) and this approach was preferred to the arbitrary choice of parameters derived for organisms other than *C. riparius*.

2.6 Statistics

Data were analyzed using Student's t-test, analysis of variance (one-way ANOVA followed by post-hoc Tukey's HSD test) and linear regression. All data were checked for normality and homogeneity of variance by the Shapiro-Wilk test and the Bartlett test,

respectively. If conditions for ANOVA were not fulfilled, the data were log-transformed and reanalysed. For all statistical tests, the significance level was set at 0.05 and calculations were performed using the software package R (v 2.9.0).

3. Results

3.1. Quality control

Figures of merit and overall accuracy of Cr analysis in aqueous matrices were analogous to those reported in Vignati et al. (this issue). Percentage Cr recovery from LKSD-4 was about 120% when based on the concentrated HNO₃/concentrated HCl extractable fraction ($21 \pm 2 \mu\text{g g}^{-1}$) and 75% when based on total Cr content determined by X-ray diffraction ($33 \pm 6 \mu\text{g g}^{-1}$; n=2 in both cases). For WQB3, measured Cr concentrations were $96 \pm 8 \%$ (n=5) of the indicative value of $119 \mu\text{g g}^{-1}$. Triplicate analyses of sediment subsamples from selected cages agreed to 10% or better. Chromium recovery from TORT-2 ($0.77 \pm 0.15 \mu\text{g g}^{-1}$) was $96 \pm 3 \%$ (n=3). Due to the insufficient biological material, no replicate analyses of exposed chironomids were possible. However, the small standard deviation on the three replicates of TORT-2 material suggests an acceptable precision in the digestion of the biological material.

3.2 Bioaccumulation under in situ and laboratory conditions

Chromium concentrations in sediments recovered inside the S-WISS1.1 units were homogeneous among the cages and across the four days of exposure ($325 \pm 35 \mu\text{g g}^{-1}$, n=11). The Cr content of homogenised sediments from Manioway used for the laboratory exposures was uniform ($150 \pm 14 \mu\text{g g}^{-1}$ d.w., n=12), albeit about half of that measured in the sediment recovered from in situ cages. Chromium levels in sand used for control exposures were $1.2 \pm 0.5 \mu\text{g g}^{-1}$ (n=12) in the field and $0.8 \pm 0.1 \mu\text{g g}^{-1}$ (n=12) in the laboratory (Fig. S3).

Bioaccumulation kinetic patterns in chironomids exposed to Maniowy sediments were similar for laboratory and in situ exposures, with a marked increase over the first day followed by a plateau in accumulated Cr levels. Daily and overall variability were however higher for organisms exposed in situ (Fig. 2).

Figure 2. Cr accumulation vs. time in *Chironomus riparius* larvae exposed to sand and sediment in experiments performed in situ (Maniowy Bay, June 2007) and in the laboratory using sand and field collected sediment. In all experiments, each point represents the measure of Cr internal concentration obtained for the pooled organisms retrieved in each exposure system. Concentrations in chironomids were gut corrected according to Hare et al. (1989).

At plateau (days 2–4), chironomids exposed in the laboratory to homogenised sediments from Maniowy accumulated an average Cr concentration of $78 \pm 16 \mu\text{g g}^{-1}$ ($n=12$), comparable to the concentration of $98 \pm 43 \mu\text{g g}^{-1}$ ($n=11$) measured in chironomids exposed in situ to undisturbed field sediments using the S-WISS1.1 system. At the opposite, bioaccumulation kinetic patterns in chironomids exposed to clean sand showed a marked difference between the laboratory and in situ exposures. Indeed, no increase in Cr concentration was observed during laboratory exposures with control chironomids accumulating less than $1 \mu\text{g g}^{-1}$ (Fig. 2). On the contrary, chironomids exposed in cages filled with clean sand but placed on the reservoir bottom close to the S-WISS1.1 devices accumulated an average of $77 \pm 25 \mu\text{g Cr g}^{-1}$ ($n=12$). This value is comparable to the concentrations obtained for chironomids exposed to the Maniowy sediments, whether in laboratory experiments or after in situ exposures (Fig. 2).

Results obtained by exposing 4th instar larvae in situ or in the laboratory also compared favourably with those of 2nd instar larvae exposed for 7 days to homogenised sediments in the laboratory (Fig. 3). Indeed, three days after the start of any type of exposure protocol using young 4th instar larvae (see sections 2.2 and 2.3), all larvae were 9 days old after hatching (Fig. S1). Thus, 4th instar larvae exposed over 3 days to sediments in situ or in the laboratory were as old as 4th instar larvae remaining at the end of the 7-day bioaccumulation test in the laboratory which was started with young 2nd instar larvae (Fig. S1).

For 3 days exposures of 4th instar larvae, Cr concentrations in organisms exposed to sediment and sand (i.e., controls) were $128 \pm 33 \mu\text{g g}^{-1}$ vs. $101 \pm 21 \mu\text{g g}^{-1}$ for in situ experiments and $77 \pm 5 \mu\text{g g}^{-1}$ vs. $0.42 \pm 0.07 \mu\text{g g}^{-1}$ for laboratory experiments (Fig. 3). A statistically significant difference in Cr accumulation (Tukey HDS test on log-transformed data, $p < 0.05$) existed between controls and exposed chironomids only in laboratory experiments. Despite the variability in the Cr content of sediment substrates (Fig. S3), Cr accumulation was not significantly different between organisms caged in the field (control

and exposed) and organisms exposed to homogenised Maniowy sediments in the laboratory (Tukey HDS test on log-transformed data, $p>0.05$). For 7 days experiments started using 2nd instar larvae, measured Cr accumulation at day 7 was $66\pm8 \mu\text{g g}^{-1}$ and $0.67\pm0.17 \mu\text{g g}^{-1}$ in organisms exposed to sediments and control sand, respectively. Chromium accumulation in 2nd instar larvae exposed to Maniowy sediment for 7 days was significantly lower than in 4th instar larvae exposed for 3 days to undisturbed field sediments (Tukey HDS test on log-transformed data, $p<0.05$), but did not show significant differences with other experimental conditions (Fig. 3).

Figure 3. Cr accumulation at day 3 in *Chironomus riparius* larvae exposed to sand and sediment in experiments performed in situ (Maniowy Bay, June 2007) and in the laboratory using field collected material. The 4-day tests were initiated with young fourth instar larvae, while the 7-day tests were initiated with young second instar larvae. Error bars represent 1 standard deviation ($n=3$). Internal concentrations that do not share the same letter

are significantly different (ANOVA followed by post-hoc Tukey's HSD test, $p < 0.05$). The dotted line symbolizes the level below which no effects were observed by considering the results obtained for growth rates (see Fig. S2).

3.3. Modelling chromium bioaccumulation via sediment ingestion

The modelled Cr accumulation via sediment ingestion by *C. riparius* exposed to Maniowy sediments did not exceed $1 \mu\text{g g}^{-1}$ for neither in situ nor laboratory exposures. These values were 1% (or less) of measured ones and strongly underestimated the actual Cr burden in both scenarios (Fig. 4a, 4b). Because the biomimetic approach used to derive the AE of 0.1% employed in model calculation (see section 2.5) may overlook other factors resulting in higher assimilation efficiencies (Vignati et al., this issue), additional model calculations were performed using AE values of 1, 2 and 5%. These AEs were reported in studies on Cr uptake by various organisms and from different substrates including sediments and algae (Baumann and Fisher, 2011; Roditi and Fisher, 1999; Wang et al., 1997). Even for the maximum AE value of 5%, modelled concentrations were 40–60% lower than the measured ones (Figure 4a, b).

Chironomids exposed to clean sand in laboratory experiments did not show any appreciable Cr bioaccumulation suggesting that neither the ingestion of sand substrate (in field controls) nor the added Tetramin® food (all experimental conditions) contributed to the observed Cr burdens. Measured Cr concentration in laboratory control chironomids actually decreased from $0.9 \pm 0.5 \mu\text{g g}^{-1}$ at day 1 to $0.4 \pm 0.06 \mu\text{g g}^{-1}$ at day 4 (data not shown). At $t=0$, control chironomids had a Cr content of about $2 \mu\text{g g}^{-1}$, which was diluted by the ingestion of 'clean' food and by growth. On the other hand, measured concentration in field control organisms were comparable to those of organisms exposed to Maniowy sediments (Fig. 3). As it will be discussed in detail in section 4.1, we surmise that accumulation from freshly

deposited SPM was responsible for the Cr burden measured in the field control chironomids and also contributed the majority of accumulated Cr to organisms exposed in situ to unperturbed sediments.

Figure 4. Comparison of measured vs. modelled Cr concentrations in *Chironomus riparius* exposed to undisturbed Maniowy sediments in situ (S-WISS1.1 cages – panel A) and to homogenised Maniowy sediments in the laboratory (panel B). All values are in $\mu\text{g g}^{-1}$. MOD, modelled accumulation of Cr obtained using different values for assimilation efficiency (AE) in formula 1; SPM, Cr accumulation in field control organisms (assumed to originate from ingestion of freshly deposited Suspended Particulate Matter — see text for details); Correction

‘on site’ and ‘S-WISS’, modelled accumulation using formula 1 and corrected for changes in bioavailability due to sediment manipulation according to Ferrari et al. (2014). The dashed lines show the theoretical 1:1 correspondence between modelled and measured concentrations.

3.4 Effects on chironomids

Survival of chironomids in laboratory controls was almost complete (19.6 ± 0.7 individuals; $n=12$) over the 4 days of exposure and only slightly reduced (18.1 ± 2.2 individuals, $n=12$) for exposure to Maniowy sediments in the laboratory. Average survival in field controls was 16.3 ± 3.7 individuals ($n=12$); significantly higher than in organisms directly caged on sediments in situ (9.8 ± 2.7 individuals, $n=11$). In this last case, no temporal trend in survival was observed.

The length of alive chironomids increased linearly and significantly ($R^2 > 0.84$, $p < 0.0001$) over time during the 4-day exposure both in the laboratory and in the field (Fig S2). In the laboratory, the growth rate of the chironomids exposed to sand was 0.1499 ± 0.0054 cm d⁻¹; not significantly different from the growth rate of 0.1421 ± 0.0096 cm d⁻¹ for the chironomids exposed to homogenised Maniowy sediments (ANOVA, $p > 0.05$; Fig. S2). In the field, the growth rate of the chironomids exposed to sand (0.1273 ± 0.0065 cm d⁻¹) was significantly lower than the value obtained for control chironomids in the laboratory (ANOVA, $p = 0.016$; Fig. S2). Growth rate further and significantly decreased to 0.092 ± 0.0114 cm d⁻¹ when the chironomids were exposed to the undisturbed sediments in situ using S-WISS1.1 (ANOVA, $p = 0.011$; Fig. S2).

Concerning the 7-day bioaccumulation test, no significant effect was observed between controls and exposed chironomids. The lowest number of alive chironomids per beaker ($n=3$) was 19 and 18 in controls and exposed organisms, respectively. The mean length at the end of

exposure period was 1.01 ± 0.02 cm in controls and 0.99 ± 0.05 cm in chironomids exposed to the homogenised Maniowy sediment (t-test, $p > 0.05$).

4. Discussion

4.1. Importance of SPM as an additional exposure route to contaminants in the field

The comparable Cr body burden in chironomids exposed in situ to undisturbed field sediments and to clean sand (Figs. 2 and 3) suggests the presence of an additional source of Cr for the organisms exposed in the field. The lower accumulation of Cr in chironomids exposed in the laboratory to homogenised sediments from Maniowy also supports the hypothesis that one (or more) additional source of Cr exists in the field. A possible contribution of Tetramin® to Cr bioaccumulation can be ruled out by the absence of Cr accumulation in control laboratory chironomids. We surmise that freshly deposited SPM entering the exposure cages can act as an important source of Cr to surface deposit feeders exposed in the field at the site of Maniowy. Besides the strong Cr accumulation by control chironomids exposed in situ, several observations collectively form a basis for evidence to support this hypothesis as detailed in the next paragraphs.

The lower accumulation by laboratory-exposed chironomids likely reflects the lower Cr concentration in homogenised sediments used for laboratory experiments compared with undisturbed field sediments (Fig. S3). The differences between homogenised and undisturbed (surface) sediments may be explained by the vertical decrease of Cr contamination in Maniowy sediments (Vignati et al., this issue). However, Ferrari et al. (2014) showed that, for sediments collected from the same site in Lake Geneva, Cr bioaccumulation was about 2- to 3-fold higher for chironomids exposed to homogenised sediments in the laboratory (sediment Cr content $25\text{--}30 \mu\text{g g}^{-1}$) than to undisturbed field sediments using S-WISS1.1 type cages ($28\text{--}32 \mu\text{g g}^{-1}$ of Cr) or to homogenised sediments placed back in the field ($27\text{--}32 \mu\text{g g}^{-1}$ of

Cr). In the study of Ferrari et al. (2014), temperature differences between laboratory (21 °C) and field (15 °C) alone could explain the observed differences. Sediment manipulation is also known to trigger a series of reactions that increase the bioavailability of sediment-bound contaminants (Vandeghechuchte et al., 2013); in disagreement with our observations. Overall, these previous observations suggest that the higher accumulation of Cr in chironomids exposed in the field to undisturbed Maniowy sediments requires a supplementary source of bioavailable Cr.

As observed in previous research on metal accumulation by chironomids (Dabrin et al., 2012; De Jonge et al., 2010; Martin et al., 2008), uptake of aqueous Cr species from overlying water and pore water is unlikely to be the main responsible for the observed bioaccumulation patterns in chironomids. In laboratory experiments, homogenized Maniowy sediments were covered with filtered water from Lake Geneva; which has a low Cr content (Edder et al., 2008; Kottelat, 2008) and should not contribute much to Cr accumulation. The occurrence of higher Cr body burdens in field control chironomids compared with organisms exposed to Maniowy sediments in the laboratory can therefore be considered indicative of Cr uptake from the overlying waters of Maniowy. This pattern was actually observed after 3 and 4 days of exposure (Figs. 2 and 3). However, differences in accumulated Cr concentrations were only 30% of the total body burdens, confirming the need for another source of Cr to explain the observed bioaccumulation in field controls. Furthermore, field control chironomids were exposed on a substrate of (initially) clean sand so that Cr uptake from pore water is unlikely to have contributed markedly to the elemental burden of these organisms.

Similar conclusions can be obtained by examining the total filterable Cr concentrations measured in overlying waters inside S-WISS1.1 cages (Table S1). These concentrations showed some variability between days 1+2 ($3.9 \pm 1.4 \mu\text{g L}^{-1}$; n=6) and 3+4 ($35 \pm 29 \mu\text{g L}^{-1}$; n=6), but Cr accumulation by chironomids was not directly proportional to filterable Cr

concentration inside the cages (Table S1). DGT measurements over 89 hours (slightly shorter than the 96 hours of chironomid exposure for logistic reasons) did not show any marked change in labile Cr in overlying or interstitial waters either (Fig. S4). The concentration of Cr(VI) measured in overlying waters of cages recovered on days 3 and 4 (Table S1) was $0.59 \pm 0.321 \mu\text{g L}^{-1}$ ($n=6$, range: $0.076\text{--}1.04 \mu\text{g L}^{-1}$) and comparable to the levels measured in pore waters recovered from sediment cores sampled in Maniowy (Vignati et al., this issue). Otherwise stated, caging did not markedly change Cr(VI) concentrations (which could have had consequences on Cr bioavailability) and DGT-labile Cr, a proxy for bioavailable trivalent Cr, remained constant during chironomids' exposure. Changes in pH, conductivity and oxygen content within the S-WISS1.1 cages are also too limited (Ferrari et al., 2014) to result in significant increases of Cr oxidation from field sediments. Considering the egestion rate of Cr assimilated via the dietary pathway (i.e., the k_{es} parameter in formula 1) would have further reduced the modelled Cr body burden. Overall, Cr uptake from freshly deposited SPM remains the hypothesis best supported by the available data to explain a) the observed Cr accumulation in field controls and b) the higher Cr accumulation by chironomids exposed to Maniowy sediment in situ compared with laboratory exposures to homogenised sediment.

4.2. A unifying framework for modelling, experimental results and field observations

The hypothesis that ingestion of freshly deposited SPM (section 4.1) represents a significant route of Cr accumulation for chironomids exposed in situ allows to reconcile modelling calculations and experimental results to a great extent. Summing accumulation via SPM (i.e., the experimental values measured for field control cages) to the quantity of Cr accumulated from sediment ingestion (using an AE of 0.1% as determined by the biomimetic approach – see section 2.5) markedly improves the agreement between modelled and measured bioaccumulation for organisms exposed to Maniowy sediments in situ (Fig. 4a).

With this procedure, modelled concentrations remain within 20% of measured ones for 2, 3 and 4 days of exposure, again supporting the hypothesis that SPM is a major route of exposure in field conditions. The $\pm 20\%$ differences are within the possible contribution of Cr uptake via the aqueous route; as indicated by the comparison between laboratory exposure to Maniowy sediments and in situ control cages (see section 4.1).

On the other hand, a contribution of freshly deposited SPM to Cr accumulation cannot be invoked to reduce the discrepancies between modelled and experimental Cr concentrations for chironomids exposed to Maniowy sediments in the laboratory (Fig. 4b). Accumulation via aqueous exposure, if any, would not lead to a better agreement between modelled and experimental results either (see sections 3.3 and 4.1). It is however possible that sediment manipulation during the setup of laboratory exposure may increase Cr bioavailability for chironomids. In the case of the Vidy Bay (a polluted bay in Lake Geneva, Switzerland/France), Ferrari et al. (2014) reported that Cr bioaccumulation by *C. riparius* in the laboratory was about twice than in specimen of the same species exposed in the field (S-WISS1.1 cages). Chromium accumulation in the laboratory was even 4–5 times higher than in organisms exposed on site in cages containing the same homogenised field sediments used in the laboratory. However, including a correction factor of 2 (in situ vs. laboratory) or 4.5 (on site vs. laboratory) into modelled bioaccumulation values for chironomids exposed to Maniowy sediment in the laboratory does not improve the agreement between observed and predicted results to any appreciable extent (Fig. 4b).

Another factor that may increase Cr bioavailability in laboratory exposure following sediment manipulation is the association of Cr with Fe/Mn oxides in Maniowy sediments (Vignati et al., this issue). However, even assuming a Cr assimilation efficiency of 5% for chironomids exposed to Maniowy sediments in the laboratory, modelled results still remain about 15% of the measured accumulation at day 4 (Fig. 4b). One last option is that the

development of an active Mn redox cycle at the sediment-water interface might have led to the formation of highly bioavailable Cr(VI) via oxidation of Cr(III) in the laboratory exposure beakers (Landrot et al., 2012). While the available experimental data do not allow to test this hypothesis, it appears that chromium bioaccumulation in the laboratory occurred to a different extent, and possibly via different routes, compared with real-field situations. Otherwise stated, the use of homogenised sediment to test for contaminant bioavailability may change the relative importance of the various exposure routes and lead to results not directly applicable in the field (Ferrari et al., 2014; Simpson and Batley, 2007; Wang et al., 2004).

It is also instructive to examine how well Cr accumulation by caged specimen of *C. riparius* can mimic Cr accumulation by indigenous chironomids. As already seen, consideration of Cr uptake via ingestion of freshly deposit SPM is necessary to reconcile DYMBAM calculations and experimentally measured accumulation for specimen of *C. riparius* exposed in situ to Maniowy sediment (Fig. 4a). However, even after accounting for the SPM route, the average accumulation of Cr by *C. riparius* inside S-WISS1.1 cages ($98 \pm 23 \mu\text{g g}^{-1}$; arithmetic mean $\pm$ one standard deviation for 4 days of exposure, $n=11$) remains about 3 times lower than accumulation by indigenous chironomids recovered from Maniowy sediments during the same period ($315 \mu\text{g g}^{-1}$ for the pool of chironomids sampling in June 2007; see Vignati et al., this issue). This difference may originate from individual indigenous chironomids weighing, on average, about 3.5 times as much as S-WISS1.1 exposed specimen on day 4 (1.43 vs. 0.388 mg). Larger individuals would ingest larger sediment quantities to meet their nutritional needs, while simultaneously accumulating more Cr from contaminated sediments. Furthermore, ingestion rate is inversely related to the organic matter content of a given sediment. Contrary to caged *C. riparius*, indigenous chironomids could not rely on Tetramin® as an additional source of Cr-free carbon; which could again have led to higher ingestion rates and, eventually, higher Cr body burdens. These considerations equally apply to

the ingestion of freshly deposited SPM and do not invalidate the hypothesis of SPM as the main Cr source to chironomids. Even if not performed in the same experimental conditions, it is interesting to note that Bervoets et al. (2004) found a very good agreement in Cr internal concentrations between caged and resident larvae, especially in the most contaminated sites.

4.3. Effect of chromium on the growth of chironomids

Individuals of *C. riparius* exposed to homogenised Maniowy sediments in the laboratory for 4 or 7 days did not exhibit higher mortality or reduced growth rate compared with control chironomids (section 3.4 and Fig. S2), indicating that Maniowy sediments were not toxic in standardized conditions. The absence of effects in laboratory exposures may be linked with sediment homogenisation. The Cr content of Maniowy sediments rapidly decreases below 5–6 cm (Vignati et al., this issue) because the Czorsztyn reservoir was flooded in 1997 and a thin layer of Cr-contaminated sediments overlies a relatively compact layer of former soil with low Cr content.

On the other hand, exposure to undisturbed sediments in the field resulted in reduced survival and growth rate compared with laboratory and field controls as well as with homogenised Maniowy sediments tested in the laboratory (section 3.4 and Fig. S2). During the in situ exposure period, water temperature in the field was around 21 °C and hence similar to the optimal temperature used for laboratory experiments. Moreover, chironomids have been daily fed ad libitum in all laboratory and in situ tests, thus preventing the impact of limiting food availability during in situ exposure. The differences between field controls and field exposed chironomids are therefore indicative of some kind of adverse effect of Maniowy sediments. Furthermore, the difference between field and laboratory controls also points to adverse effect linked to the presence of freshly deposited SPM which appears as an important route of exposure to Cr for chironomids during in situ experiments.

When considering the quantity of Cr accumulated by specimen of *C. riparius* in relation to the growth results, it can be assumed that no effect on growth can be observed up to a total concentration of $77 \mu\text{g g}^{-1}$ (Fig. 3 and Fig. S2). This concentration could correspond to a Critical Body Residue (CBR). The CBR is the concentration of a chemical bioaccumulated in an organism above which ecotoxicity effects appear (McCarty and MacKay, 1993). The tentative CBR value of $77 \mu\text{g g}^{-1}$ agrees with the data of Méndez-Fernández et al. (2013) who determined a CBR of approx. $34 \mu\text{g g}^{-1}$ for *Tubifex tubifex* in tests with natural sediments spiked with $\text{K}_2\text{Cr}_2\text{O}_7$. Upon entering living cells, Cr(VI) is rapidly converted to Cr(III) (Viti et al., 2014), so that a comparison of CBR for the two redox forms of Cr remains meaningful.

Recent studies have demonstrated that metal body burdens in relative resistant invertebrates such as chironomids can be used as predictors of ecological effects of metals on aquatic ecosystems. Bervoets et al. (2016) have linked threshold body burdens of metals in fourth instar larvae of indigenous *Chironomus* sp. to effects on resident macroinvertebrate communities. Depending on the descriptor chosen for the structure of the macroinvertebrate community, the threshold values ranged from 10 to $69 \mu\text{g g}^{-1}$ d.w. of Cr accumulated in chironomids. It is therefore possible that Cr contamination at Maniowy adversely affects sensitive taxa of macroinvertebrates. Furthermore, the tentative CBR value obtained for *C. riparius* in the present study suggests that Cr levels above $77 \mu\text{g g}^{-1}$ d.w. can also affect relatively pollution-tolerant organisms such as chironomids themselves. These findings are not at odds with the thresholds proposed by Bervoets et al. (2016) because laboratory-reared organisms are likely more sensitive than adapted local indigenous species.

However, the bioaccumulation results obtained in this study showed a large variability (Fig. 2) which hampers a clear definition of the actual CBR for Cr in *C. riparius*. Other studies also suggest that metals have toxic effects only when their uptake rate exceeds their combined rates of efflux and detoxification (Casado-Martinez et al., 2010) or the fraction of

metabolically available metal increases (Rainbow, 2007). Knowledge of the subcellular fate of metals is therefore required to better understand the physiological processes underlying their bioaccumulation and toxicity. Such methods have been developed for larvae of chironomids (Gimbert et al., 2016; Péry et al., 2008) and will have to be considered to further clarify the metabolically available concentration of Cr that can affect the development of chironomids at different stages in real-field situations.

5. Conclusions

Comparable levels of Cr accumulation by specimen of *Chironomus riparius* exposed to undisturbed field sediments in situ and to clean sand on-site suggest that ingestion of freshly deposited SPM can be the main route of Cr accumulation by surface deposit feeders in real field conditions. This additional route of exposure is not accounted for in standardized laboratory experiments or, as yet, by biodynamic bioaccumulation modelling, so that both approaches will lead to an underestimation of Cr accumulation compared with field conditions. Changes in the bioavailability of sediment-bound Cr also seem to occur during sediment sampling and handling, with the correction of such experimental artefacts being far from straightforward. While both laboratory and in situ experiments provide evidence for the availability of Cr in aquatic system impacted by tannery wastewaters, in situ investigations appear necessary to obtain an adequate understanding of Cr bioavailability under real environmental conditions.

A tentative threshold of Cr accumulation for effects on the growth of *C. riparius* is proposed at $77 \mu\text{g g}^{-1}$ d.w. Further studies using toxicokinetics approaches and subcellular fractionation methods are however required to refine the actual concentration of Cr linked to the biologically active sites and leading to adverse effects in these benthic organisms. Extrapolation from the model organism *C. riparius* to other species of indigenous

chironomids also represents a future challenge to better link laboratory and in situ results with real field situations.

Acknowledgements

This research was supported by the Swiss National Fund (grants 200020-101844 and 200020-117942). The authors express their sincerest thanks to Michal Krzyszkowski and the direction of the Niedzica Hydro-Electric Power Plant (Poland) for their precious cooperation. We are indebted to Mr. Przemek Kramin and Mr. Tomasz Smereczynski for handling of S-WISS1.1 in situ exposure systems in the Czorsztyn reservoir. We also thank our colleagues Vincent Chanudet, Régis Kottelat and Stephanie Jüstrich from the Institute F.-A. Forel for their help during field work.

References

- Aharchaou I, Py JS, Cambier S, Loizeau JL, Cornelis G, Rousselle P, Battaglia E, Vignati DAL. 2018. Chromium hazard and risk assessment: New insights from a detailed speciation study in a standard test medium. *Environmental Toxicology and Chemistry*. 37(4):983-992.
- Baumann Z, Fisher NS. 2011. Modeling metal bioaccumulation in a deposit-feeding polychaete from labile sediment fractions and from pore water. *Science of The Total Environment*. 409(13):2607-2615.
- Bencheikh-Latmani R, Obraztsova A, Mackey MR, Ellisman MH, Tebo BM. 2007. Toxicity of Cr(III) to *Shewanella* sp. strain mr-4 during Cr(VI) reduction. *Environ Sci Technol*. 41:214-220.

621 Bervoets L, De Bruyn L, Van Ginneken L, Blust R. 2003. Accumulation of ¹³⁷Cs by larvae of
 622 the midge *Chironomus riparius* from sediment: effect of potassium. Environ Toxicol
 623 Chem. 22(7):1589-1596.

624 Bervoets L, De Jonge M, Blust R. 2016. Identification of threshold body burdens of metals for
 625 the protection of the aquatic ecological status using two benthic invertebrates.
 626 Environmental Pollution. 210(Supplement C):76-84.

627 Bervoets L, Meragalli G, De Cooman W, Goddeeris B, Blust R. 2004. Caged midge larvae
 628 (*Chironomus riparius*) for the assessment of metal bioaccumulation from sediment in
 629 situ. Environ Toxicol Chem. 23:443-454.

630 Bobrowski A, Kapturski P, Zarębski J, Dominik J, Vignati DAL. 2012. Catalytic adsorptive
 631 stripping voltammetric determination of chromium(VI) in overlying and interstitial
 632 waters isolated from sediments contaminated by tannery waste. Analytical Letters.
 633 45(5-6):495-507.

634 Burton Jr GA, Rosen G, Chadwick DB, Greenberg MS, Taulbee WK, Lotufo GR, Reible DD.
 635 2012. A sediment ecotoxicity assessment platform for in situ measures of chemistry,
 636 bioaccumulation and toxicity. Part 1: System description and proof of concept.
 637 Environmental Pollution. 162:449-456.

638 Casado-Martinez CM, Smith BD, Luoma SN, Rainbow PS. 2010. Metal toxicity in a
 639 sediment-dwelling polychaete: Threshold body concentrations or overwhelming
 640 accumulation rates? Environmental Pollution. 158(10):3071-3076.

641 Chatterjee N, Luo Z. 2010. Cr-(III)-organic compounds treatment causes genotoxicity and
 642 changes in DNA and protein level in *Saccharomyces cerevisiae*. Ecotoxicology.
 643 19(4):593-603.

644 Crane M, Burton GA, Culp JM, Greenberg MS, Munkittrick KR, Ribeiro R, Salazar MH, St-
645 Jean SF. 2007. Review of aquatic in situ approaches for stressor and effect diagnosis.
646 Integrated Environ Assess Manag. 3(2): 234-245.

647 Dabrin A, Durand CL, Garric J, Geffard O, Ferrari BJD, Coquery M. 2012. Coupling
648 geochemical and biological approaches to assess the availability of cadmium in
649 freshwater sediment. Science of The Total Environment. 424:308-315.

650 De Flora S, Bagnasco M, Serra D, Zancchi P. 1990. Genotoxicity of chromium compounds.
651 A review. Mutation Research/Reviews in Genetic Toxicology. 238(2):99-172.

652 De Jonge M, Blust R, Bervoets L. 2010. The relation between acid volatile sulfides (avs) and
653 metal accumulation in aquatic invertebrates: Implications of feeding behavior and
654 ecology. Environmental Pollution. 158(5):1381-1391.

655 Dominik J, Vignati DAL, Koukal B, Pereira de Abreu M-H, Kottelat R, Szalinska E, Bas B,
656 Bobrowski A. 2007. Speciation and environmental fate of chromium in rivers
657 contaminated with tannery effluents. Engineerig Life Science. 7:155-169.

658 Edder P, Ortell D, Klein A, Ramseier S. 2008. Metals and organic micropollutants in Geneva
659 Lake waters and sediments. Rapp. Comm. int. prot. eaux Léman contre pollut.,
660 Campagne 2007, 57-84.

661 Fairbrother A, Wenstel R, Sappington K, Wood W. 2007. Framework for metals risk
662 assessment. Ecotoxicology and Environmental Safety. 68(2):145-227.

663 Ferrari BJD, Faburé J. 2017. Field assessment of reproduction-related traits of chironomids
664 using a newly developed emergence platform (e-board). Ecotoxicology and
665 Environmental Safety. 137:186-193.

666 Ferrari BJD, Vignati DAL, Dominik J. 2014. Bioaccumulation kinetics and effects of
667 sediment-bound contaminants on chironomids in deep waters: New insights using a
668 low-disturbance in situ system. Environmental Technology. 35(4):456-469.

669 Gimbert F, Geffard A, Guédron S, Dominik J, Ferrari BJD. 2016. Mercury tissue residue
670 approach in *Chironomus riparius*: Involvement of toxicokinetics and comparison of
671 subcellular fractionation methods. *Aquatic Toxicology*. 171:1-8.

672 Hare L, Campbell PGC, Tessier A, Belzile N. 1989. Gut sediment in a burrowing mayfly
673 (Ephemeroptera, *Hexagenia limbata*): their contribution to animal trace element
674 burdens, their removal, and the efficacy of a correction for their presence. *Can J Fish*
675 *Aquat Sci*. 46:451-456.

676 Khwaja AR, Singh R, Tandon SN. 2001. Monitoring of Ganga water and sediments vis-à-vis
677 tannery pollution at Kanpur (India): A case study. *Environmental Monitoring and*
678 *Assessment*. 68(1):19-35.

679 Kottelat R. 2008. Caractérisation physico-chimique de microcosmes alimentés en continu et
680 leur utilisation dans l'étude des voies d'exposition de Cd et Cr (III) chez "*Daphnia*
681 *magna*". *Terre et Environnement, Université de Genève*; 73: 191 p.

682 Koukal B, Dominik J, Vignati D, Arpagaus P, Santiago S, Ouddane B, Benaabidate L. 2004.
683 Assessment of water quality and toxicity of polluted rivers Fez and Sebou in the
684 region of Fez (Morocco). *Environ Poll*. 131:163-172.

685 Kováčik J, Babula P, Hedbavny J, Kryštofová O, Provazník I. 2015. Physiology and
686 methodology of chromium toxicity using alga *Scenedesmus quadricauda* as model
687 object. *Chemosphere* 120: 23-30.

688 Landrot G, Ginder-Vogel M, Livi K, Fitts JP, Sparks DL. 2012. Chromium(III) Oxidation by
689 Three Poorly-Crystalline Manganese(IV) Oxides. 1. Chromium(III)-Oxidizing
690 Capacity. *Environmental Science & Technology* 46: 11594-11600.

691 Leghouchi E, Laib E, Guerbet M. 2009. Evaluation of chromium contamination in water,
692 sediment and vegetation caused by the tannery of Jijel (Algeria): a case study.
693 *Environmental Monitoring and Assessment* 153: 111-117.

694 Lynch J. 1990. Provisional elemental composition values for eight new geochemical lake
 695 sediment and stream sediment reference materials LKSD-1, LKSD-2, LKSD-3,
 696 LKSD-4, STSD-1, STSD-2, STSD-3, STSD-4. *Geostandards Newsletter* 14: 153-167.
 697 Martin S, Proulx I, Hare L. 2008. Explaining metal concentrations in sympatric chironomus
 698 species. *Limnology and Oceanography* 53(2):411-419.
 699 McCarty LS, MacKay D. Enhancing ecotoxicological modeling and assessment. 1993.
 700 *Environ. Sci. Technol.* 27: 1719-1728.
 701 Méndez-Fernández L, Martínez-Madrid M, Rodriguez P. 2013. Toxicity and critical body
 702 residues of cd, cu and cr in the aquatic oligochaete tubifex tubifex (müller) based on
 703 lethal and sublethal effects. *Ecotoxicology* 22(10): 1445-1460.
 704 Michailova P, Petrova N, Sella G, Bovero S, Ramella L, Regoli F, Zelano V. 2001. Genotoxic
 705 effects of chromium on polytene chromosomes of *Chironomus riparius* Meigen 1804
 706 (Diptera, Chironomidae). *Caryologia* 54(1): 59-71.
 707 Michailova P, Szarek-Gwiazda E, Kownacki A, Warchalowska-Sliwa E. 2011. Biodiversity
 708 of chironomidae (Diptera) and genome responses to trace metals in the environment.
 709 *Pestycydy/Pesticides* 1-4: 41-48.
 710 Péry ARR, Geffard A, Conrad A, Mons R, Garric J. 2008. Assessing the risk of metal
 711 mixtures in contaminated sediments on *Chironomus riparius* based on cytosolic
 712 accumulation. *Ecotoxicology and Environmental Safety* 71: 869-873.
 713 Péry ARR, Mons R, Flammarion P, Lagadic L, Garric J. 2002. A modeling approach to link
 714 food availability, growth, emergence, and reproduction for the midge *Chironomus*
 715 *riparius*. *Environ. Toxicol. Chem.* 21: 2507-2513.
 716 Ponti B, Bettinetti R, Dossi C, Vignati DAL. 2014. How reliable are data for the ecotoxicity
 717 of trivalent chromium to daphnia magna? *Environmental Toxicology and Chemistry*
 718 33(10): 2280-2287.

719 Rainbow PS. 2007. Trace metal bioaccumulation: Models, metabolic availability and toxicity.
 720 Environment International 33: 576-582.

721 Rasmussen JB. 1984. Comparison of gut contents and assimilation efficiency of fourth instar
 722 larvae of two coexisting chironomids, *Chironomus riparius* Meigen and
 723 *Glyptotendipes paripes* (Edwards). Can. J. Zoology 62: 1022-1026.

724 Roditi HA, Fisher NS. 1999. Rates and routes of trace elements uptake in zebra mussel.
 725 Limnol. Oceanogr. 44: 1730-1749.

726 Simpson SL, Batley GE. 2007. Predicting metal toxicity in sediments: A critique of current
 727 approaches. Integrated Environmental Assessment and Management 3: 18-31.

728 Szalinska E, Dominik J, Vignati DAL, Bobrowski A, Bas B. 2010. Seasonal transport pattern
 729 of chromium(III and VI) in a stream receiving wastewater from tanneries. Applied
 730 Geochemistry 25: 116-122.

731 Vandegehuchte MB, Nguyen LTH, De Laender F, Muyssen BTA, Janssen CR. 2013. Whole
 732 sediment toxicity tests for metal risk assessments: on the importance of equilibration
 733 and test design to increase ecological relevance. Environmental Toxicology and
 734 Chemistry 32: 1048-1059.

735 Vignati DAL, Dominik J, Beye ML, Pettine M, Ferrari BJD. 2010. Chromium(VI) is more
 736 toxic than chromium(III) to freshwater algae: A paradigm to revise? Ecotoxicology
 737 and Environmental Safety 73: 743-749.

738 Vignati DAL, Ferrari BJD, Dominik J. 2007. Laboratory-to-field extrapolation in aquatic
 739 sciences. Environ. Sci. Technol. 41: 1067-1073.

740 Vignati DAL, Ferrari BJD, Roulier JL, Coquery M, Szalinska E, Bobrowski A, Czaplicka A,
 741 Dominik J. Chromium bioavailability in aquatic systems impacted by tannery
 742 wastewaters. Part 1: understanding chromium accumulation by indigenous
 743 chironomids. Science of The Total Environment ; this issue.

- Viti C, Marchi E, Decorosi F, Giovannetti L. 2014. Molecular mechanisms of Cr(VI) resistance in bacteria and fungi. FEMS Microbiology Reviews 38: 633-659.
- Wang F, Goulet RR, Chapman PM. 2004. Testing sediment biological effects with the freshwater amphipod *Hyaella azteca*: the gap between laboratory and nature. Chemosphere 57: 1713-1724.
- Wang W-X, Griscom SB, Fisher NS. 1997. Bioavailability of Cr(III) and Cr(VI) to marine mussels from solute and particulate pathways. Environ. Sci. Technol. 31: 603-611.

Supporting information

Chromium bioavailability in aquatic systems impacted by tannery wastewaters. Part 2: new insights from laboratory and *in situ* testing with *Chironomus riparius* Meigen (Diptera, Chironomidae)

Ferrari B.J.D.^{a,b}, Vignati D.AL.^{a,c}, Roulier J.-L.^d, Coquery M.^d, Szalinska E.^e, Bobrowski A.^f,
Czaplicka A.^g, Dominik, J.^{a,h}

i) Department F.-A. Forel for Environmental and Aquatic Sciences, University of Geneva,
Uni Carl Vogt, 66 boulevard Carl-Vogt CH-1211 Geneva, Switzerland

j) Swiss Centre for Applied Ecotoxicology Eawag-EPFL (Centre Ecotox), EPFL-ENAC-
IIE-GE, Station 2, 1015 Lausanne, Switzerland

k) Université de Lorraine, CNRS, LIEC, F-57000 Metz, France

l) Irstea, UR RiverLy, centre de Lyon-Villeurbanne, F-69625 Villeurbanne, France

m) Department of Environment Protection, Faculty of Geology, Geophysics and
Environmental Protection, AGH University of Science and Technology, 30 A.
Mickiewicza Av. , 30-059 Krakow, Poland.

n) Department of Building Materials Technology Faculty of Materials Science and Ceramics,
AGH University of Science and Technology, 30 A. Mickiewicza Av. , 30-059 Krakow,
Poland

o) Department of Water Supply, Sewerage and Environmental Monitoring, Cracow
University of Technology, 24 Warszawska ul., 31-155 Krakow, Poland

p) Institute of Marine Science - National Research Council (ISMAR-CNR) Arsenale - Tesa
104, Castello 2737/F, 30122 Venice, Italy

*Corresponding Author: benoit.ferrari@centreecotox.ch

Correction of sediment ingestion rate (IR) for temperature

Temperature correction for sediment ingestion rates was calculated according to the formula:

$$Q_{10} = \left(\frac{R_2}{R_1} \right)^{\left(\frac{10}{T_2 - T_1} \right)} \quad (A)$$

where: Q_{10} (dimensionless) is the rate of increase of a reaction for a 10 degrees temperature increase (in °C or K), R_1 and R_2 are the reaction rates at T_1 and T_2 , respectively (with $T_1 < T_2$). Web-based calculator (http://www.physiologyweb.com/calculators/q10_calculator.html) was used to estimated temperature-adjusted sediment ingestion rates entering an arbitrary Q_{10} value of 2 in formula (A) (see below for the rationale behind this choice). Other input values for formula A were the reference temperature of 10°C used by Bervoets et al. (2003), the corresponding IR value of 0.325 g g⁻¹ d⁻¹ for *C. riparius* and the measured field and laboratory temperature of 21°C for our study. A temperature corrected value of IR equal to 0.697 g g⁻¹ day⁻¹ resulted from this calculations.

The Q_{10} factor of 2 was selected based on (Croteau et al., 2002; Gresens, 2001) who studied the effect of temperature on Cd accumulation and sediment ingestion and digestion. It is important to note that uncertainties exist as to the influence of temperature on sediment ingestion rates and contaminant accumulation by chironomids. In particular, the relationship between temperature and sediment ingestion rate is not necessarily a monotonous one (Gresens, 2001) and its determination would require experimental approaches outside the scope of the present work.

Table S1. Concentration of total Cr measured in overlying waters recovered inside each of the three S-WISS cages on days 1, 2, 3, and 4; concentrations of hexavalent Cr in the same waters; and concentration of total Cr in the chironomids exposed within each cage.

Day	Replicate	Water		Chironomids
		Total Cr	Cr(VI)	Total Cr ^a
		($\mu\text{g L}^{-1}$)	($\mu\text{g L}^{-1}$)	($\mu\text{g g}^{-1}$) d.w.
1	1	3.42	n.a.	68.1
1	2	6.31	n.a.	150
1	3	4.31	n.a.	79.2
2	1	2.48	n.a.	29.9
2	2	2.87	n.a.	63.0
2	3	3.93	n.a.	127
3	1	93.5	0.57	125
3	2	26.1	0.61	162
3	3	21.8	0.46	97.3
4	1	23.8	1.04	54.9
4	2	29.3	0.08	n.a.
4	3	18.7	0.76	125

a) Concentrations corrected for sediment content in the gut (see main text for details)

Figure S1.

Figure S1. Chronological step for implementing the bioaccumulation and toxicity tests with the non-biting midge larvae *Chironomus riparius*. For the present study, experiments were started either with young second instar larvae (7 day laboratory tests) or with young fourth instar larvae (4 day laboratory and in situ tests). Numbers correspond to the post-hatching days needed to reach the end of the corresponding test. Note that 2nd instar larvae used in 7-day toxicity test and 4th instar larvae used in bioaccumulation tests will have the same post-hatching age at day 9.

Figure S2.

Figure S2. Evolution of the length (a) and weight (b) of 4th instar *Chironomus riparius* larvae as a function of time in bioaccumulation experiments performed in the field (Maniow Bay, June 2007) and in the laboratory using field collected material. Blue diamonds correspond to the measured responses of the chironomids exposed to the sand, while the red squares to the measured responses of those exposed to Maniow sediments. In figures (a), each point represents the average of the lengths obtained by individually measuring live specimen recovered daily from each exposure batch. The lines represent linear regressions and parameters b1, b2, b3 and b4 their slopes (means $\pm$ 1 s.e., in cm d⁻¹) defined as the representative growth rates (g) for the different conditions. Slopes that do not share the same capital letter are significantly different (ANOVA, $p < 0.05$). In the figures (b), each point represents the weight of the pooled live chironomids. The lines represent exponential regressions and parameters c1, c2, c3 and c4 their slopes (means $\pm$ 1 s.e., in d⁻¹). R^2 and p indicate the goodness-of-fit and the statistical significance of each fitted regression.

Figure S3.

Figure S3. Total Cr concentrations (mg kg⁻¹ d.w.) in sediments used in bioaccumulation experiments performed in situ (Maniowy Bay, June 2007) and in the laboratory using field collected material. Data are the arithmetic mean $\pm$ one standard deviation (n=12) of Cr concentration in each system of exposure retrieved during the 4 days of exposure *in situ* or in the laboratory. Note the log scale on the vertical axis.

Figure S4

Figure S4. Temporal and vertical variability of DGT-labile Cr concentrations in interstitial (negative depth values; in cm) and overlying (positive depth values; in cm) at Maniowy bay in June 2007. Four different DGT probes were exposed at the same time and recovered after 22, 45, 68 and 89 hours of exposures. Exposure location was close to the cages for chironomids and is assumed to be representative of Cr behaviour within the cages.