



Multiradionuclide evidence for an extreme solar proton event around 2,610 B.P. (~660 BC)

Paschal O'Hare^a, Florian Mekhaldi^a, Florian Adolphi^{a,b}, Grant Raisbeck^c, Ala Aldahan^d, Emma Anderberg^a, Jürg Beer^e, Marcus Christl^f, Simon Fahrni^f, Hans-Arno Synal^f, Junghun Park^g, Göran Possnert^h, John Southonⁱ, Edouard Bard^j, ASTER Team^{j,1}, and Raimund Muscheler^{a,2}

^aDepartment of Geology—Quaternary Sciences, Lund University, 22362 Lund, Sweden; ^bClimate and Environmental Physics & Oeschger Centre for Climate Change Research, Physics Institute, University of Bern, 3012 Bern, Switzerland; ^cCentre de Sciences Nucléaires et de Sciences de la Matière, CNRS, Université Paris-Saclay, 91405 Orsay, France; ^dDepartment of Geology, United Arab Emirates University, Al Ain, United Arab Emirates; ^eDepartment of Surface Waters, Swiss Federal Institute of Aquatic Science and Technology, 8600 Dübendorf, Switzerland; ^fLaboratory of Ion Beam Physics, Swiss Federal Institute of Technology, 8093 Zurich, Switzerland; ^gKorea Institute of Geoscience and Mineral Resources, 34132 Daejeon, Korea; ^hTandem Laboratory, Uppsala University, 751 20 Uppsala, Sweden; ⁱKeck/Accelerator Mass Spectrometry (AMS) Laboratory, University of California, Irvine, CA 92697-3100; and ^jCNRS, Institut de Recherche pour le Développement (IRD), Institut National de la Recherche Agronomique, Coll France, UMR 34 Centre Européen de Recherche et d'Enseignement des Géosciences de l'Environnement (CEREGE), Aix-Marseille University, 13545 Aix-en-Provence, France

Edited by Lennard A. Fisk, University of Michigan, Ann Arbor, MI, and approved February 4, 2019 (received for review September 13, 2018)

Recently, it has been confirmed that extreme solar proton events can lead to significantly increased atmospheric production rates of cosmogenic radionuclides. Evidence of such events is recorded in annually resolved natural archives, such as tree rings [carbon-14 (¹⁴C)] and ice cores [beryllium-10 (¹⁰Be), chlorine-36 (³⁶Cl)]. Here, we show evidence for an extreme solar event around 2,610 years B.P. (~660 BC) based on high-resolution ¹⁰Be data from two Greenland ice cores. Our conclusions are supported by modeled ¹⁴C production rates for the same period. Using existing ³⁶Cl ice core data in conjunction with ¹⁰Be, we further show that this solar event was characterized by a very hard energy spectrum. These results indicate that the 2,610-years B.P. event was an order of magnitude stronger than any solar event recorded during the instrumental period and comparable with the solar proton event of AD 774/775, the largest solar event known to date. The results illustrate the importance of multiple ice core radionuclide measurements for the reliable identification of short-term production rate increases and the assessment of their origins.

solar storms | radionuclides | ice cores | solar proton events

Our Sun sometimes produces highly energetic particles, which are accelerated either by magnetic reconnection in solar flares or by shock waves associated with coronal mass ejections. Such energetic particles then follow trajectories along the interplanetary magnetic field lines, which together with the location of the event on the Sun, determine whether these particles hit the Earth's atmosphere. These phenomena are referred to as solar proton events (SPEs). Such events represent a threat to modern society in terms of communication and navigation systems, space technologies, and commercial aircraft operations (1, 2). Therefore, better understanding the possible magnitudes and occurrence frequency of such events is of great importance for safeguarding space technologies and modern technological infrastructure. During the past ~60 y, these events have been recorded instrumentally and are commonly described quantitatively by their fluence (the number of incident particles per area integrated over the event) of protons with kinetic energy greater than 30 MeV (F_{30}). Sometimes, SPEs are so energetic (large fluence >100 MeV; F_{100}) that they can lead to increased counts in surface-based neutron monitors known as ground-level enhancements (GLEs). The largest of these GLEs occurred on February 23, 1956 (3) (also called GLE05 according to the numbering of the instrumentally observed GLEs: www.nmdb.eu/nest/gle_list.php) and is estimated to have had an F_{30} of 1.8×10^9 protons per 1 cm^2 [$F_{100} = 3.0 \times 10^8$ protons per 1 cm^2 (4)]. The relatively short instrumental record does not allow for robust estimates of the frequency of extreme SPEs, and it cannot be used to reliably define the upper limit of our Sun's eruptive capacity.

While the use of ice core nitrate to document these events has been rejected (5–7), an extended record of the fluence, frequency, and energy distribution of SPEs can be obtained through the analysis of cosmogenic radionuclides, such as beryllium-10 (¹⁰Be), carbon-14 (¹⁴C), and chlorine-36 (³⁶Cl) (8–10). These radionuclides are mainly produced via a nuclear cascade triggered by galactic cosmic rays reaching the Earth's atmosphere on average with much higher kinetic energy than the solar protons. Incoming galactic cosmic rays are modulated by the heliomagnetic and geomagnetic fields, with the strength of this modulation changing from decadal to millennial timescales (11–13). However, strong SPEs can lead to large fluxes of solar protons, causing a short-term rapid increase in the atmospheric production of radionuclides, which are subsequently stored in environmental archives, such as tree rings (¹⁴C) and ice cores (¹⁰Be and ³⁶Cl). A recent study by Mekhaldi et al. (9) used a series of ice core records (14) to confirm a solar origin for two rapid increases in $\Delta^{14}\text{C}$ (¹⁴C/¹²C corrected for fractionation and decay relative to a standard) in AD 774/775 and AD 993/994 first identified in tree rings (15–18). Mekhaldi et al. (9) have proposed that the stronger

Significance

This study provides evidence of an enormous solar storm around 2,610 B.P. It is only the third such event reliably documented and is comparable with the strongest event detected at AD 774/775. The event of 2,610 years B.P. stands out because of its particular signature in the radionuclide data [i.e., carbon-14 (¹⁴C) data alone does not allow for an unequivocal detection of the event]. It illustrates that present efforts to find such events based solely on ¹⁴C data likely lead to an underestimated number of such potentially devastating events for our society. In addition to ¹⁴C data, high-resolution records of beryllium-10 and chlorine-36 are crucial for reliable estimates of the occurrence rate and the properties of past solar proton events.

Author contributions: R.M. designed research; P.O. and F.M. performed research; A.A., E.A., J.B., M.C., S.F., H.-A.S., J.P., G.P., J.S., E.B., and A.T. contributed data; P.O., F.M., F.A., G.R., and R.M. analyzed data; and P.O., F.M., and R.M. wrote the paper with input from all coauthors.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

This open access article is distributed under Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND).

¹A complete list of the ASTER Team can be found in *SI Appendix*.

²To whom correspondence should be addressed. Email: Raimund.Muscheler@geol.lu.se.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1815725116/-DCSupplemental.

Published online March 11, 2019.




Fig. 2. Multiradionuclide measurements for the 2,610-y B.P. (~660 BC) event. (A) Time series for the newly measured NGRIP ^{10}Be concentration (red curve, left axis) with corresponding measurement error margins and estimated natural baseline (dashed red line). Baseline concentration for ^{10}Be is calculated as the average ^{10}Be concentration for the measured period excluding the three peak values that span about 2.3 y. The red envelope represents the ^{10}Be production range attributable to a solar modulation Φ varying between 500 and 1,200 MeV, which corresponds to a typical modern 11-y cycle (36). This estimate assumes that ^{10}Be variations in Greenland ice cores vary proportionally to the global average ^{10}Be production rate changes as supported by ^{10}Be – ^{14}C comparison studies (29). NGRIP ^{10}Be concentration measurements have been overlaid on the modeled ^{14}C production rate inferred from the data shown in Fig. 1 (gray curve, right axis) with 1σ uncertainties (gray error bars). The ^{14}C production rate is normalized to pre-industrial absolute production rates. (B) Time series for ^{10}Be (red curve, left axis) (ref. 26 and this study) and ^{36}Cl concentrations measured in the GRIP ice core (blue curve, right axis) (21), with associated measurement errors (1σ) and calculated baseline concentration for ^{10}Be and ^{36}Cl (dashed blue line). Red and blue envelopes are as per A but considering the data's lower resolution for ^{10}Be and ^{36}Cl , respectively. All ice core data are plotted on the timescale according to ref. 29. Please note that the timescale in A is stretched as indicated by the lines between the panels.

production only (31). Furthermore, in the case of solar protons, we expect radionuclide production almost exclusively in the stratosphere due to the comparably lower energies of the solar particles (32). Due to the relatively long lifetime of ^{10}Be in the stratosphere, the stratosphere can be considered well mixed within each hemisphere (31). The results of Heikkilä et al. (31) indicate that the relative contribution of stratospheric ^{10}Be to the ^{10}Be deposition in Greenland is very close to the global average stratospheric ^{10}Be

contribution to the global production and deposition (69 vs. 65%) (Table 2) (“control run” in ref. 31), supporting the assumption that we can expect the relative changes in radionuclide deposition in Greenland to be close to the relative changes in global average production. Radionuclide concentration/production values exceeding 3σ of the calculated natural background level around 2,610 y B.P. are assumed to be related to the event and have been integrated into a single year (with calculated baseline removed). These integrated values are represented as enhancement values (Table 1) and include errors incorporating uncertainties in measurement as well as a baseline variability, which results from the 11-y solar modulation variability and noise inherent in the data. The NGRIP ^{10}Be measurements indicate a concentration peak factor (enhancement divided by baseline estimate) of 2.52 ± 0.91 , and the GRIP ^{10}Be and ^{36}Cl data indicate concentration peak factors of 4.99 ± 0.99 and 6.36 ± 1.36 , respectively. The GRIP ^{10}Be numbers were calculated the same way as for the corresponding ^{36}Cl data (same resolution and number of data points). To obtain the $^{36}\text{Cl}/^{10}\text{Be}$ of the enhancement region, we have used two procedures. For the case of ^{10}Be in NGRIP and ^{36}Cl in GRIP, we used the procedure of Mekhaldi et al. (9), which is to subtract the baseline contribution under the peak region using the SD of ^{36}Cl in GRIP and ^{10}Be in NGRIP. With this calculation, we get an enhancement of 1.27 ± 0.36 . However, for the case of both isotopes in GRIP, we can take advantage of the fact that variable factors, such as accumulation, solar variability, and duration of the enhancement region, are the same for both isotopes. We calculate the $^{36}\text{Cl}/^{10}\text{Be}$ ratio (R) for each of the sample pairs in Fig. 2B. As expected, this ratio outside the peak region is constant within experimental uncertainties ($R = 0.168 \pm 0.019$). Thus, when subtracting the baseline contribution of ^{36}Cl , we can use the ^{36}Cl baseline estimated from ^{10}Be (^{36}Cl baseline = $R \times ^{10}\text{Be}$ baseline). With this approach, we get that $^{36}\text{Cl}/^{10}\text{Be}$ in the enhancement region is equal to 1.27 ± 0.28 . The smaller uncertainty compared with using two different cores results from the fact that the baseline contributions under the peak regions are not independent. This emphasizes the advantages of having the two isotopes measured in the same core.

The peaks in both ^{10}Be and ^{36}Cl concentrations are synchronous with the peak in ^{14}C production rate after correction for known timescale offsets (29) (*SI Appendix*). However, the uncertainty in estimating the peak amplitude from ^{14}C is large, since the enhancement is not well defined in the ^{14}C record. This global signature increase, which is somewhat reminiscent of AD 774/775 and AD 993/994, strongly points to the occurrence of a strong short-term global increase in radionuclide production rates around 2,610 y B.P.

The three radionuclide records used in this study all display slightly different patterns in terms of the peak duration ranging from 2 to 6 y, despite the expected rapid nature of cosmic ray

Table 1. Summary of results for the 2,610-y B.P. event

Variable	NGRIP ^{10}Be	GRIP ^{10}Be	GRIP ^{36}Cl
Baseline, atoms per gram	$1.60 \pm 0.50 \times 10^4$	$1.23 \pm 0.13 \times 10^4$	$2.08 \pm 0.3 \times 10^3$
Integrated enhancement, atoms per gram	$4.03 \pm 0.76 \times 10^4$	$6.16 \pm 0.95 \times 10^4$	$13.2 \pm 2.4 \times 10^3$
Peak factor	2.52 ± 0.91	4.99 ± 0.93	6.36 ± 1.36
$^{36}\text{Cl}/^{10}\text{Be}$	2.52 ± 0.99	1.27 ± 0.28	

Estimates of the baseline production, enhancement (integrated peak concentration above baseline), and enhancement factor for ^{10}Be and ^{36}Cl (atoms per gram ice) for the 2,610-y B.P. event as shown in Fig. 2. The peak factor relates the integrated radionuclide enhancement to the baseline radionuclide production over a year. Uncertainties are based on error propagation, including measurement errors and baseline variability of 1σ . The peak factor for the NGRIP ^{10}Be data contains an additional uncertainty, since one ^{10}Be measurement just before the ^{10}Be increase was lost. Furthermore, the baseline calculation for NGRIP might be less robust due to the relatively short dataset. The error calculation for GRIP $^{36}\text{Cl}/^{10}\text{Be}$ is explained in the text. The $^{36}\text{Cl}/^{10}\text{Be}$ ratio for NGRIP ^{10}Be is based on ^{10}Be from NGRIP and ^{36}Cl from GRIP.

Table 2. Relative $^{36}\text{Cl}/^{10}\text{Be}$ ratios and estimated fluences for the SPE events discussed in this study

SPE	GLE no.	Relative $^{36}\text{Cl}/^{10}\text{Be}$ ratio	$F > 30$ MeV protons per 1 cm^2	$F > 100$ MeV protons per 1 cm^2	$F > 360$ MeV protons per 1 cm^2
February 23, 1956	GLE05	1.2	1.8×10^9	3.0×10^8	6.0×10^7
January 20, 2005 (SPE05)	GLE69	1.5	2.0×10^8	6.0×10^7	1.5×10^7
2,610 y B.P. (~660 BC)		1.4 ± 0.3	$2.09 (\pm 0.75) \times 10^{10}$	$6.3 (\pm 2.28) \times 10^9$	$1.57 (\pm 0.56) \times 10^9$
AD 774/775		1.8 ± 0.2	$2.82 (\pm 0.25) \times 10^{10}$	$8.5 (\pm 0.75) \times 10^9$	$2.12 (\pm 0.18) \times 10^9$
AD 993/994		2.1 ± 0.4	$1.02 (\pm 0.21) \times 10^{10}$	$3.1 (\pm 0.64) \times 10^9$	$7.65 (\pm 0.16) \times 10^8$

Ratios shown are based on computations of the annual mean production of ^{10}Be and ^{36}Cl by a series of large SPEs measured between 1956 and 2005 (4) (a full list is in *SI Appendix*). The ratios calculated by Mekhaldi et al. (9) for the AD 774/775 and AD 993/994 are included along with the ratio for the 2,610-y B.P. event calculated in this study. The fluence (F) values of protons per 1 cm^2 above 30, 100, and 360 MeV for the instrumentally observed events are from Webber et al. (4). The events are ordered according to the estimated spectral hardness.

events. The 2.3-y long peak in the highest-resolution NGRIP ^{10}Be record can be attributed to a rapid production rate increase and the subsequent transport from the stratosphere, where radionuclides are mainly produced, to its geological archive (33, 34). ^{14}C shows an unexpectedly long enhancement that lasts about 4–6 y. Within the relatively large errors, the ^{14}C data alone cannot be used to pinpoint the source of the radionuclide enhancement. A reason could be the relatively low sensitivity of the atmospheric $\Delta^{14}\text{C}$ to short-term enhancements in the ^{14}C production rate and the corresponding large errors in the reconstructed ^{14}C production rate. In the case of GRIP ^{10}Be and ^{36}Cl , the broad ~6-y peak can be attributed to the low resolution of the dataset owing to sample size requirements of the original GRIP project. The ^{36}Cl enhancement reported here is comparable with ^{36}Cl peaks around AD 775 and AD 994 (9). The estimates for the relative ^{10}Be and ^{36}Cl increases are listed in Table 1. The production of ^{36}Cl was more enhanced during the 2,610-y B.P. event compared with ^{10}Be . This pattern is in accordance with the expected production signature of cosmogenic radionuclides by solar energetic protons (4, 35). As shown in Fig. 24, the short-term rapid increase in ^{10}Be cannot be explained by typical solar modulation. More specifically, the red band in Fig. 24 represents the ^{10}Be production range attributable to a solar modulation Φ varying between 500 and 1,200 MeV, which is the range of variance estimated for the past 60 y (36). Fig. 24 shows that the three data points assumed here to be related to the same event are all well above this production range. Modeled ^{14}C production for the 2,610-y B.P. period supports the increased production rate, although the peak in modeled ^{14}C production has large uncertainties and appears broader than expected as discussed above.

Discussion

Excluding Other Sources/Support for a Solar Origin. The Sun impacts the radionuclide production rate indirectly by modulating the galactic cosmic ray flux reaching the Earth’s atmosphere, most notably in the form of the 11-y solar cycle (37). The 2,610-y B.P. production increases in ^{14}C , ^{10}Be , and ^{36}Cl described in this study cannot be explained by this modulation for several reasons. First, there is a theoretical limit for the production increase that is reached when solar shielding of galactic cosmic rays shifts from average shielding (which corresponds to the estimated baseline) to no shielding as mentioned above. The red and blue bands in Fig. 2 show the typical variability in radionuclide production rates over an 11-y solar cycle based on the past 60 y. If we assume that one such cycle is interrupted by a period of completely absent solar shielding, we would expect an ~65% increase in the radionuclide production rate exceeding the upper limit of this range (red and blue bands in Fig. 2) (36). Both the high-resolution NGRIP ^{10}Be and GRIP ^{36}Cl concentration increase exceed this theoretical threshold, while the GRIP ^{10}Be data are at that limit within uncertainty. Furthermore, the lower resolution of the GRIP data and the smoothing during transport and deposition of ^{10}Be and ^{36}Cl lead to a reduced amplitude in the

radionuclide data compared with the actual production rate increase. Second, the effect of solar modulation on the production of ^{10}Be and ^{36}Cl is very similar (i.e., leading to changes of the $^{10}\text{Be}/^{36}\text{Cl}$ ratios even during varying solar or geomagnetic shielding of galactic cosmic rays of less than 1%) (38). The larger enhancement of ^{36}Cl compared with ^{10}Be for the 2,610-y B.P. event, therefore, points to another process, likely a production rate increase due to lower-energy solar particles, as the ratio is expected to increase if solar protons significantly contribute to the radionuclide production rate (4).

The incorporation of ^{14}C into the carbon cycle results in an attenuation of high-frequency peaks in the atmospheric ^{14}C concentrations. This leads to larger errors of the reconstructed ^{14}C production rate changes and therefore, at least in this case, to difficulties in robustly identifying the SPE event even with precisely measured ^{14}C data (Fig. 24). Due to the relatively large errors of the inferred ^{14}C production rate, a solar modulation origin of the ^{14}C peak cannot be excluded. However, the $\Delta^{14}\text{C}$ peak around 2,610 y B.P. (Fig. 1) exceeds significantly the $\Delta^{14}\text{C}$ variations during an 11-y cycle, which are typically in the range of 4‰ (25). The differences to ^{10}Be could be attributed to ^{14}C measurement uncertainties, which result in the large error band of the reconstructed ^{14}C production rate, carbon cycle uncertainties, and/or transport effects on ^{10}Be . Furthermore, the ^{10}Be and ^{14}C production rate differences around the 2,610-y B.P. peak are similar to the ^{10}Be – ^{14}C differences before and after the peak (Fig. 24), indicating the challenges of robustly identifying short-term production rate increases with individual radionuclide records.

Aside from solar modulation and the occurrence of extreme solar events, several theories have been proposed as responsible for rapid short-term increases in cosmogenic radionuclide production. One such theory involves a short gamma ray burst resulting in irradiation of the atmosphere by a flux of high-energy gamma rays (18, 39, 40). Based on production cross-section measurements, Raisbeck et al. (41) estimated an about four times larger relative production rate increase in ^{14}C compared with ^{10}Be for gamma ray-induced radionuclide production. Therefore, as with the AD 774/775 event, we can reject this hypothesis, since it contradicts our data.

The ^{10}Be data from this study, when analyzed in conjunction with existing GRIP ^{36}Cl records, support the hypothesis of an extreme SPE being responsible for a sharp peak in radionuclide production around 2,610 B.P. The initial investigation of a sharp change in $\Delta^{14}\text{C}$ in the IntCal13 dataset combined with modeled ^{14}C production based on annual ^{14}C tree ring measurements adds support to our hypothesis but also, leaves open questions regarding ^{14}C measurements alone to robustly infer past solar storm events.

Spectral Hardness. For a SPE to result in a measurable increase in cosmogenic radionuclide production, its proton fluence is expected to be large (4). This is illustrated by the fact that no SPE in the instrumental period has been unequivocally linked to

calculate the ^{14}C production rates to correct for postproduction effects as a result of ^{14}C entering the carbon cycle (SI Appendix).

^{10}Be Data. Before this study, there were no published high-resolution ^{10}Be data for the 2,610-y B.P. period. Completing the GRIP ^{10}Be record with a resolution of about 3 y supports the exceptional GRIP ^{36}Cl increase around 2,610 y B.P. The NGRIP ice was sampled at a constant resolution of ~ 11 cm, which resulted in an average temporal resolution of about 0.75 y. That is with the exception of two samples, which were sampled at a resolution of 18.3 cm (1.2-y resolution) due to less ice available, which would hinder quality measurements (SI Appendix).

^{36}Cl Data. Existing ^{36}Cl measurements in the GRIP ice core (21), which have an average resolution of ~ 6 y, were investigated for the same period. Due to the greater sensitivity of the ^{36}Cl production rate for lower-energy solar protons (4), ^{36}Cl is, therefore, of particular interest when investigating SPEs.

ACKNOWLEDGMENTS. We thank Carmen Vega for advice concerning the preparation of ^{10}Be samples for AMS measurements. We also thank three anonymous reviewers and the editorial team for constructive comments on the manuscript. This work was supported by a Royal Physiographic Society of Lund grant (to F.M.) and Swedish Research Council Grant DNR2013-8421 (to

R.M.). F.A. was supported by Swedish Research Council Grant DNR2016-00218. J.P. was supported by the Basic Research Project of the Korea Institute of Geoscience and Mineral Resources funded by the Ministry and Information and Communications Technology of Korea. NGRIP is directed and organized by the Center of Ice and Climate at the Niels Bohr Institute and the US NSF. It is supported by funding agencies and institutions in Belgium (Fonds de la Recherche Scientifique-Communauté Française de Belgique and Research Foundation - Flanders), Canada (Natural Resources Canada/Geological Survey of Canada), China (Chinese Academy of Sciences), Denmark (Danish Natural Science Research Council and FIST), France [Institut Français pour la Recherche et la Technologie Polaires, Institut Polaire Français Paul Emile Victor, CNRS/Institut National des Sciences de l'Univers (INSU), Commissariat à l'Énergie Atomique, and Agence Nationale de la Recherche (ANR)], Germany (Alfred Wegener Institut), Iceland (The Icelandic Centre for Research), Japan (Ministry of Education, Culture, Sports, Science and Technology and National Institute of Polar Research), Korea (Korea Polar Research Institute), The Netherlands (Netherlands Organization for Scientific Research/Section Earth and Life Sciences), Sweden (Swedish Research Council and Swedish Polar Research Secretariat), Switzerland (Swiss National Science Foundation), the United Kingdom (Natural Environment Research Council), and the United States (US NSF Polar Programs). The French AMS national facility ASTER (CEREGE, Aix-en-Provence, France) is supported by the INSU/CNRS, the ANR through the "Projets thématiques d'excellence" Program for the "Equipements d'excellence" ASTER-CEREGE action, and IRD.

- NRC (2008) *Severe Space Weather Events: Understanding Societal and Economic Impacts: A Workshop Report* (National Academies Press, Washington, DC).
- Shea M, Smart D (2012) Space weather and the ground-level solar proton events of the 23rd solar cycle. *Space Sci Rev* 171:161–188.
- Meyer P, Parker E, Simpson J (1956) Solar cosmic rays of February, 1956 and their propagation through interplanetary space. *Phys Rev* 104:768.
- Webber W, Higbie P, McCracken K (2007) Production of the cosmogenic isotopes ^3H , ^7Be , ^{10}Be , and ^{36}Cl in the Earth's atmosphere by solar and galactic cosmic rays. *J Geophys Res Space Phys* 112:A10106.
- Wolff EW, et al. (2012) The Carrington event not observed in most ice core nitrate records. *Geophys Res Lett* 39:L08503.
- Mekhaldi F, et al. (2017) No coincident nitrate enhancement events in polar ice cores following the largest known solar storms. *J Geophys Res Atmos* 122:11900–11913.
- Sukhodolov T, et al. (2017) Atmospheric impacts of the strongest known solar particle storm of 775 AD. *Sci Rep* 7:45257.
- Lal D, Peters B (1967) Cosmic ray produced radioactivity on the earth. *Handbuch der Physik*, ed Flüggé S (Springer, Berlin), Vol 46/2, pp 551–612.
- Mekhaldi F, et al. (2015) Multiradionuclide evidence for the solar origin of the cosmic-ray events of AD 774/5 and 993/4. *Nat Commun* 6:8611.
- Usoskin IG, Solanki SK, Kovaltsov GA, Beer J, Kromer B (2006) Solar proton events in cosmogenic isotope data. *Geophys Res Lett* 33:L08107.
- Beer J, Siegenthaler U, Oeschger H, Bonani G, Finkel R (1988) Information on past solar activity and geomagnetism from Be-10 in the Camp Century ice core. *Nature* 331:675–679.
- Muscheler R, Adolphi F, Herbst K, Nilsson A (2016) The revised sunspot record in comparison to cosmogenic radionuclide-based solar activity reconstructions. *Sol Phys* 291:3025–3043.
- Vonmoos M, Beer J, Muscheler R (2006) Large variations in Holocene solar activity: Constraints from ^{10}Be in the Greenland Ice Core Project ice core. *J Geophys Res Space Phys* 111:A10105.
- Sigl M, et al. (2015) Timing and climate forcing of volcanic eruptions for the past 2,500 years. *Nature* 523:543–549.
- Güttlér D, et al. (2015) Rapid increase in cosmogenic ^{14}C in AD 775 measured in New Zealand kauri trees indicates short-lived increase in ^{14}C production spanning both hemispheres. *Earth Planet Sci Lett* 411:290–297.
- Jull A, et al. (2014) Excursions in the ^{14}C record at AD 774–775 in tree rings from Russia and America. *Geophys Res Lett* 41:L3004–3010.
- Miyake F, Masuda K, Nakamura T (2013) Another rapid event in the carbon-14 content of tree rings. *Nat Commun* 4:1748.
- Miyake F, Nagaya K, Masuda K, Nakamura T (2012) A signature of cosmic-ray increase in AD 774–775 from tree rings in Japan. *Nature* 486:240–242.
- Park J, Southon J, Fahrni S, Creasman PP, Mewaldt R (2017) Relationship between solar activity and $\Delta 14\text{C}$ peaks in AD 775, AD 994, and 660 BC. *Radiocarbon* 59:1147–1156.
- Yiou F, et al. (1997) Beryllium 10 in the Greenland Ice Core Project ice core at Summit, Greenland. *J Geophys Res* 102:26783–26794.
- Wagner G, et al. (2000) Reconstruction of the geomagnetic field between 20 and 60 kyr BP from cosmogenic radionuclides in the GRIP ice core. *Nucl Instrum Methods Phys Res B* 172:597–604.
- Thomas BC, Melott AL, Arkenberg KR, Snyder BR (2013) Terrestrial effects of possible astrophysical sources of an AD 774–775 increase in ^{14}C production. *Geophys Res Lett* 40:L237–1240.
- Usoskin I, et al. (2013) The AD775 cosmic event revisited: The Sun is to blame. *Astron Astrophys* 552:L3.
- Reimer PJ, et al. (2013) IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years cal BP. *Radiocarbon* 55:1869–1887.
- Stuiver M, Braziunas TF (1993) Sun, ocean, climate and atmospheric $^{14}\text{CO}_2$: An evaluation of causal and spectral relationships. *Holocene* 3:289–305.
- Muscheler R, et al. (2004) Changes in the carbon cycle during the last deglaciation as indicated by the comparison of ^{10}Be and ^{14}C records. *Earth Planet Sci Lett* 219:325–340.
- Raisbeck GM, et al. (1981) Cosmogenic $^{10}\text{Be}/^{7}\text{Be}$ as a probe of atmospheric transport processes. *Geophys Res Lett* 8:1015–1018.
- Heikkilä U, Beer J, Feichter J (2008) Modeling cosmogenic radionuclides ^{10}Be and ^7Be during the maunder minimum using the ECHAM5-HAM general circulation model. *Atmos Chem Phys* 8:2797–2809.
- Adolphi F, Muscheler R (2016) Synchronizing the Greenland ice core and radiocarbon timescales over the Holocene—Bayesian wiggle-matching of cosmogenic radionuclide records. *Clim Past* 12:15–30.
- Muscheler R, Beer J, Kubik PW, Synal H-A (2005) Geomagnetic field intensity during the last 60,000 years based on ^{10}Be and ^{36}Cl from the Summit ice cores and ^{14}C . *Quat Sci Rev* 24:1849–1860.
- Heikkilä U, Beer J, Feichter J (2009) Meridional transport and deposition of atmospheric ^{10}Be . *Atmos Chem Phys* 9:515–527.
- Poluianov SV, Kovaltsov GA, Mishev AL, Usoskin IG (2016) Production of cosmogenic isotopes ^7Be , ^{10}Be , ^{14}C , ^{22}Na , and ^{36}Cl in the atmosphere: Altitudinal profiles of yield functions. *J Geophys Res Atmos* 121:8125–8136.
- Heikkilä U (2007) Modeling of the atmospheric transport of the cosmogenic radionuclides ^{10}Be and ^7Be using the ECHAM5-HAM general circulation model. PhD dissertation ETH No. 17516 (ETH, Zurich).
- Lal D, Peters B (1967) Cosmic ray produced radioactivity on the Earth. *Handbuch der Physik*, ed Flüggé S (Springer, Berlin), pp 551–612.
- Castagnoli G, Lal D (1980) Solar modulation effects in terrestrial production of carbon-14. *Radiocarbon* 22:133–158.
- Herbst K, Muscheler R, Heber B (2017) The new local interstellar spectra and their influence on the production rates of the cosmogenic radionuclides ^{10}Be and ^{14}C . *J Geophys Res Space Phys* 122:23–34.
- Schwabe M (1844) Sonnenbeobachtungen im Jahre 1843. Von Herrn Hofrath Schwabe in Dessau. *Astron Nachr* 21:233.
- Masarik J, Beer J (1999) Simulation of particle fluxes and cosmogenic nuclide production in the Earth's atmosphere. *J Geophys Res Atmos* 104:12099–12111.
- Pavlov A, et al. (2013) AD 775 pulse of cosmogenic radionuclides production as imprint of a Galactic gamma-ray burst. *Mon Not R Astron Soc* 435:2878–2884.
- Hambaryan V, Neuhäuser R (2013) A Galactic short gamma-ray burst as cause for the ^{14}C peak in AD 774/5. *Mon Not R Astron Soc* 430:32–36.
- Raisbeck GM, et al. (1992) ^{10}Be deposition at Vostok, Antarctica, during the last 50,000 years and its relationship to possible cosmogenic production variations during this period. *The Last Deglaciation: Absolute and Radiocarbon Chronologies*, eds Bard E, Broecker WS (Springer, Erice, Italy), 1st Ed, Vol 2, pp 127–140.
- Baroni M, Bard E, Petit J-R, Magand O, Bourlès D (2011) Volcanic and solar activity, and atmospheric circulation influences on cosmogenic ^{10}Be fallout at Vostok and Concordia (Antarctica) over the last 60 years. *Geochim Cosmochim Acta* 75:7132–7145.
- Beer J, et al. (1990) Use of ^{10}Be in polar ice to trace the 11-year cycle of solar activity. *Nature* 347:164–166.
- Berggren AM, et al. (2009) A 600-year annual ^{10}Be record from the NGRIP ice core, Greenland. *Geophys Res Lett* 36:L11801.
- Beer J, McCracken K, Steiger R (2012) *Cosmogenic Radionuclides: Theory and Applications in the Terrestrial and Space Environments* (Springer Science & Business Media, Berlin).
- Siegenthaler U, Heimann M, Oeschger H (1980) ^{14}C variations caused by changes in the global carbon cycle. *Radiocarbon* 22:177–191.
- Siegenthaler U (1983) Uptake of excess CO_2 by an outcrop-diffusion model of the ocean. *J Geophys Res Oceans* 88:3599–3608.