



Complete transcriptome assembly and annotation of a critically important amphipod species in freshwater ecotoxicological risk assessment: *Gammarus fossarum*



Domenico R. Caputo^{a,1}, Samuel C. Robson^{b,1}, Inge Werner^c, Alex T. Ford^{a,*}

^a Institute of Marine Sciences, School of Biological Sciences, University of Portsmouth, Ferry Road, Portsmouth PO4 9LY, UK

^b Centre for Enzyme Innovation, St. Michael's Building, University of Portsmouth, White Swan Road, Portsmouth PO1 2DT, UK

^c Swiss Centre for Applied Ecotoxicology, Eawag - EPFL, Überlandstrasse 133, 8600 Dübendorf, Switzerland

ARTICLE INFO

Handling Editor: Hefa Cheng

Keywords:

Gammarus fossarum

RNA sequencing

De novo assembly

Serotonin pathway

ABSTRACT

Because of their crucial role in ecotoxicological risk assessment, amphipods (Crustacea) are commonly employed as model species in a wide range of studies. However, despite their ecological importance, their genome has not yet been completely annotated and molecular mechanisms underlying key pathways, such as the serotonin pathway, in development of ecotoxicological biomarkers of exposure to neuroactive pharmaceuticals are still poorly understood. Furthermore, genetic similarities and discrepancies with other model arthropods (e.g., *Drosophila melanogaster*) have not been completely clarified. In this report, we present a new transcriptome assembly of *Gammarus fossarum*, an important amphipod species, widespread in Central Europe. RNA-Seq with Illumina HiSeq technology was used to analyse samples extracted from total internal tissues. We used the Trinity and Trinotate software suites for transcriptome assembly and annotation, respectively. The quality of this assembly and the affiliated targeted homology searches greatly enrich the molecular knowledge on this species. Because of the lack of publicly available molecular information on the serotonin pathway, we also highlighted sequence homologies and divergences of the genes encoding the serotonin pathway components of the well-annotated arthropod *D. melanogaster*, and Crustacea with the corresponding genes of our assembly. An inferior number of hits was found when running a BLAST analysis of both *D. melanogaster* and Crustacea mRNA sequences encoding serotonin receptors available in GenBank against the total assembly, compared to other serotonin pathway components. A lack of information on important components for serotonin biosynthesis and vesicle endocytosis (i.e., tryptophan hydroxylase and vesicular monoamine transporter) in Crustacea was also brought to light. Our results will provide an extensive transcriptional resource for this important species in ecotoxicological risk assessment and highlight the need for a more detailed categorization of neuronal pathways components in invertebrates.

1. Introduction

Identifying new sentinel species and acquiring molecular data for biomarker analyses are fundamental objectives in today's ecotoxicological research. Given the number and diversity of chemicals released into natural environments, the need for a deeper understanding of the molecular mechanisms taking place in native species in response to exposure to these substances is increasing more and more. Fortunately, the degree of specificity and above all the number of biomolecules that can be simultaneously analysed in a given species have exponentially increased over the last decade due to the use of

multi-omics platforms, such as next generation sequencing (NGS) technologies (Simmons et al., 2015). In research on aquatic species for instance, -omics platforms have allowed the development of new biomarkers of exposure to a considerably higher number of chemicals, allowing an even more precise description of the molecular effects in response to the exposure to a range of ecologically dangerous substances (Hauser-Davis et al., 2012; Jayapal, 2012; Barrera and Ariza, 2017; Poynton et al., 2018). With particular regard to freshwater species, numerous studies have been carried out to evaluate the molecular effects of a large spectrum of substances unremoved by wastewater treatment plants (WWTPs). These include endocrine disruptor

* Corresponding author at: Institute of Marine Sciences, School of Biological Sciences, University of Portsmouth, Ferry Road, Portsmouth PO4 9LY, UK.
E-mail address: alex.ford@port.ac.uk (A.T. Ford).

¹ These authors contributed equally to this work.

compounds (Bahamonde et al., 2014; Schneider et al., 2015; Gouveia et al., 2018), pharmaceuticals (Sanchez et al., 2011; Pascoe et al., 2003; Brandão et al., 2013), metals (Atli and Canli, 2007; Gismondi et al., 2017; Lebrun et al., 2017) and polychlorinated biphenyls (Rylander et al., 1998; Leroy et al., 2010) on various species of fish and invertebrates. Other important studies focusing on acquiring genomic sequences or applying multiple omics platforms to elucidate unclear biological mechanisms in species of ecotoxicological interest are also available (Trapp et al., 2016; Macher et al., 2017; Poynton et al., 2018). Because of their ecological relevance, invertebrates and in particular amphipods, are commonly employed as test organisms in ecotoxicological assessments. Among amphipods, the genus *Gammarus* represents the greatest number of epigeal freshwater species distributed throughout the Northern Hemisphere (Trapp et al., 2014a, 2014b). They are commonly used as model species in aquatic ecotoxicology for several reasons (Kunz et al., 2010). First, they are widespread and found throughout a large habitat range, where they often occur at high densities. Second, they occupy a large trophic repertoire as: herbivores, predators, and detritivores playing a major role in leaf-litter breakdown processes (Dangles and Guérol, 2001). They also constitute a food reserve for macroinvertebrates and fish. Gammarids can be easily maintained in the laboratory or used in field bioassays (Kunz et al., 2010), in which it is possible to assess the impact of pollutants by measuring molecular markers related to diverse modes of action, such as neurotoxicity (Xuereb et al., 2009), as well as by using life-history-trait reproductive features (Geffard et al., 2010). In particular, *Gammarus fossarum* has been shown as notably useful in ecotoxicology studies due to an intrinsic sensitivity to anthropogenic pollutants (Trapp et al., 2016; Wigh et al., 2017) and a spatial distribution across Central Europe (Straškraba, 1962; Meijering, 1971), making them ideal for freshwater risk assessment in the European geographical area. As a basic premise, a detailed elucidation of the molecular pathways is key to understanding the effects of pollutants on exposed organisms. In the literature, there are several studies that specifically applied sequencing platforms to *G. fossarum*. For example, Weiss et al. (2014) used sequencing of the genes encoding rRNA 16s and *cytochrome c oxidase 1* (CO1) for a taxonomic purpose. Trapp et al. (2014a, 2014b) used a proteogenomics strategy to generate a molecular report specifically on the reproductive tissues of female individuals of *G. fossarum* and Macher et al. (2017) sequenced the complete mitochondrial genome of this amphipod species. However, the lack of publicly available molecular information on amphipod physiological pathways compared to other model species is still clear. This is particularly true when considering neurological pathways, especially the molecular processes at the serotonergic synapse (Wu and Cooper, 2012). Behavioural studies that use pharmaceuticals acting on the serotonin pathway (e.g., antidepressants) have demonstrated a high pleiotropism of this neurotransmitter in invertebrates. In particular, feeding (Yeoman et al., 1994), swimming (Satterlie and Norekian, 1995), beating of cilia (Gosselin, 1961), reproduction (Fong, 1998), egg laying (Muschamp and Fong, 2001) and other behaviours such as aggressive motivation and activity (Huber et al., 1997; Tierney et al., 2004) are among the main behaviours altered by the most commonly used antidepressants. Furthermore, studies focusing on the impact of antidepressants, especially selective serotonin re-uptake inhibitors (SSRIs) (e.g., fluoxetine, fluvoxamine and sertraline) on aquatic organisms have been increasing (Johnson et al., 2007; Minagh et al., 2009; Demeestere et al., 2010; Guler and Ford, 2010; Styrisshave et al., 2011) and in most cases, molecular information in *D. melanogaster* (e.g., orthologous genes and proteins) are used as a reference to investigate the metabolic pathways in other arthropods, such as amphipods (Bossus et al., 2014; Trapp et al., 2014a, 2014b; Poynton et al., 2018). Therefore, a more detailed knowledge of the molecular processes taking place in the serotonergic synapse and a clear elucidation of the genetic differences of the main components (i.e., enzymes for serotonin biosynthesis, synaptic re-uptake and receptors) compared to other well-annotated arthropods such

as *D. melanogaster*, will have a highly positive impact on the development of new ecotoxicological biomarkers. This aspect becomes even more important when considering the historical contribution of crustaceans in the synaptic physiology research. For instance, the presence of many parallels found in central synaptic physiology between crustaceans and vertebrates (Wu and Cooper, 2012) and the amount of knowledge about the basic principles underlying the generation, maintenance, and modulation of rhythmically active behaviours (e.g., walking, chewing and breathing) in humans, gained through studies on crustaceans (Selverston and Moulins, 1987; Harris-Warrick et al., 1992; Marder et al., 1995; Selverston et al., 1998; Nusbaum et al., 2001; Skiebe, 2001; Cooke, 2002; Fénelon et al., 2003; Selverston, 2005; Selverston and Ayers, 2006; Marder and Bucher, 2007; Stein, 2009; Christie et al., 2010; Blitz and Nusbaum, 2011; Christie, 2011; Dickinson et al., 2016). While much work has focused on examining the behavioural effects of neuroactive drugs active on the serotonergic synapse in invertebrates, commonly released in aquatic environments thorough wastewaters, (De Lange et al., 2006; Guler and Ford, 2010; Bossus et al., 2014), many physiological and molecular details remain largely unknown. Although there are studies that have applied high-throughput omics technologies on various species of amphipods (Short et al., 2014a, 2014b; Gouveia et al., 2017), the genomes of many organisms belonging to this species have not yet been completely annotated. To our knowledge the only study applying a next-generation sequencing approach, in order to give an overall report on the *G. fossarum* transcriptome was performed by Cogne et al., (2019). The authors sequenced the transcriptome of 7 different taxonomic groups of *G. fossarum* and combined the sequencing data with a high-throughput proteomics analysis, in order to provide a broad molecular dataset on this amphipod. Our report will significantly add to the gene discovery work on this critically important species in ecotoxicological risk assessment as well as providing an overview on the publicly available sequence resources on the serotonin pathway in the sub-species A of *G. fossarum*. In this study, we employed a *de novo* transcriptome assembly approach to identify functionally relevant transcripts and explored the serotonin pathway.

2. Materials and methods

2.1. Sampling

The sampling was performed in mid-September 2017 at the Eulach river. The sampling site is located in Elgg, Switzerland at the following geographic coordinates: 47°30'04.23"N – 8°51'09.40"E. *G. fossarum* individuals were collected from beneath stones and leaves at the bottom of the stream, using a standard kick-net method. A net with 1 mm mesh size was used. The amphipods were removed from the net using forceps and sorted, separating the species of interest from leaves and other invertebrates. The animals were placed into 10 L buckets containing stream water and quickly transported to the laboratory where they were placed in controlled conditions. The amphipods were placed in glass tanks filled with continuously aerated stream water (20 cm depth). Incubation conditions were 16 ± 2 °C with a 12/12 light-dark cycle (Blarer and Burkhardt-Holm, 2016). Gammarids were fed *ad libitum* with alder leaves (*Alnus glutinosa*) collected at the sampling site. After a 24 h period, RNA was extracted.

2.2. Total RNA extraction

Total RNA was extracted from *G. fossarum* total internal tissues using RNeasy® Mini Kit (Qiagen, Hombrechtikon, Switzerland), following manufacturer's instructions. A total of 100 amphipod dissections were conducted, but in order to increase the RNA yield 5 independent amphipods were pooled per replicate, resulting in a total of 20 distinct pools. Each pool was considered an independent biological replicate for downstream analyses. Sampling was conducted independently for both

males and females (10 male pools and 10 female pools). Fresh amphipods were anaesthetised for 10–15 min in a 5% (v/v) clove oil solution prepared in water and washed in DEPC water (Sigma-Aldrich, Shnellendorf, Germany) prior to dissection, to remove any residual debris. Dissections were performed under a stereo-binocular (x3–4 magnification; SZ2 –ILST, Olympus), using stainless steel forceps. Heads were removed from the body, allowing an easier removal of the internal tissues. Internal tissues were washed in DEPC water and placed in 1.5 mL tubes previously cooled. Tubes were quickly snap-frozen in liquid nitrogen and placed at -80°C until RNA extractions. For each tube, one pre-treated stainless-steel bead (Qiagen, Hilden, Germany) and 350 μL of lysis buffer plus 10 μL β -mercaptoethanol (Sigma-Aldrich, Buchs, Switzerland) were added and the samples were immediately placed into the adaptors of a Tissue Lyser II[®] machine (Qiagen, Hilden, Germany). Stainless-steel beads were previously subject to two treatments of 15% (v/v) H_2O_2 and 70% (v/v) ethanol washings followed by 20 min of UV irradiation, to remove any potential chemical/biological contamination. Three 20 sec mechanical stirring cycles at 30 Hz speed were performed, in order to disrupt the tissues and homogenize the cell suspension. The lysed tissue samples were centrifuged at full speed at room temperature for 3 mins to separate the cell debris from the supernatant and the supernatants were transferred to fresh tubes. Total RNA was then extracted from *G. fossarum* tissue samples. RNA concentrations and purity were assessed measuring the absorbance at 230, 260 and 280 nm using Nanodrop ND-1000 (Witec, Littau, Switzerland). Finally, RNA integrity was checked using an Agilent 2100 Bioanalyzer (Agilent Technologies, Wahren, Germany) assay. All RNA samples displayed low background signal with sharp peaks corresponding to intact ribosomal RNA.

2.2.1. RNA sequencing

An Illumina TruSeq Stranded mRNA library kit was used to generate cDNA libraries for each of the 20 RNA samples, which were run on an Illumina HiSeq 2500 sequencer by GATC Biotech (Konstanz, Germany) to generate paired-end 150 bp reads.

2.2.2. Quality control

Quality control of the raw reads was performed using FastQC v0.11.7 (Andrews, 2015). Species-specific sequence purity was assessed by using a multiple genome alignment approach, by mapping reads against a database of different model species taken from the Ensembl database (Aken et al., 2017), a draft transcriptome for *Gammarus chevreuxi* (Truebano et al., 2016), and all RefSeq entries for *Gammarus fossarum* using MGA v1.4 (Hadfield and Eldridge, 2014). Read trimming was performed using Trim Galore v0.4.4 (Krueger, 2012) using the following parameters “–illumina -q 20 –stringency 5 -e 0.1 –length 20 –trim-n” to remove Illumina adapter sequence contamination, and to trim reads for ambiguous or low-quality base calls.

2.2.3. Assembly and annotation

Reads were combined across the data set and used to generate a putative transcriptome assembly using Trinity v2.5.1 (Grabherr et al., 2011) with parameters “–seqType fq –max_memory 100G –CPU 24 –min_contig_length 200 –min_kmer_cov 1 –SS_lib_type RF –verbose –full_cleanup”. Raw sequencing data were deposited in the NCBI Sequence Read Archive (SRA) and the complete transcriptome was deposited in the Transcriptome Shotgun Assembly (TSA) database. These data have been collected under BioProject accession code PRJNA556212. Unique transcript sequences were clustered into potential alternatively spliced isoform groups and paralogous “genes”. TransDecoder v5.0.2 (Haas et al., 2013) was run using default parameters to identify open reading frames (ORF) of 100 amino acids or more within transcripts, and putative protein amino acid sequences were produced. Transcripts were annotated against the Universal Protein Knowledge Base (UniProtKB) SwissProt database (The UniProt Consortium, 2017) using BLAST (Altschul, 1990), either at the protein

level by taking the TransDecoder derived peptide sequence (using “blastp”) or from the translated nucleotide sequence directly if no ORF was identified (using blastx). Additional annotation was performed against the Protein family (Pfam) database (Finn et al., 2013) using HMMER (Finn et al., 2011), Clusters of Orthologous Groups of proteins (eggNOG) database (Huerta-Cepas et al., 2015), the Kyoto Encyclopedia of Genes and Genomes (KEGG) database (Kanehisa, 2000), and Gene Ontology (GO) database (Ashburner et al., 2000). Results were collated into a single output table using Trinotate v3.02 (<http://trinotate.github.io/>). Transcriptome completeness was assessed by comparing the assembly against a database of metazoan universal single copy orthologs using BUSCO v2.0 (Simão et al., 2015).

2.2.4. *G. fossarum* sub-species assignment

A sequence alignment of the transcript sequences against the complete *G. fossarum* mitochondrial genome (Macher et al., 2017) was conducted using Blast2GO 5 Basic software (Conesa et al., 2005), in order to identify transcripts corresponding to mitochondrial genes that may allow for taxonomic assignment (i.e., rRNA 16S and CO1) (Müller, 2000; Weiss et al., 2014). The following parameters were used for the analysis:

- E-value: 1.0E^{-3}
- Number of blast hits: 20
- Word size: 11
- HSP length cutoff: 33

BLAST was then used to align putative mitochondrial transcripts against the NCBI non-redundant database (Altschul et al., 1990) for “amphipods – taxid:6821”, to assign both *G. fossarum* sub-species and CO1 type.

2.3. Serotonin pathway survey

Serotonin-specific nucleotide sequences were identified from GenBank (Benson et al., 2017) and were filtered to identify serotonin-specific mRNAs for enzymes, transporters and receptors, belonging to either *D. melanogaster* or Crustacea. In cases where multiple transcript variants were identified, all were considered, including *in silico* predicted transcripts. cDNA and single exon sequences were excluded. Genes encoding serotonin non-specific components (i.e., non-specific ion channels, G proteins, cytochrome p450 enzymes, receptors specific for other neurotransmitters) were not included and their corresponding boxes in Fig. 3 were left blank. A BLAST search for each nucleotide sequence was conducted against our transcriptome using BLAST2GO (Conesa et al., 2005) with parameters as described above to identify *G. fossarum*-specific serotonin genes. Fig. 1 shows a schematic overview of the workflow used, including both laboratory and bioinformatics procedures.

3. Results

3.1. *G. fossarum* de novo transcriptome assembly

Sequencing generated a total of 325,393,762 paired end reads across the 20 samples. Base calling quality was excellent across the entire data set, with the vast majority of reads showing average quality scores greater than 30. Mapping of reads against a range of different target genomes identified similar mapping rates for each of the 20 samples, with around 50% of reads mapping to either *G. fossarum*, *G. pulex* or *E. marinus*, whilst the remaining reads importantly did not map to any other model organism indicating no sign of cross-contamination. Trimming of reads to remove Illumina adapters, ambiguous base calls and poor-quality bases resulted in the loss of only 0.1% of reads. In total, 324,958,898 quality-trimmed reads were used for *de novo* transcriptome assembly. The final assembly consisted of 680,840




Fig. 1. Overview of workflow. Schematic flowchart showing the molecular biology and bioinformatic methods and software employed in this study. Blue steps are wet-lab procedures, red steps are *in silico* and black steps represent the outputs. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

Table 1

Summary statistics of *G. fossarum* transcriptome assembly. The table shows the main parameters describing the overall structure and quality of the assembly.

Trinity outputs	
Total Trinity 'genes'	407,060
Total Trinity transcripts	680,840
Percent GC	44.68%
Longest contig	31,653 bp
Shortest contig	100 bp
Greater than 10 Kb	352
Greater than 5 Kb	5,389
Greater than 2 Kb	94,819
Transcript contig N10	4,671 bp
Transcript contig N20	3,075 bp
Transcript contig N30	2,184 bp
Transcript contig N40	1,531 bp
Transcript contig N50	1,026 bp
Median contig length	324 bp
Average contig	628.1 bp
Total assembled bases	427,679,404

transcripts, clustered into 407,060 genes by the Butterfly portion of the Trinity algorithm. The transcriptome consisted of a total of 427,679,404 bp and showed a GC content of 44.68%. Transcripts ranged in size from 100 bp to 31,653 bp, with a median transcript size of 628.16 bp and an N50 of 1,026 bp (Table 1). The Ex90N50 (the N50 score for the transcripts accounting for 90% of the total normalized expression data) was 1922 bp and is accounted for by only 47,563 (6.8%) of the transcripts. Mapping of reads to this assembly showed a mapping rate of 65.7%. Transcriptome completeness was assessed by comparing against a database of 978 metazoan universal single-copy orthologs using BUSCO (Simão et al., 2015). In this assembly, 942 (96.4%) of the universal single-copy orthologs were present in a complete form, with 335 (34.3%) showing a single copy and 607 (62.1%) showing 2 or more copies. 19 (1.9%) were found in a fragmented form whilst only 17 (1.7%) were missing. This suggests that this assembly represents a very complete transcriptome for *G. fossarum*. Annotation against the UniProt/Swissprot database identified candidate hits for only 80,476 (11.8%) of the transcripts. However, 53.5% of transcripts with an identified open-reading frame (ORF) showed a hit, whilst only 5.6% of transcripts with no ORF showed a hit. The vast majority of non-

ORF transcripts (78.7%) were shorter in length than 500 bp (Fig. 2). This suggests that the majority of non-annotated non-coding transcripts are likely a result of fragmented RNA. Annotation for all transcripts identified in this assembly can be seen in Table S1 (Supplementary Information).

3.2. *G. fossarum* sub-species assignment

Previous studies have reported strong genetic differences among populations of *G. fossarum* (Siegismund, 1988; Siegismund and Müller, 1991; Müller, 2000). Although this amphipod has been a subject of controversial debate in taxonomy for decades (e.g., Roux, 1971; Goedmakers, 1972; Jazdzewski, 1977; Goedmakers, 1980; Scheepmaker and van Dalfsen, 1989; Pinkster and Scheepmaker, 1994; Müller, 1998; Westram et al., 2011), to date at least 3 main sub-species have been identified based on nucleotide differences within the mitochondrial gene encoding rRNA 16S (Müller, 2000). An additional taxonomic system in *G. fossarum* based on the differences in the gene encoding cytochrome oxidase 1 (CO1) was proposed by Weiss et al., (2014). In this study, a BLAST analysis of the transcriptome assembly was conducted against the complete *G. fossarum* mitochondrial genome (Macher et al., 2017). Putative mitochondrial transcript sequences from our assembly were aligned against the amphipod NCBI nucleotide database using BLAST in order to identify transcripts corresponding to rRNA 16S and CO1. Among the 20 putative mitochondrial transcripts, one (TRINITY_DN95725_c7_g2_i2) showed a hit against “*Gammarus fossarum* type A mitochondrial partial 16S rRNA gene, haplotype A14” suggesting that these data are specific to sub-species assignment (*G. fossarum* A) (Müller, 2000) (Table 2). Additionally, another transcript (TRINITY_DN93353_c3_g3_i1) showed a hit against “*Gammarus fossarum* isolate Gfos_47 cytochrome oxidase subunit 1 (COI) gene, partial cds; mitochondrial” and “*Gammarus fossarum* isolate Gfos_45 cytochrome oxidase subunit 1 (COI) gene, partial cds; mitochondrial” (Table 2). One other transcript (TRINITY_DN93353_c3_g1_i3) also showed hits against CO1, but from the amphipod taxa *Chiltoniidae*.

3.3. Serotonin pathway survey

To further test the depth and accuracy of this dataset, we aimed to identify serotonin-related transcripts by comparing against known serotonin-related genes from the NCBI database GenBank (Benson et al.,




Fig. 2. Annotation rates for both the transcripts with an ORF and with no ORF found. The plot shows the number of transcripts with an ORF (right) and with no ORF (left) that show a hit against the UniProt/Swissprot database. The transcripts were ordered by their length (x-axis).




Fig. 3. KEGG serotonergic synapse diagram. mRNA sequences belonging to *D. melanogaster* and Crustacea for each of the serotonin-specific enzymes, transporters and receptors were identified from GenBank. Sequence alignment of these nucleotide sequences against the transcriptome assembly was conducted using BLAST to identify putative serotonin-specific enzymes in the *G. fossarum* transcriptome. Red: at least one hit in both *D. melanogaster* and Crustacea. Blue: at least one hit in Crustacea. Brown: at least one hit in *D. melanogaster*. Green: at least one hit in human. Grey: no hit. Blank: not serotonin-specific gene. *: Results refer to 5-HT_{1A} only. **: Results refer to 5-HT_{4,6} only. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

2017). Given the well-known effects of exposure to antidepressants (a pharmaceutical class specifically acting on the serotonin pathway) on aquatic organisms (Guler and Ford, 2010; Styrisshave et al., 2011; Bossus et al., 2014; Chen et al., 2018), a detailed exploration of these

specific transcripts was of interest from an ecotoxicological perspective. *D. melanogaster* is often used as a reference to investigate the metabolic pathways in other arthropods, such as amphipods (Bossus et al., 2013; Trapp et al., 2014a, 2014b; Poynton et al., 2018). However, mRNA

Table 2
G. fossarum sub-type assignment BLAST parameters. A BLAST analysis of the complete G. fossarum mitochondrial genome was conducted against the total assembly, in order to identify the transcripts corresponding to rRNA 16S and CO1 genes. The table shows the BLAST parameters of the best hits obtained when re-BLASTing the transcripts coding rRNA 16S and CO1 genes in the NCBI database.

Transcript ID	Max score	Total score	Query cover	E-value	% identity	Best hit description	Accession
TRINITY_DN95725_c7_g2_i2 *	725	725	17%	0.0	100%	G. fossarum type A mitochondrial partial 16S rRNA gene, haplotype A14	AJ269600.1
TRINITY_DN93353_c3_g3_i1	1110	1110	73%	0.0	100%	Gammarrus fossarum isolate Gfos_47 cytochrome oxidase subunit 1 (COI) gene, partial cds; mitochondrial	MG986797.1
	1110	1110	73%	0.0	100%	Gammarrus fossarum isolate Gfos_45 cytochrome oxidase subunit 1 (COI) gene, partial cds; mitochondrial	MG986795.1

* Top 2 hits for TRINITY_DN95725_c7_g2_i2 were “Gammarrus fossarum mitochondrial, complete genome” and “Gammarrus fossarum isolate NEU07 16S ribosomal RNA gene, partial sequence; mitochondrial”. The table shows the third hit, which was useful for the sub-species assignment (G. fossarum A), according to Müller (2000) classification.

sequences encoding important serotonin receptors (e.g., 5-HT_{3,4,5,6}) were not available in GenBank for neither *D. melanogaster* nor Crustacea, so human-derived sequences were also included in the analysis (green boxes in Fig. 3). Fig. 3 shows the reference pathway of the serotonergic synapse from the KEGG database (Kanehisa & Goto, 2000). Table S2 (Supplementary Information) shows the details of this analysis, including the number of BLAST hits for each transcript, the GenBank accession code of the sequence matches, and the transcript ID.

3.3.1. 5-HT biosynthesis components

Serotonin is synthesized from the amino acid tryptophan in two enzymatic steps: hydroxylation of tryptophan by the neuronal form of tryptophan hydroxylase (TPH), which generates 5-hydroxytryptophan (5-HTP), and conversion to 5-hydroxytryptamine (5-HT) by the enzyme DOPA-decarboxylase (DDC) (Daubert and Condron, 2010). No mRNA sequences encoding TPH were available in GenBank for Crustacea, but 3 out of 4 sequences encoding this enzyme in *D. melanogaster* showed a putative hit in the *G. fossarum* transcriptome. No hit was found for the enzyme DDC, represented in GenBank by 3 and 2 mRNA sequences for *D. melanogaster* and Crustacea, respectively.

3.3.2. 5-HT transport proteins

After synthesis, the vesicular monoamine transporter (VMAT) transports 5-HT into vesicles for storage. Upon vesicle fusion with the plasma membrane, 5-HT is released where it interacts with autoreceptors located on the releasing cell or heteroreceptors, serotonin receptors located on other cell types. The serotonin transporter (SERT) transports 5-HT back into the releasing cell where it is likely to be re-packaged for release by VMAT or degraded by monoamine oxidase A (MAO-A) located on the outer mitochondrial membrane (Daubert and Condron, 2010). Similarly to TPH, no mRNA sequences encoding the vesicle monoamine transporter VMAT were found in Crustacea. However, each of the 4 sequences encoding this carrier protein found for *D. melanogaster* gave a hit with at least one transcript. mRNA sequences encoding the re-uptake protein SERT were found for both *D. melanogaster* and Crustacea and all showed hits with at least one transcript (Fig. 3).

3.3.3. 5-HT receptors

To date, at least fourteen different serotonin receptor subtypes have been identified and are grouped into seven families in human (5-HT₁–5-HT₇) (Daubert and Condron, 2010). Although some of these receptors are present in different isoforms in *D. melanogaster* and Crustacea (Wu and Cooper, 2012), they mediate the whole range of serotonin's effects and are distributed throughout the body (Daubert and Condron, 2010). All of the serotonin receptors are G-protein-coupled receptors, except for the 5-HT₃ ligand-gated ion channel (Nichols and Nichols, 2008). Compared to other components of the serotonergic synapse, fewer hits were found against *D. melanogaster* and Crustacea mRNA sequences encoding 5-HT receptors. In fact, of all of the presynaptic receptors, 5-HT_{1A} was the only one that gave at least one hit in Crustacea. In contrast, 5-HT_{1B}, 5-HT₂, 5-HT_{2A}, 5-HT_{5a} and 5-HT₇ gave no hits for Crustacea nor for *D. melanogaster* (Fig. 3). No mRNA sequences were found in GenBank for 5-HT_{1C}, 5-HT_{1F} and 5-HT₃ for *D. melanogaster* nor Crustacea. Interestingly, 5-HT₃, 5-HT₄ and 5-HT₆ gave at least one hit when using human sequences (Fig. 3).

3.3.4. 5-HT inactivation enzyme

An additional serotonin inactivation mechanism is mediated by the enzyme monoamine oxidase A (MAO-A). This enzyme is located on the outer mitochondrial membrane and is responsible for degradation of 5-HT at the presynaptic terminal (Daubert and Condron, 2010). Surprisingly, no mRNA sequences were found in GenBank for *D. melanogaster* and none of the 4 sequences found for Crustacea gave hits.

4. Discussion

Much has changed in recent years, and researchers are now more aware of the importance of obtaining detailed molecular information in species of ecotoxicological interest (Simmons et al., 2015). The search for new molecular biomarkers for the evaluation of the status of natural habitats, as well as the study of specific pathways affected as a result of anthropogenic activities, has increased exponentially (Pascoe et al., 2003; Atli and Canli, 2007; Leroy et al., 2010; Sanchez et al., 2011; Brandão et al., 2013; Bahamonde et al., 2014; Schneider et al., 2015; Gismondi et al., 2017; Lebrun et al., 2017; Gouveia et al., 2018).

In parallel, the use of “-omics” platforms has greatly increased the depth of molecular analyses, allowing the acquisition of data from hundreds of thousands of molecules simultaneously. This provides a valuable resource for ecotoxicological research (Simmons et al., 2015). Particular attention has been given to a sub-group of Crustacea, amphipods, due to their sensitivity to aquatic pollutants (Trapp et al., 2015; Wigh et al., 2017) and their central role in the freshwater food web (Dangles and Guérol, 2001). Furthermore, the potential of amphipods to bioaccumulate toxic substances is likely among the main reasons for their crucial role in ecotoxicological research. Munz et al., (2018) found higher concentrations of frequently detected risk-driving substances (i.e., various classes of pharmaceuticals, corrosion inhibitors, biocides, pesticides and personal care products) in *Gammarus fossarum* and *Gammarus pulex* sampled downstream of several Swiss WWTPs compared to upstream sites. In a recent study by Müller et al. (2019), the authors performed a sampling of *G. pulex* amphipods from several shores of the United Kingdom and assessed the potential presence of various dangerous chemicals within them. Surprisingly, illicit drugs such as cocaine and ketamine, as well as banned pesticides and pharmaceuticals, were found. However, the lack of molecular information on these species still represents a limiting factor and a more detailed genomic annotation is fundamental to highlighting homologies and discrepancies compared with other model organisms. Although some progress has already been made, for instance, the parallels to vertebrate central synaptic physiology of phenomenon described at Crustacea neuromuscular junctions (Wu and Cooper, 2012), there is still a lot of work to be done.

4.1. *G. fossarum* sub-species assignment

A correct sub-type assignment of the amphipod *G. fossarum* is essential. In fact, the genetic differences between the sub-types are considered strong enough to prevent crossbreeding in a natural setting (Müller, 1998; Wiese, 2013). Although Pinkster and Scheepmaker (1994) were able to obtain *G. fossarum* F1 juveniles from *ex-situ* crossbreeding experiments, Müller, the author who differentiated *G. fossarum* into 3 distinct cryptic species (A, B, C) (Müller, 1998, 2000), found out that sub-types A and B are to be considered reproductively isolate in a natural setting (Müller, 1998). An additional taxonomic system in *G. fossarum*, based on the differences in the faster evolving gene encoding *cytochrome oxidase 1* (CO1) was proposed by Weiss et al. (2014). In this study, we have attributed the *G. fossarum* sub-type A, haplotype A14 to our species using a BLAST analysis of our assembly against the complete *G. fossarum* mitochondrial genome (Macher et al., 2017). Although the BLAST analysis did not allow an unambiguous assignment of the CO1 type, it revealed CO1-45 and CO1-47 as most probable CO1 types of the amphipod subject of study.

4.2. Serotonin pathway survey

Considering that the historical contribution of Crustacea in synaptic physiology is unsurpassed (Atwood, 1976; Wiese, 2013) and much is still unclear concerning the synaptic molecular mechanisms within this taxon (Wu and Cooper, 2012), we also carried out a manual annotation of the genes encoding the serotonin-related molecular components

present in the final assembly. We focused on this particular neurotransmitter because of the lack of general molecular information in invertebrates in contrast with the wide range of documented behavioural and transcriptional effects of antidepressants released in aquatic environments (Nentwig, 2007; Guler and Ford, 2010; Bossus et al., 2014; Ford and Fong, 2016; Estévez-Calvar et al., 2017). The acquisition of further molecular details underlying the serotonin pathway in amphipods will also be useful in the field of invertebrate neurobiology. In fact, it has been shown that, similarly to neuroactive drugs, some parasite species (e.g., trematodes) are able to manipulate their invertebrate host's behaviour through modification of monoamine pathways, such as the serotonin pathway (Guler et al., 2015). We therefore aimed to use BLAST to identify putative candidate genes from our assembly that match serotonin-specific mRNA sequences available in GenBank for *D. melanogaster*, a widely used arthropod for genomic annotation (Bossus et al., 2013; Trapp et al., 2014a, 2014b; Poynton et al., 2018), and Crustacea.

Interestingly, the 5-HT transporter SERT was the only component of this pathway to show matches from both *D. melanogaster* and Crustacea within our assembly. Although further investigations will be needed to confirm this, we speculate that the re-uptake carriers at synaptic level might have a higher rate of evolutionary conservation compared to other components of the serotonergic synapse. In comparison, no mRNA sequences encoding the degradation enzyme MAO-A were found in GenBank for *D. melanogaster*, and none of the sequences found for Crustacea gave hits against our assembly.

Surprisingly, no mRNA sequences encoding the first enzyme of the serotonin biosynthetic chain (TPH) were found in GenBank for Crustacea, but all splicing variants of this enzyme in *D. melanogaster* sequences gave at least one hit against our assembly, highlighting the homology of this enzyme in *D. melanogaster* and amphipods. However, none of the splicing variants found for the second serotonin biosynthesis enzyme in *D. melanogaster*, DOPA-decarboxylase, gave hits. Although two predicted nucleotide sequences encoding DDC enzyme were found for Crustacea, neither gave hits in the *G. fossarum* assembly. At present, we are unable to clarify whether this enzyme may have additional isoforms in Crustacea not present in GenBank, or whether sequence differences within this taxon are stronger than expected.

When looking at the serotonin receptors sequences, 5-HT_{1A} was the only receptor that gave at least one hit against Crustacea, but not *D. melanogaster* sequences. 5-HT_{1B}, 5-HT₂, 5-HT_{2A} and 5-HT_{5a} gave no hits against either the *D. melanogaster* or the Crustacea sequences. No mRNA sequences were found in GenBank for 5-HT_{1C-F} or 5-HT₃ for either *D. melanogaster* and Crustacea. 5-HT₇ gave no hits against *D. melanogaster* and Crustacea sequences.

It has previously been shown that the classification of serotonin receptors in humans is not totally applicable to invertebrates (Wu and Cooper, 2012). For instance, 5-HT₄ and 4-HT₇ are shown to have different splicing variants (Hoyer, Hannon and Martin, 2002) and 5-HT₂ can have different RNA-edited isoforms (Burns et al., 1997; Niswender et al., 1998). This analysis highlights the fact that further studies on a more precise classification of receptors for serotonin in invertebrates is strongly needed, considering the well documented effects that pharmaceutical waste targeting the serotonin pathway can have on aquatic species (Guler and Ford, 2010; Styryshave et al., 2011; Bossus et al., 2014; Chen et al., 2018). 5-HT_{1A} was the only receptor that gave at least one hit when using Crustacea mRNA sequences found in GenBank. Despite no sequences being found for 5-HT₃, we did identify hits in our assembly for human 5-HT₄ and 5-HT₆ mRNA sequences. Therefore, the lower hit-rate observed for serotonin receptors compared to other molecular components of the pathway might be due to a scarcity of genetic information in Crustacea rather than a strong evolutionary divergence of the 5-HT receptor sequences of this taxon.

Taken together, these findings reveal that molecular comparisons between the serotonin pathway of different organisms, or even within Crustacea taxon, may not be possible for all components. However, the

amount of genetic information available still represents a limiting factor. Given the importance of the serotonin pathway in ecotoxicological research, further studies on the classification of 5-HT receptors in invertebrates as well as their pharmaceutical targets will be fundamental.

5. Conclusions

The amphipod species *G. fossarum* represents an important organism in ecotoxicological risk assessment, and yet there remains a significant lack of genomic information. To further add to our current understanding, we used RNA sequencing and *de novo* assembly to generate a complete transcriptome of *G. fossarum* - type A. Our dataset will provide an additional resource of genomic information in this poorly annotated species and will represent a reference source for further and more focused molecular analysis on this and other amphipod species. Given the importance of the serotonin pathway in behavioural and neurological studies on invertebrates (Wu and Cooper, 2012; Guler et al., 2015) an exploration of the publicly available mRNA sequences involved in this pathway in *D. melanogaster* and Crustacea was also conducted. An inferior number of hits was found when running sequence alignment of our assembly against both *D. melanogaster* and Crustacea mRNA sequences encoding serotonin receptors available in GenBank, compared to other serotonin pathway components. This may indicate a lack of publicly available sequence information on important components of the serotonin pathway in crustaceans, for instance tryptophan hydroxylase and vesicular monoamine transporter, suggesting the need for a more extensive categorization of neuronal pathway components in key invertebrate species and important model species for both ecotoxicological and neurological studies.

Declaration of Competing Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Acknowledgements

We thank Dr. Melanie Fischer (aQuaTox - Solutions, Dübendorf, Switzerland), Andrea Schifferli and Thomas Bucher (Swiss Centre for Applied Ecotoxicology, Dübendorf, Switzerland) for their valuable technical support in sampling and RNA extractions. This project was financially supported by a bursary awarded by the Swiss Centre for Applied Ecotoxicology (Switzerland) and the University of Portsmouth (Portsmouth, UK). Dr Robson was partially funded through an Expanding Excellence in England (E3) grant from Research England.

Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.envint.2019.105319>.

References

Aken, Bronwen L., Achuthan, Premanand, Akanni, Wasiu, Ridwan Amodé, M., Bernsdorff, Friederike, Bhai, Jyothish, Billis, Konstantinos, Carvalho-Silva, Denise, Cummins, Carla, Clapham, Peter, Gil, Laurent, Girón, Carlos García, Gordon, Leo, Hourlier, Thibaut, Hunt, Sarah E., Janacek, Sophie H., Juettemann, Thomas, Keenan, Stephen, Laird, Matthew R., Lavidas, Ilias, Maurel, Thomas, McLaren, William, Moore, Benjamin, Murphy, Daniel N., Nag, Rishi, Newman, Victoria, Nuhn, Michael, Ong, Chuang Kee, Parker, Anne, Patricio, Mateus, Riat, Harpreet Singh, Sheppard, Daniel, Sparrow, Helen, Taylor, Kieron, Thormann, Anja, Vullo, Alessandro, Walts, Brandon, Wilder, Steven P., Zadissa, Amonida, Kostadima, Myrto, Martin, Fergal J., Muffato, Matthieu, Perry, Emily, Ruffier, Magali, Staines, Daniel M., Trevanion, Stephen J., Cunningham, Fiona, Yates, Andrew, Zerbino, Daniel R., Flicek, Paul, 2017. Ensembl. Nucl. Acids Res. 45(D1), D635–D642.

Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J., 1990. Basic local alignment search tool. J. Mol. Biol. 215 (3), 403–410.

Andrews, S., 2015. A quality control tool for high throughput sequence data. 2010. URL: <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>.

Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K., Dwight, S.S., Eppig, J.T., Harris, M.A., 2000. Gene ontology: tool for the unification of biology. Nat. Genet. 25 (1), 25.

Atli, G., Canli, M., 2007. Enzymatic responses to metal exposures in a freshwater fish *Oreochromis niloticus*. Comp. Biochem. Physiol. Part C: Toxicol. Pharmacol. 145 (2), 282–287.

Atwood, H.L., 1976. Organization and synaptic physiology of crustacean neuromuscular systems. Progr. Neurobiol. 7, 291–391.

Bahamonde, P.A., Tetreault, G.R., McMaster, M.E., Servos, M.R., Martyniuk, C.J., Munkittrick, K.R., 2014. Molecular signatures in rainbow darter (*Etheostoma caeruleum*) inhabiting an urbanized river reach receiving wastewater effluents. Aquatic Toxicol. 148, 211–220.

Barrera, T.G., Ariza, J.L.G., 2017. New Trends in Aquatic Pollution Monitoring: From Conventional Biomarkers to Environmental Proteomics. In Environmental Problems in Marine Biology. CRC Press, pp. 166–187.

Benson, Dennis A., Cavanaugh, Mark, Clark, Karen, Karsch-Mizrachi, Ilene, Lipman, David J., Ostell, James, Sayers, Eric W., 2017. GenBank. Nucl. Acids Res. 45 (D1), D37–D42.

Blarer, P., Burkhardt-Holm, P., 2016. Microplastics affect assimilation efficiency in the freshwater amphipod *Gammarus fossarum*. Environ. Sci. Pollut. Res. 23 (23), 23522–23532.

Blitz, D.M., Nusbaum, M.P., 2011. Neural circuit flexibility in a small sensorimotor system. Curr. Opin. Neurobiol. 21 (4), 544–552.

Bossus, M.C., Guler, Y.Z., Short, S.J., Morrison, E.R., Ford, A.T., 2014. Behavioural and transcriptional changes in the amphipod *Echinogammarus marinus* exposed to two antidepressants, fluoxetine and sertraline. Aquatic Toxicol. 151, 46–56.

Brandão, F.P., Rodrigues, S., Castro, B.B., Gonçalves, F., Antunes, S.C., Nunes, B., 2013. Short-term effects of neuroactive pharmaceutical drugs on a fish species: biochemical and behavioural effects. Aquatic Toxicol. 144, 218–229.

Burns, C.M., Chu, H., Rueter, S.M., Hutchinson, L.K., Canton, H., Sanders-Bush, E., Emeson, R.B., 1997. Regulation of serotonin - 2C receptor G-protein coupling by RNA editing. Nature 387, 303–308.

Chen, H., Zeng, X., Mu, L., Hou, L., Yang, B., Zhao, J., Schlenk, D., Dong, W., Xie, L., Zhang, Q., 2018. Effects of acute and chronic exposures of fluoxetine on the Chinese fish, topmouth gudgeon *Pseudorasbora parva*. Ecotoxicol. Environ. Safety 160, 104–113.

Christie, A.E., 2011. Crustacean neuroendocrine systems and their signaling agents. Cell Tissue Res. 345, 41–67.

Christie, A.E., Stemmler, E.A., Dickinson, P.S., 2010. Crustacean neuropeptides. Cell Mol. Life Sci. 67, 4135–4169.

Cogne, Y., Degli-Esposti, D., Pible, O., Gouveia, D., François, A., Bouchez, O., Eché, C., Ford, A., Geffard, O., Armengaud, J., Chaumot, A., 2019. De novo transcriptomes of 14 gammarid individuals for proteogenomic analysis of seven taxonomic groups. Scientific Data 6 (1), 1–7.

Conesa, A., Götz, S., García-Gómez, J.M., Terol, J., Talón, M., Robles, M., 2005. Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. Bioinformatics 21 (18), 3674–3676.

Cooke, I.M., 2002. Reliable, responsive pacemaking and pattern generation with minimal cell numbers: the crustacean cardiac ganglion. Biol. Bull. 202 (2), 108–136.

Dangles, O., Guérol, F., 2001. Linking shredders and leaf litter processing: insights from an acidic stream study. Int. Rev. Hydrobiol.: J. Covering Aspects Limnol. Mar. Biol. 86 (4–5), 395–406.

Daubert, E.A., Condron, B.G., 2010. Serotonin: a regulator of neuronal morphology and circuitry. Trends Neurosci. 33 (9), 424–434.

De Lange, H.J., Noordoven, W., Murk, A.J., Lüring, M.F.L.L.W., Peeters, E.T.H.M., 2006. Behavioural responses of *Gammarus pulex* (Crustacea, Amphipoda) to low concentrations of pharmaceuticals. Aquatic Toxicol. 78 (3), 209–216.

Demeestere, K., Petrović, M., Gros, M., Dewulf, J., Van Langenhove, H., Barceló, D., 2010. Trace analysis of antidepressants in environmental waters by molecularly imprinted polymer-based solid-phase extraction followed by ultra-performance liquid chromatography coupled to triple quadrupole mass spectrometry. Anal. Bioanal. Chem. 396 (2), 825–837.

Dickinson, P.S., Qu, X., Stanhope, M.E., 2016. Neuropeptide modulation of pattern-generating systems in crustaceans: comparative studies and approaches. Curr. Opin. Neurobiol. 41, 149–157.

Estévez-Calvar, N., Canesi, L., Montagna, M., Faimali, M., Piazza, V., Garaventa, F., 2017. Adverse effects of the SSRI antidepressant sertraline on early life stages of marine invertebrates. Mar. Environ. Res. 128, 88–97.

Fénelon, V., Le Feuvre, Y., Bem, T., Meyrand, P., 2003. Maturation of rhythmic neural network: role of central modulatory inputs. J. Physiol.-Paris 97 (1), 59–68.

Finn, R.D., Clements, J., Eddy, S.R., 2011. HMMER web server: interactive sequence similarity searching. Nucl. Acids Res. 39 (suppl_2), W29–W37.

Finn, R.D., Bateman, A., Clements, J., Coggill, P., Eberhardt, R.Y., Eddy, S.R., Heger, A., Hetherington, K., Holm, L., Mistry, J., Sonnhammer, E.L., 2013. Pfam: the protein families database. Nucl. Acids Res. 42 (D1), D222–D230.

Fong, P.P., 1998. Zebra mussel spawning is induced in low concentrations of putative serotonin reuptake inhibitors. Biol. Bull. 194, 143–149.

Ford, A.T., Fong, P.P., 2016. The effects of antidepressants appear to be rapid and at environmentally relevant concentrations. Environ. Toxicol. Chem. 35 (4), 794–798.

Geffard, O., Xuereb, B., Chaumot, A., Geffard, A., Biagianti, S., Noël, C., Abbaci, K., Garric, J., Charmantier, G., Charmantier-Daures, M., 2010. Ovarian cycle and embryonic development in *Gammarus fossarum*: application for reproductive toxicity assessment. Environ. Toxicol. Chem. 29 (10), 2249–2259.

Gismond, E., Thomé, J.P., Urien, N., Uher, E., Baiwir, D., Mazzucchi, G., De Pauw, E.,

- Fechner, L.C., Lebrun, J.D., 2017. Ecotoxicoproteomic assessment of the functional alterations caused by chronic metallic exposures in gammarids. *Environ. Pollut.* 225, 428–438.
- Goedmakers, A., 1972. *Gammarus fossarum* Koch, 1835: redescription based on Bijdragen tot de Dierkunde 42 (2), 125.
- Goedmakers, A., 1980. Microgeographic races of *Gammarus fossarum* Koch, 1836. *Crustaceana. Supplement* 216–224.
- Gosselin, R.E., 1961. The cilioexcitatory activity of serotonin. *J. Cell. Comp. Physiol.* 58, 17–25.
- Gouveia, D., Chaumot, A., Charnot, A., Almunia, C., François, A., Navarro, L., Armengaud, J., Salvador, A., Geffard, O., 2017. Ecotoxicoproteomics for aquatic environmental monitoring: First in situ application of a new proteomics-based multi-biomarker assay using caged amphipods. *Environ. Sci. Technol.* 51 (22), 13417–13426.
- Gouveia, D., Bonneton, F., Almunia, C., Armengaud, J., Quéau, H., Degli-Esposti, D., Geffard, O., Chaumot, A., 2018. Identification, expression, and endocrine-disruption of three ecdysone-responsive genes in the sentinel species *Gammarus fossarum*. *Scientific Reports* 8 (1), 3793.
- Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat. Biotechnol.* 29 (7), 644.
- Guler, Y., Ford, A.T., 2010. Anti-depressants make amphipods see the light. *Aquatic Toxicol.* 99 (3), 397–404.
- Guler, Y., Short, S., Etxabe, A.G., Sherhod, C.M., Kille, P., Ford, A.T., 2015. Impacts of a newly identified behaviour-altering trematode on its host amphipod: from the level of gene expression to population. *Parasitology* 142 (12), 1469–1480.
- Haas, B.J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P.D., Bowden, J., Couger, M.B., Eccles, D., Li, B., Lieber, M., MacManes, M.D., 2013. De novo transcriptome sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nat. Protocols* 8 (8), 1494.
- Hadfield, J., Eldridge, M.D., 2014. Multi-genome alignment for quality control and contamination screening of next-generation sequencing data. *Front. Genet.* 5, 31.
- Harris-Warrick, R.M., Marder, E., Selverston, A.I., Moulins, M., 1992. Dynamic Biological Networks: The Stomatogastric Nervous System. MIT press Cambridge.
- Hauser-Davis, R.A., de Campos, R.C., Zioli, R.L., 2012. Fish metalloproteins as biomarkers of environmental contamination. In: *Reviews of Environmental Contamination and Toxicology*, vol. 218. Springer, Boston, MA, pp. 101–123.
- Hoyer, D., Hannon, J.P., Martin, G.R., 2002. Molecular, pharmacological and functional diversity of 5-HT receptors. *Pharmacol. Biochem. Behav.* 71 (4), 533–554.
- Huber, R., Smith, K., Delago, A., Isaksson, K., Kravitz, E.A., 1997. Serotonin and aggressive motivation in crustaceans: altering the decision to retreat. *Proc. Natl. Acad. Sci.* 94 (11), 5939–5942.
- Huerta-Cepas, J., Szklarczyk, D., Forslund, K., Cook, H., Heller, D., Walter, M.C., Rattei, T., Mende, D.R., Sunagawa, S., Kuhn, M., Jensen, L.J., 2015. eggNOG 4.5: a hierarchical orthology framework with improved functional annotations for eukaryotic, prokaryotic and viral sequences. *Nucl. Acids Res.* 44 (D1), D286–D293.
- Jayapal, M., 2012. Integration of next-generation sequencing based multi-omics approaches in toxicogenomics. *Front. Genet.* 3, 88.
- Jazdzewski, K., 1977. Remarks on the morphology of *Gammarus fossarum* Koch, 1835, and *Gammarus kischineffensis* Schellenberg, 1937. *Crustaceana. Supplement*, pp. 201–211.
- Johnson, D.J., Sanderson, H., Brain, R.A., Wilson, C.J., Solomon, K.R., 2007. Toxicity and hazard of selective serotonin reuptake inhibitor antidepressants fluoxetine, fluvoxamine, and sertraline to algae. *Ecotoxicol. Environ. Safety* 72 (1), 128–139.
- Kanehisa, M., Goto, S., 2000. KEGG: kyoto encyclopedia of genes and genomes. *Nucl. Acids Res.* 28 (1), 27–30.
- Krueger, F., 2012. Trim Galore! A quality trimming tool for high throughput sequence data. Available online at: https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/.
- Kunz, P.Y., Kienle, C., Gerhardt, A., 2010. *Gammarus* spp. in aquatic ecotoxicology and water quality assessment: toward integrated multilevel tests. In: *Reviews of Environmental Contamination and Toxicology*, vol. 205. Springer, New York, NY, pp. 1–76.
- Lebrun, J.D., Uher, E., Fechner, L.C., 2017. Behavioural and biochemical responses to metals tested alone or in mixture (Cd-Cu-Ni-Pb-Zn) in *Gammarus fossarum*: From a multi-biomarker approach to modelling metal mixture toxicity. *Aquatic Toxicol.* 193, 160–167.
- Leroy, D., Haubruge, E., De Pauw, E., Thomé, J.P., Francis, F., 2010. Development of ecotoxicoproteomics on the freshwater amphipod *Gammarus pulex*: identification of PCB biomarkers in glycolysis and glutamate pathways. *Ecotoxicol. Environ. Safety* 73 (3), 343–352.
- Macher, J.N., Leese, F., Weigand, A.M., Rozenberg, A., 2017. The complete mitochondrial genome of a cryptic amphipod species from the *Gammarus fossarum* complex. *Mitochondrial DNA Part B* 2 (1), 17–18.
- Marder, E., Bucher, D., 2007. Understanding circuit dynamics using the stomatogastric nervous system of lobsters and crabs. *Annu. Rev. Physiol.* 69, 291–316.
- Marder, E., Christie, A.E., Kilman, V.L., 1995. Functional organization of cotransmission systems: lessons from small nervous systems. *Invertebrate Neurosci.* 1 (2), 105–112.
- Meijering, M.P.D., 1971. Die Gammarusfauna der Schlitzladerfließgewässer. *Arch. Hydrobiol.* 68 (4), 575–608.
- Miller, Thomas H., Ng, Keng Tiong, Bury, Samuel T., Bury, Sophie E., Bury, Nicolas R., Barrona, Leon P., 2019. Biomonitoring of pesticides, pharmaceuticals and illicit drugs in a freshwater invertebrate to estimate toxic or effect pressure. *Environ. Int.* 129, 595–606.
- Minagh, E., Hernan, R., O'Rourke, K., Lyng, F.M., Davoren, M., 2009. Aquatic ecotoxicity of the selective serotonin reuptake inhibitor sertraline hydrochloride in a battery of freshwater test species. *Ecotoxicol. Environ. Safety* 72 (2), 434–440.
- Müller, J., 1998. Genetic population structure of two cryptic *Gammarus fossarum* types across a contact zone. *J. Evol. Biol.* 11, 79–101.
- Müller, J., 2000. Mitochondrial DNA variation and the evolutionary history of cryptic *Gammarus fossarum* types. *Mol. Phylogenet. Evol.* 15 (2), 260–268.
- Munz, N.A., Fu, Q., Stamm, C., Hollender, J., 2018. Internal concentrations in gammarids reveal increased risk of organic micropollutants in wastewater-impacted streams. *Environ. Sci. Technol.* 52 (18), 10347–10358.
- Muschamp, J.W., Fong, P.P., 2001. Effects of the serotonin receptor ligand methiothepin on reproductive behaviour of the freshwater snail *Biomphalaria glabrata*: reduction of egg laying and induction of penile erection. *J. Exp. Zool.* 289, 202–207.
- Nentwig, G., 2007. Effects of pharmaceuticals on aquatic invertebrates. Part II: The antidepressant drug fluoxetine. *Arch. Environ. Contamin. Toxicol.* 52 (2), 163–170.
- Nichols, D.E., Nichols, C.D., 2008. Serotonin receptors. *Chem. Rev.* 108 (5), 1614–1641.
- Niswender, C.M., Sanders-Bush, E., Emeson, R.B., 1998. Identification and characterization of RNA editing events within the 5-HT_{2C} receptor. *Ann. New York Acad. Sci.* 861 (1), 38–48.
- Nusbaum, M.P., Blitz, D.M., Swensen, A.M., Wood, D., Marder, E., 2001. The roles of cotransmission in neural network modulation. *Trends Neurosci.* 24 (3), 146–154.
- Pascoe, D., Karntant, W., Müller, C.T., 2003. Do pharmaceuticals affect freshwater invertebrates? A study with the cnidarian *Hydra vulgaris*. *Chemosphere* 51 (6), 521–528.
- Pinkster, S., Scheepmaker, M., 1994. Hybridization experiments and the taxonomy of *Gammarus* (Amphipoda): a contribution to the understanding of controversial results. *Crustaceana* 129–143.
- Poynton, H.C., Hasenbein, S., Benoit, J.B., Sepulveda, M.S., Poelchau, M.F., Hughes, D.S., Murali, S.C., Chen, S., Glastad, K.M., Goodisman, M.A., Werren, J.H., 2018. The Toxicogenome of *Hyalella azteca*: a model for sediment ecotoxicology and evolutionary toxicology. *Environ. Sci. Technol.* 52 (10), 6009–6022.
- Roux, A.L., 1971. Les Gammarus du groupe pulex. Essai de systématique biologique. II. Quelques caractéristiques écologiques et physiologiques. *Arch. Zool. exp. gén* 112, 471–503.
- Rylander, L., Strömberg, U., Dyremark, E., Östman, C., Nilsson-Ehle, P., Hagmar, L., 1998. Polychlorinated biphenyls in blood plasma among Swedish female fish consumers in relation to low birth weight. *Am. J. Epidemiol.* 147 (5), 493–502.
- Sanchez, W., Sremski, W., Piccini, B., Palluel, O., Maillot-Marechal, E., Betoulle, S., Jaffal, A., Ait-Aissa, S., Brion, F., Thybaud, E., Hinfrey, N., 2011. Adverse effects in wild fish living downstream from pharmaceutical manufacture discharges. *Environ. Int.* 37 (8), 1342–1348.
- Satterlie, R.A., Norekian, T.P., 1995. Serotonergic modulation of swimming speed in the pteropod mollusc *Clione limacina*. III. Cerebral neurons. *J. Exp. Biol.* 198, 917–930.
- Scheepmaker, M., van Dalfsen, J., 1989. Genetic differentiation in *Gammarus fossarum* and *G. caparti* (Crustacea, Amphipoda) with reference to *G. pulex* in north-western Europe. *Bijdragen Tot De Dierkunde* 59, 127–139.
- Schneider, I., Oehlmann, J., Oetken, M., 2015. Impact of an estrogenic sewage treatment plant effluent on life-history traits of the freshwater amphipod *Gammarus pulex*. *J. Environ. Sci. Health, Part A* 50 (3), 272–281.
- Selverston, A.I., 2005. A neural infrastructure for rhythmic motor patterns. *Cell. Mol. Neurobiol.* 25 (2), 223–244.
- Selverston, A.I., Ayers, J., 2006. Oscillations and oscillatory behavior in small neural circuits. *Biol. Cybernet.* 95 (6), 537.
- Selverston, A., Elson, R., Rabinovich, M., Huerta, R., Abarbanel, H., 1998. Basic principles for generating motor output in the stomatogastric ganglion. *Ann. New York Acad. Sci.* 860 (1), 35–50.
- Selverston, A.I., Moulins, M., 1987. *The Crustacean Stomatogastric System*. Springer, Berlin Heidelberg.
- Short, S., Yang, G., Kille, P., Ford, A.T., 2014b. Vitellogenin is not an appropriate biomarker of feminisation in a crustacean. *Aquatic Toxicol.* 153, 89–97.
- Short, S., Yang, G., Guler, Y., Green Etxabe, A., Kille, P., Ford, A.T., 2014a. Crustacean intersexuality is feminization without demasculinization: implications for environmental toxicology. *Environ. Sci. Technol.* 48 (22), 13520–13529.
- Siegmund, H.R., 1988. Genetic differentiation in populations of the freshwater amphipods *Gammarus roeselii* and *Gammarus fossarum*. *Heredity* 109 (2), 269–276.
- Siegmund, H.R., Müller, J., 1991. Genetic structure of *Gammarus fossarum* populations. *Heredity* 66 (3), 419.
- Simão, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E.V., Zdobnov, E.M., 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* 31 (19), 3210–3212.
- Simmons, D.B., Benskin, J.P., Cosgrove, J.R., Duncker, B.P., Ekman, D.R., Martyniuk, C.J., Sherry, J.P., 2015. Omics for aquatic ecotoxicology: Control of extraneous variability to enhance the analysis of environmental effects. *Environ. Toxicol. Chem.* 34 (8), 1693–1704.
- Skiepe, P., 2001. Neuropeptides are ubiquitous chemical mediators: using the stomatogastric nervous system as a model system. *J. Exp. Biol.* 204 (12), 2035–2048.
- Stein, W., 2009. Modulation of stomatogastric rhythms. *J. Comp. Physiol. A* 195 (11), 989–1009.
- Stráškaba, M., 1962. Amphipoden der Tschechoslowakei nach den Sammlungen von Prof. Hrabě. I. Věstník Československé Zoologické Společnosti 26 (2), 117–145.
- Styrishave, B., Halling-Sørensen, B., Ingerslev, F., 2011. Environmental risk assessment of three selective serotonin reuptake inhibitors in the aquatic environment: a case study including a cocktail scenario. *Environ. Toxicol. Chem.* 30 (1), 254–261.
- The UniProt Consortium, 2017. UniProt: the universal protein knowledgebase. *Nucleic acids research* 45, no. D1 (2016), pp. D158–D169.
- Tierney, A.J., Greenlaw, M.A., Dams-O'Connor, K., Aig, S.D., Perna, A.M., 2004. Behavioral effects of serotonin and serotonin agonists in two crayfish species,

- Procambarus clarkii* and *Orconectes rusticus*. *Comp. Biochem. Physiol. Part A Mol. Integr. Physiol.* 139, 495–502.
- Trapp, J., Armengaud, J., Pible, O., Gaillard, J.C., Abbaci, K., Habtoul, Y., Chaumot, A., Geffard, O., 2014a. Proteomic investigation of male *Gammarus fossarum*, a freshwater crustacean, in response to endocrine disruptors. *J. Proteome Res.* 14 (1), 292–303.
- Trapp, J., Geffard, O., Imbert, G., Gaillard, J.C., Davin, A.H., Chaumot, A., Armengaud, J., 2014b. Proteogenomics of *Gammarus fossarum* to document the reproductive system of amphipods. *Mol. Cell. Proteom.* 13 (12), 3612–3625.
- Trapp, J., Almunia, C., Gaillard, J.C., Pible, O., Chaumot, A., Geffard, O., Armengaud, J., 2016. Proteogenomic insights into the core-proteome of female reproductive tissues from crustacean amphipods. *J. Proteom.* 135, 51–61.
- Truebano, M., Tills, O., Spicer, J.I., 2016. Embryonic transcriptome of the brackishwater amphipod *Gammarus chevreuxi*. *Mar. Genom.* 28, 5–6.
- Weiss, M., Macher, J.N., Seefeldt, M.A., Leese, F., 2014. Molecular evidence for further overlooked species within the *Gammarus fossarum* complex (Crustacea: Amphipoda). *Hydrobiologia* 721 (1), 165–184.
- Westram, A.M., Jokela, J., Baumgartner, C., Keller, I., 2011. Spatial distribution of cryptic species diversity in European freshwater amphipods (*Gammarus fossarum*) as revealed by pyrosequencing. *PLoS one* 6 (8), e23879.
- Wiese, K. (Ed.), 2013. *The Crustacean Nervous System*. Springer Science & Business Media.
- Wigh, A., Geffard, O., Abbaci, K., Francois, A., Noury, P., Bergé, A., Vulliet, E., Domenjoud, B., Gonzalez-Ospina, A., Bony, S., Devaux, A., 2017. *Gammarus fossarum* as a sensitive tool to reveal residual toxicity of treated wastewater effluents. *Sci. Total Environ.* 584, 1012–1021.
- Wu, W.H., Cooper, R.L., 2012. Serotonin and synaptic transmission at invertebrate neuromuscular junctions. *Exp. Neurobiol.* 21 (3), 101–112.
- Xuereb, B., Chaumot, A., Mons, R., Garric, J., Geffard, O., 2009. Acetylcholinesterase activity in *Gammarus fossarum* (Crustacea Amphipoda): intrinsic variability, reference levels, and a reliable tool for field surveys. *Aquatic Toxicol.* 93 (4), 225–233.
- Yeoman, M.S., Pieneman, A.W., Ferguson, G.P., Ter Maat, A., Benjamin, P.R., 1994. Modulatory role for the serotonergic cerebral giant cells in the feeding system of the snail, *Lymnaea*. I. Fine wire recording in the intact animal and pharmacology. *J. Neurophysiol.* 72, 1357–1371.