



## Hypolimnetic oxygen depletion rates in deep lakes: Effects of trophic state and organic matter accumulation

Thomas Steinsberger<sup>1</sup>, Robert Schwefel<sup>2,3</sup>, Alfred Wüest<sup>1,3</sup>, Beat Müller<sup>1\*</sup>

<sup>1</sup>Eawag, Swiss Federal Institute of Aquatic Science and Technology, Kastanienbaum, Switzerland

<sup>2</sup>UC Santa Barbara, Santa Barbara, California

<sup>3</sup>Physics of Aquatic Systems Laboratory, Margaretha Kamprad Chair, École Polytechnique Fédérale de Lausanne, Institute of Environmental Engineering, Lausanne, Switzerland

### Abstract

This study investigated the consumption of oxygen ( $O_2$ ) in 11 European lakes ranging from 48 m to 372 m deep. In lakes less than ~100 m deep, the main pathways for  $O_2$  consumption were organic matter (OM) mineralization at the sediment surface and oxidation of reduced compounds diffusing up from the sediment. In deeper lakes, mineralization of OM transported through the water column to the sediment represented a greater proportion of  $O_2$  consumption. This process predominated in the most productive lakes but declined with decreasing total phosphorous (TP) concentrations and hence primary production, when TP concentrations fell below a threshold value of ~10 mg P  $m^{-3}$ . Oxygen uptake by the sediment and the flux of reduced compounds from the sediment in these deep lakes were 7.9–10.6 and 0.6–3.6  $mmol\ m^{-2}\ d^{-1}$ , respectively. These parameters did not depend on the lake's trophic state but did depend on sedimentation rates for the primarily allochthonous or already degraded OM. These results indicate that in lakes deeper than ~100 m, mineralization of autochthonous OM is mostly complete by the time of sedimentary burial. This explains why hypolimnetic  $O_2$  concentrations improve more rapidly following TP load reduction in deeper lakes relative to shallower lakes, where larger sediment-based  $O_2$  consumption by settled OM and release of reduced substances may inhibit the restoration of hypolimnetic  $O_2$  concentrations.

Sufficient  $O_2$  concentrations in the hypolimnia of lakes are essential for maintaining habitats, supporting fish reproductive cycles, and ensuring reliable drinking water sources. Oxygen concentrations are thus a key parameter for lake management. In the middle of the 20th century, increasing phosphorous (P) concentrations contributed to drastic increases in primary production and subsequent transport of OM to deep hypolimnetic waters. Aerobic mineralization of OM and the release of anaerobically produced compounds such as methane ( $CH_4$ ) and ammonium ( $NH_4^+$ ) into bottom waters are common contributors to hypolimnetic  $O_2$  depletion. Lacustrine OM mineralization generally occurs in three different domains. These include (1) the water column during settling of organic particles (Livingstone and Imboden 1996), (2) sediment surfaces subject to intensive remineralization and biological reworking (Burns 1995), and (3) underlying, deeper sediments that host gradual but continuous production of reduced substances (Carignan and Lean 1991; Steinsberger

et al. 2017).  $O_2$  consumption rates for each of these locations have been respectively termed (1) water column mineralization rate in the hypolimnion (WCM), (2) sediment oxygen uptake (SOU), and (3) flux of reduced compounds from the sediment,  $F_{red}$ , as converted to  $O_2$  consumption equivalents (Matzinger et al. 2010; Müller et al. 2012). The areal hypolimnetic mineralization (AHM) represents the sum of all three  $O_2$  consumption processes over the whole hypolimnion.

For fully productive (eutrophic) lakes of less than ~100 m depth, SOU (OM mineralization on the sediment surface) and  $F_{red}$  (reduced compounds diffusing from interstitial fluids in deeper sediments) represent the primary sites of  $O_2$  consumption (Matthews and Effler 2006; Müller et al. 2012). In deeper (>100 m) lakes such as the Great Laurentian Lakes, WCM and SOU govern  $O_2$  depletion (Charlton 1980). Studies of Lakes Michigan (Thomsen et al. 2004), Erie (Edwards et al. 2005; Smith and Matisoff 2008), and Superior (Li et al. 2012) have further demonstrated the importance of SOU and/or WCM as  $O_2$  sinks in these lakes. A study of the highly productive and deep Lake Geneva by Schwefel et al. (2018) showed that, as with the Great Lakes,  $O_2$  consumption primarily occurred in the water column rather than near or in the sediment.

\*Correspondence: beat.mueller@eawag.ch

Additional Supporting Information may be found in the online version of this article.

In general, deep lakes such as Baikal or Superior store enough O<sub>2</sub> in their large hypolimnia to prevent anoxia in bottom waters, assure aerobic OM remineralization and limit the production of reduced substances (Müller et al. 2005; Li et al. 2012). Even in the very productive and deep Lake Geneva, which exhibits a TP concentration of ~ 24 mg P m<sup>-3</sup> (average from 2008 to 2018) and a carbon assimilation rate of ~ 250 g C m<sup>-2</sup> yr<sup>-1</sup> (Anneville and Pelletier 2000; Soomets et al. 2019), sediment characteristics resemble those of an oligotrophic lake with low TOC content (~ 1%), very small fluxes of CH<sub>4</sub> and NH<sub>4</sub><sup>+</sup> from the sediment and limited O<sub>2</sub> consumption at the sediment–water interface (Steinsberger et al. 2017; Schwefel et al. 2018). The high fraction of WCM (~ 70%) on the total AHM diminishes the oxidizing capacity of the settling material on its way to the sediment. Given the mostly, already degraded OM, SOU only contributed about 27% to AHM. The O<sub>2</sub> made available by a large hypolimnion for extensive oxic mineralization of settling OM results in only a marginal amount of OM burial in sediments.

In shallower lakes (< 100 m depth) with similar productivity, shorter settling distances may limit (oxic) mineralization in the water column. This could lead to accumulation and burial of large volumes of mostly un-mineralized OM in sediments. The subsequent slow but persistent release of reduced compounds generated by anaerobic OM degradation further consumes O<sub>2</sub> and often impedes later restoration measures (Wehrli et al. 1997).

In shallow lakes, this “sediment memory effect” or “sediment legacy” and associated release of reduced substances from the sediment may account for up to 80% of the O<sub>2</sub> depletion in the hypolimnion even with measures aimed at reestablishing oligotrophic conditions (Matthews and Effler 2006; Carey et al. 2018). Although this effect has been documented in lakes of up to ~ 100 m depth, its potential impact on deeper lakes is not as well understood (Matzinger et al. 2010). In most deep lakes, the contribution of different

O<sub>2</sub> sinks to overall hypolimnion O<sub>2</sub> depletion remains debated or unknown.

This study aimed to determine the location and intensity of O<sub>2</sub> consumption in eight lakes of > 100 m depth and three lakes of < 100 m depth. We determined O<sub>2</sub> consumption rates at the sediment surface, fluxes of reduced compounds and O<sub>2</sub> depletion rates in the water column using lake monitoring data, O<sub>2</sub> microprofile measurements and simultaneous sediment pore-water analysis. Oxygen consumption was characterized and interpreted in terms of spatial location, trophic state, hypolimnion depth, and accumulation of OM.

## Material and methods

### Site description

Sediment cores of the lakes characterized in Table 1 were retrieved at the deepest locations of each lake except for Lakes Maggiore, Lucerne, and Constance. For Lake Maggiore, sediment cores were collected from 273 m depth at the deepest site of the Swiss part of the lake. For Lake Lucerne, sediment was collected in the deepest part of the Vitznau basin at 150 m water depth. For Lake Constance, sediment was collected at 150 m depth in the main basin. All sediment cores were collected during the lake’s stratified period (between May and October). Table S1 summarizes sampling details, location, and dates. The two shallowest lakes, Hallwil and Baldegg, both experience recurring hypolimnetic hypoxia caused by high primary production. Both lakes are artificially aerated during summer stratification and mixing has been supported by bubble plumes during overturn since the 1980s. Hypolimnetic O<sub>2</sub> sinks are thus influenced by artificially elevated O<sub>2</sub> concentrations introduced by lake restoration efforts.

### Sampling procedure

For each sampling site, four sediment cores were collected with a gravity corer. These cores were subjected to individual

**Table 1.** Physiographic and limnological characteristics of the lakes investigated by this study.  $A_H$  and  $V_H$  represent the hypolimnion area and volume, respectively.  $z_H$  is the mean hypolimnion depth ( $V_H/A_H$ ). The hypolimnion was defined as the water body below 15 m water depth.

Lake	Max. depth (m)	Volume (km <sup>3</sup> )	Surface (km <sup>2</sup> )	$A_H$ (km <sup>2</sup> )	$V_H$ (km <sup>3</sup> )	$z_H$ (m)
Maggiore (MAG)	372	37.6	213	196	34.5	176
Geneva (GEN)	310	89.1	582	534	80.8	151
Brienz (BRI)	260	5.0	29.8	27.8	4.59	165
Constance (CON)	253	47.8	482	403	41.4	103
Thun (THU)	217	6.46	47.9	43.9	5.78	132
Lucerne (LUC)	214	2.30	22	21.1	1.98	93.9
Neuchâtel (NEU)	154	13.8	218	156	11.1	71.3
Walensee (WAL)	151	2.51	24.1	22.8	2.15	94.3
Aegeri (AEG)	81	0.35	7.21	6.4	0.26	46.6
Baldegg (BAL)	66	0.17	5.22	4.53	0.13	27.6
Hallwil (HAL)	48	0.29	9.95	8.58	0.19	22.6

measurements of sediment pore water, CH<sub>4</sub>, O<sub>2</sub> micro-optode profiles, and bulk analysis of sediment chemical and physical properties.

Sediment cores subjected to bulk analysis were stored in a cold room (4°C) until processing. The cores were sampled as 0.5 to 1 cm sections, weighed and freeze-dried. Water content was calculated from the weight difference before and after drying. Sediment porosity was estimated from water content, sediment density, and TOC content (Och et al. 2012). After grinding with an agate mortar, the sediment was further analyzed for TOC/TN with a Euro EA 3000 (HEKAtech). Total organic carbon mass accumulation rates (TOC-MAR) were determined for each site at depths between 1 and 10 cm sediment depths in 5 mm spatial resolution. Estimation of TOC-MAR followed detailed methods outlined in Steinsberger et al. (2017). Sedimentation rates were derived from literature sources or estimated from <sup>210</sup>Pb and <sup>137</sup>Cs measurements as reported from prior analyses of the respective lake sediments.

For CH<sub>4</sub> and sediment pore-water analysis, 60-cm-long PVC liners with predrilled and taped holes (diameter 1.2 cm and 2 mm, respectively) were used. Syringes with clipped tips were used to sample 2 cm<sup>3</sup> of sediment aliquots for CH<sub>4</sub> analysis. Aliquots were directly transferred to 113.7 mL flasks containing 2 mL of 7 M NaOH, then capped, stored, and prepared for GC headspace analysis. The sampling resolution was 1 cm for the upper 10 cm, 2 cm for the middle 10 cm and 3 cm from 20 to 35 cm sediment depth.

Sediment pore water was extracted using MicroRhizon filter tubes of 1.5 mm diameter (Torres et al. 2013). Punching through the taped pre-drilled holes allowed for withdrawal of 10–50 μL of sediment pore water. This pore water was transferred to a flip tube and immediately injected onto a capillary electrophoresis (CE)-C<sup>4</sup>D column (Section 1.3). Relative to the sediment–water interface, pore water was sampled every 5 mm for the upper 5 cm, every 1 cm for the following 5 cm and every 2 cm from 10 to 20 cm sediment depth.

The fourth core was used to sample bottom water for O<sub>2</sub> concentration using the Winkler method. Vertical concentration profiles of O<sub>2</sub> in the sediment cores were performed with a PreSens O<sub>2</sub> micro-optode mounted onto an automated micromanipulator (Section 1.3).

### Oxygen and sediment pore-water analysis

Sediment pore-water analysis and O<sub>2</sub> concentration profiles were measured directly on site at each lake. Sediment pore-water samples were analyzed for anions and cations with two portable CE devices, each with a capacitively coupled contactless conductivity detector (C<sup>4</sup>D) calibrated directly on site (Kubáň et al. 2007). These devices were set at 15 kV and 0.5 μA to achieve complete separation of cations (NH<sub>4</sub><sup>+</sup>, K<sup>+</sup>, Ca<sup>2+</sup>, Na<sup>+</sup>, Mg<sup>2+</sup>, Fe<sup>2+</sup>, Mn<sup>2+</sup>) and anions (Cl<sup>-</sup>, NO<sub>3</sub><sup>-</sup>, NO<sub>2</sub><sup>-</sup>, SO<sub>4</sub><sup>2-</sup>, H<sub>2</sub>PO<sub>4</sub><sup>-</sup>) within 8 min (Torres et al. 2013). All calibration standards and the background electrolyte buffer were prepared from corresponding salts with ultrapure water (Merck)

immediately prior to each sampling campaign. These were checked using a multi-cation standard (Fluka) and a multi-anion standard (Roth). Standard deviations of all calibration measurements were < 5%.

The diffusive fluxes of reduced substances ( $J_{\text{CH}_4}$ ,  $J_{\text{NH}_4}$ ,  $J_{\text{Mn(II)}}$ ,  $J_{\text{Fe(II)}}$ ) from the sediments were calculated from pore-water concentration gradients (Supporting Information Fig. S1). The one-dimensional steady-state diffusion model by Epping and Helder (1997) was adapted to calculate fluxes:

$$J_i = -\frac{D}{G} \frac{dC_i}{dx} \quad (1)$$

For  $J$  in mmol m<sup>-2</sup> d<sup>-1</sup>,  $D$  represents the diffusion coefficient based on temperature corrected values from Li and Gregory (1974),  $G$  represents porosity corrected formation factor ( $1.02 * \text{porosity}^{-1.81}$ ) for clay-silt sediment, and  $\frac{dC_i}{dx}$  represents the concentration gradient of species  $i$  (Maerki et al. 2004). Based on the redox stoichiometry (Eq. 2), the estimated fluxes were converted to O<sub>2</sub> equivalents (mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>) (Matzinger et al. 2010; Müller et al. 2012) as follows:

$$F_{\text{red}} = 2 * J_{\text{CH}_4} + 2 * J_{\text{NH}_4} + 0.5 * J_{\text{Mn(II)}} + 0.25 J_{\text{Fe(II)}} \quad (2)$$

O<sub>2</sub> measurements were conducted with a PreSens O<sub>2</sub> micro-optode with a flat tip size of 230 μm, a detection limit of 0.64 μmol O<sub>2</sub> L<sup>-1</sup> and a response time < 10 s. The micro-optode was mounted to an automated micromanipulator. O<sub>2</sub> profiles were taken between 2 cm above and 3 cm below the sediment–water interface at 100 μm intervals and a penetration speed of 100 μm s<sup>-1</sup> stopping for each measurement, completing a full profile in < 9 min. The O<sub>2</sub> penetration depth was defined as the sediment depth where O<sub>2</sub> concentrations dropped below 3 μmol O<sub>2</sub> L<sup>-1</sup>. The O<sub>2</sub> micro-optode was calibrated in the laboratory with a two-point calibration. During the on-site measurements, standard deviation for O<sub>2</sub> measurements was typically ~ 11%. Up to nine replicate measurements were taken at different positions for each core.

Simultaneous O<sub>2</sub> measurements were not performed in Lakes Hallwil, Geneva, or Constance. Data from Bierlein et al. (2017) (Lake Hallwil), Schwefel et al. (2018) (Lake Geneva), and Sobek et al. (2009) (Lake Constance) were used for comparison instead.

The SOU flux was calculated similar to the flux of reduced substances (Eq. 1). Fluxes of the other two main electron acceptors, sulfate (SO<sub>4</sub><sup>2-</sup>) and nitrate (NO<sub>3</sub><sup>-</sup>), were converted to O<sub>2</sub> equivalents and added to SOU (Müller et al. 2007):

$$\text{SOU} = J_{\text{O}_2} + 1.25 * J_{\text{NO}_3} + 2 * J_{\text{SO}_4} \quad (3)$$

### Areal hypolimnetic and water column mineralization

Based on monthly monitoring data, average AHMs for the past 10 years (2007–2017) were calculated from differences in

hypolimnetic O<sub>2</sub> content (> 15 m water depth) as measured between maximum values after winter mixing (February to April) and minimum values from October to November.

If reduced substances such as NH<sub>4</sub><sup>+</sup> and/or CH<sub>4</sub> occurred in bottom waters, they were included in the budget with their O<sub>2</sub> equivalents (Matzinger et al. 2010). WCM was calculated by subtracting SOU and  $F_{\text{red}}$  from AHM as:

$$\text{AHM} = \text{SOU} + F_{\text{red}} + \text{WCM} \quad (4)$$

$$\begin{aligned} \text{WCM} = & \text{AHM} - \left( 2 * J_{\text{CH}_4} + 2 * J_{\text{NH}_4^+} + 0.5 * J_{\text{Mn(II)}} + 0.25 * J_{\text{Fe(II)}} \right) \\ & - (J_{\text{O}_2} + 1.25 * J_{\text{NO}_3^-} + 2 * J_{\text{SO}_4^{2-}}) \end{aligned} \quad (5)$$

### Comparability and limitations of SOU, $F_{\text{red}}$ , and WCM parameters

Hypolimnetic O<sub>2</sub> sinks fundamentally depend on aerobic or anaerobic OM mineralization. Although the scale and biogeochemical processes may differ, these sinks are comparable because all fluxes are normalized to the hypolimnion area and expressed as O<sub>2</sub> equivalents (mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>). However, spatial and temporal variation may influence the individual size of the O<sub>2</sub> sinks.

Sediment-based O<sub>2</sub> sinks,  $F_{\text{red}}$ , and SOU are controlled by the amount of settled OM and are therefore sensitive to spatial effects such as sediment focusing. Gravitational slumping or wind forcing can resuspend fine-grained, organic-rich material, and transport it from littoral areas to deeper parts of lakes (Lehman 1975; Bloesch and Uehlinger 1986; Carignan and Lean 1991). Seasonal variations may also influence SOU since hypolimnetic O<sub>2</sub> concentrations tended to decrease until the end of the stratified season. A single O<sub>2</sub> measurement may therefore over- or underestimate annual average SOU values. Seasonal variation in  $F_{\text{red}}$  is less relevant since it depends only indirectly on fluctuating OM input or overlying O<sub>2</sub>

concentrations (Frenzel et al. 1990; Schwefel et al. 2018; Steinsberger et al. 2019). By contrast, seasonal variation in primary production and associated settling and mineralization throughout the hypolimnetic water column strongly influence WCM but this variation is imbedded in yearly average AHM calculations.

## Results

### Fluxes of reduced substances from the sediment


Typical sediment concentration profiles of CH<sub>4</sub>, NH<sub>4</sub><sup>+</sup>, Fe(II), Mn(II), O<sub>2</sub>, NO<sub>3</sub><sup>-</sup>, and SO<sub>4</sub><sup>2-</sup> are shown in the Supporting Fig. S1. Table 2 summarizes fluxes ( $J$ ) of all measured species converted to O<sub>2</sub> equivalents from the 11 lakes.

In lakes deeper than 100 m,  $F_{\text{red}}$  showed similar low values of between 0.6 and 3.6 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>. Shallower Lakes Hallwil, Baldegg, and Aegeri exhibited higher  $F_{\text{red}}$  values of between 4.0 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> (Lake Aegeri) and 13.8 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> (Lake Baldegg).

CH<sub>4</sub> and NH<sub>4</sub><sup>+</sup> represented the largest contributions to  $F_{\text{red}}$ .  $J_{\text{Fe(II)}}$  and  $J_{\text{Mn(II)}}$  were comparatively negligible and contributed on average only 1% and 2%, respectively, even though deeper parts of the sediment (> 10 cm depth) exhibited high sediment pore-water concentrations. Lago Maggiore sediment pore water, for example, contained 0.7 mmol Mn(II) L<sup>-1</sup> and Lake Walensee sediment pore water contained 0.8 mmol Fe(II) L<sup>-1</sup>. In lakes of > 100 m depth, NH<sub>4</sub><sup>+</sup> and CH<sub>4</sub> pore-water concentrations were at or below detection limits in the uppermost sediment layers (sometimes down to 5 cm). These values increased to 0.3–0.6 mmol L<sup>-1</sup> (NH<sub>4</sub><sup>+</sup>) and 2–7 mmol L<sup>-1</sup> (CH<sub>4</sub>) downcore (see Supporting Information Fig. S1). Except for Lago Maggiore, NH<sub>4</sub><sup>+</sup> did not reach the sediment surface but was oxidized within the uppermost centimeter. Here,  $J_{\text{CH}_4}$  represented up to 99% of  $F_{\text{red}}$  (Fig. 1). In the shallower Lakes Hallwil, Baldegg, and Aegeri (< 100 m depth),

**Table 2.** Sediment pore-water fluxes of reduced and oxidized species expressed as O<sub>2</sub> equivalents in mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>. Reduced species are summed as  $F_{\text{red}}$  and oxidants as SOU. Values of Lake Constance are averages from three sediment cores. Lakes are sorted according to decreasing mean hypolimnion depth.

Lake	$J_{\text{CH}_4}$	$J_{\text{NH}_4^+}$	$J_{\text{Fe(II)}}$	$J_{\text{Mn(II)}}$	$F_{\text{red}}$	$J_{\text{O}_2}$	$J_{\text{NO}_3^-}$	$J_{\text{SO}_4^{2-}}$	SOU
Maggiore	3.85	1.96	0.04	0.13	5.97	14.0	0.47	1.95	16.4
Geneva	1.74	0.30	0.01	0.03	2.07	4.48	0.86	2.7	8.04
Brienzen	1.40	0	0	0.02	1.42	7.06	0.48	0.33	7.87
Constance	1.31	0.54	0.04	0.11	2.02	6.78	0.53	1.28	8.59
Thun	2.25	0	0	0.05	2.30	8.24	0.32	2.05	10.6
Lucerne	0.57	0	0.03	0.03	0.62	9.08	1.37	0.50	11.0
Neuchâtel	3.06	0	0.01	0.01	3.08	9.52	0.60	0.50	10.6
Walensee	3.32	0	0.07	0.18	3.57	16.2	0.76	0.66	17.6
Aegeri	2.15	1.59	0.11	0.11	3.96	10.6	0.46	0.48	11.6
Baldegg	7.71	5.68	0.25	0.11	13.8	18.8	2.36	2.04	23.2
Hallwil	3.48	1.96	0.13	0.03	5.60	12.4	1.10	1.32	14.8



**Fig. 1.** Flux at the sediment-water-interface (SWI) expressed in O<sub>2</sub> equivalents (mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>) of the main reduced substances, NH<sub>4</sub><sup>+</sup> and CH<sub>4</sub> (shown negatively scaled as black and red bars), and oxidants, O<sub>2</sub>, SO<sub>4</sub><sup>2-</sup>, and NO<sub>3</sub><sup>-</sup> (positively scaled as light blue, green, and purple bars, respectively). Lakes were sorted according to  $F_{\text{red}}$ . Fluxes of Mn(II) and Fe(II) were negligible and are not shown.

in addition to  $J_{\text{CH}_4}$  (54% to 62%),  $J_{\text{NH}_4}$  was responsible for the remaining fraction of  $F_{\text{red}}$  (Fig. 1; ~40%).

### Sediment oxygen uptake and penetration depths

Among lakes deeper than 100 m, SOU rates ranged only from 7.9 to 11.0 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>. However, SOU values in Lakes Maggiore (16.4 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>) and Walensee (17.6 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>) were exceptionally high and resembled the SOU average of 19.0 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> measured from the shallower and more productive Lakes Hallwil and Baldegg.

All lakes showed high O<sub>2</sub> concentrations in the hypolimnion (> 0.19 mmol O<sub>2</sub> L<sup>-1</sup>, Supporting Information Fig. S2). However, penetration depths into the sediment were low (between 2.0 ± 1.0 mm in Lake Baldegg and 6.3 ± 0.8 mm in Lake Brienz, see Table 4) and did not correlate with bottom water O<sub>2</sub> concentrations.  $J_{\text{O}_2}$  ranged from 4.4 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> in Lake Geneva to 18.8 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> in Lake Baldegg (Table 2).  $J_{\text{O}_2}$  contributed an average of 78 ± 13% to total SOU.  $J_{\text{NO}_3}$  and  $J_{\text{SO}_4}$  varied between 0.3–2.4 and 0.3–2.0 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> and respectively represented 6.7% ± 3.2% and 11.2% ± 9.1% of SOU. For lakes with high hypolimnetic NO<sub>3</sub><sup>-</sup> and/or SO<sub>4</sub><sup>2-</sup> concentrations (> 0.4 mmol L<sup>-1</sup>; Lakes Geneva and Thun), combined  $J_{\text{NO}_3}$  and  $J_{\text{SO}_4}$  represented up to 45% of SOU (Fig. 1).

### Mineralization in the water column

WCM rates varied between 2.1 and 27.4 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>. As a measure of OM mineralization within the hypolimnetic water column, WCM depends on the lakes' volume and

**Table 3.** Estimation of the water column mineralization rate (WCM) calculated as the difference between AHM and the sum of SOU and  $F_{\text{red}}$ . TP is the average concentration after winter mixing over the past 10 years. For Lake Brienz, the value of 0.6 mg P m<sup>-3</sup> is calculated as the bio-available TP (Müller et al. 2006).

Lake	$F_{\text{red}}$	SOU	AHM	WCM	TP
	(mmol O <sub>2</sub> m <sup>-2</sup> d <sup>-1</sup> )				(mg P m <sup>-3</sup> )
Maggiore	6.0	16.4	26.3	3.9	11.0
Geneva	2.1	8.0	37.5	27.4	24.3
Brienz	1.4	7.9	13.1	3.8	0.6
Constance	2.0	8.6	25.0	14.4	7.5
Thun	2.3	10.6	15.0	2.1	1.8
Lucerne	0.6	11.0	20.3	8.7	4.8
Neuchâtel	3.1	10.6	24.7	11.0	9.8
Walensee	3.6	17.6	20.0	~0	3.2
Aegeri	4.0	11.6	19.7	4.1	5.4
Baldegg	13.8	23.2	42.2	5.2	24.1
Hallwil	5.6	14.8	30.9	10.5	15.8

primary production and thus its TP concentration. Oligotrophic lakes (TP < 10 mg P m<sup>-3</sup>) that have not experienced eutrophication in the past (Walensee, Lakes Thun, Brienz, and Aegeri) exhibited low WCM values of 2.1 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> (Lake Thun) and 4.2 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> (Lake Aegeri). WCM could not be estimated from Walensee because the sum of  $F_{\text{red}}$  (3.6 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>) and SOU (17.6 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>) already equaled measured AHM (20.0 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>; Table 3). Lago Maggiore similarly showed a very low WCM value (3.9 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>). Both lakes have exceptionally steep walls which cause substantial sediment focusing leading to elevated SOU and  $F_{\text{red}}$  at the central coring sites. In the oligotrophic but recovering Lake Constance, WCM values resembled those measured from the mesotrophic Lake Neuchâtel (14.4 and 11.0 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>, respectively). The highest WCM rate was observed from the most productive very deep Lake Geneva (27.4 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>). The shallower but similarly productive Lakes Baldegg and Hallwil exhibited lower respective WCM rates of 5.2 and 10.5 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> (Table 3).

## Discussion

### Fluxes of reduced substances from the sediment

Very low  $F_{\text{red}}$  values (< 0.2 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>) were previously reported in lakes > 100 m deep such as Lakes Superior (Remsen et al. 1989; Li et al. 2012), Michigan (Thomsen et al. 2004), and Baikal (Och et al. 2012; Torres et al. 2014). These lakes are characterized by low OM accumulation rates and high O<sub>2</sub> penetration depths resulting in prolonged (up to several decades) aerobic mineralization of the settled

OM. Consequently, fluxes of CH<sub>4</sub>, NH<sub>4</sub><sup>+</sup>, Fe(II) Mn(II) and hence,  $F_{\text{red}}$ , are close to zero.

The deep lakes investigated by this study (> 100 m) exhibited distinctly higher  $F_{\text{red}}$  values (average  $2.6 \pm 1.6$  mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>) and showed a consistent dependence on mean lake depth (Fig. 2) (Steinsberger et al. 2017). These were dominated by CH<sub>4</sub> fluxes which constituted up to 99% of  $F_{\text{red}}$  (Table 2, Fig. 1). NH<sub>4</sub><sup>+</sup> contributions were negligible in all lakes that have never surpassed mesotrophic production in the past. Deeper sedimentary horizons (> 10 cm) however exhibited elevated CH<sub>4</sub> and NH<sub>4</sub><sup>+</sup> concentration of up to 7.0 and 0.6 mmol L<sup>-1</sup>, respectively. These resembled values measured from river deltas or from shallower, more productive lakes (Sobek et al. 2009; Randlett et al. 2015; Steinsberger et al. 2017). In contrast to the Great Lakes, the deep peri-alpine lakes investigated by this study accumulated inorganic matter from weathering of rocks and OM from their watersheds. This gradually degrading allochthonous OM is diluted with additional mineral deposits resulting in small concentrations of TOC (Supporting Information Fig. S3). High sedimentation rates lead to rapid OM burial in the anoxic sediments. A laboratory study by Grasset et al. (2018) showed that the mineralization of allochthonous OM by methanogenesis, although slower than autochthonous OM degradation, can yield similar overall CH<sub>4</sub> production. This finding helps to explain high pore-water accumulation of CH<sub>4</sub> and NH<sub>4</sub><sup>+</sup> in deeper sediments collected from Walensee and Lago Maggiore.

Consequently, upward diffusing CH<sub>4</sub> and NH<sub>4</sub><sup>+</sup> comes into contact with the main electron acceptors O<sub>2</sub>, NO<sub>3</sub><sup>-</sup>, and SO<sub>4</sub><sup>2-</sup>, and is (partially) oxidized within the sediment. In consequence, a substantial fraction of O<sub>2</sub>, NO<sub>3</sub><sup>-</sup>, or SO<sub>4</sub><sup>2-</sup>

consumption could be attributed to  $F_{\text{red}}$  oxidation rather than by direct aerobic mineralization of OM.

In productive lakes < 100 m deep (Lakes Baldegg and Hallwil), pore-water concentration gradients of CH<sub>4</sub>, NH<sub>4</sub><sup>+</sup>, Fe(II), and Mn(II) extended to the sediment–water interface.  $F_{\text{red}}$  attained high rates of 5.6 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> (Lake Hallwil) and 13.8 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> (Lake Baldegg) fueled by the high primary production and mineralization of large amounts of TOC deposited during past anoxic periods (Steinsberger et al. 2019).


### Sediment oxygen uptake dependence on organic matter

O<sub>2</sub> is depleted by the initial aerobic degradation of OM at the sediment surface and by oxidation of reduced compounds produced during anaerobic mineralization within the sediment (Steinsberger et al. 2017). Lakes > 100 m deep exhibited SOU in the narrow 7.9–11.0 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> range. Walensee and Lago Maggiore exhibited exceptionally high SOU values of 17.6 and 16.4 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> respectively, which resemble values observed in the shallower and more productive Lakes Hallwil and Baldegg (14.8 and 23.2 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>; Table 3). The steep slopes of Walensee and Lago Maggiore may increase OM deposition rates by focusing sediment into the central parts of the basin which in turn increases sediment-related O<sub>2</sub> consumption. The sediment area relevant for SOU and  $F_{\text{red}}$  is thus substantially smaller than the hypolimnion area used for AHM estimation. Equation 4 may thus systematically underestimate WCM values for lakes having steep walls.

Sediment-related O<sub>2</sub> consumption depends on several key parameters such as bottom water O<sub>2</sub> concentrations, O<sub>2</sub> penetration depth, TOC mass accumulation rate, and OM susceptibility to mineralization (Polerecky et al. 2006; Sobek et al. 2009; Bryant et al. 2010; Gudasz et al. 2010). All lakes > 100 m deep exhibited sediment TOC contents of 0.5–1.5%, whereas shallower lakes exhibited sediment TOC concentrations of 3.4–3.5% (Supporting Information Fig. S3, Table 4). The more rapid mineralization rates for autochthonous OM relative to the more recalcitrant terrestrial OM (Sobek et al. 2009) suggest that the former is preferentially mineralized in the water column and then deposited mostly as already degraded OM onto the sediment for lakes deeper than 100 m. Schwefel et al. (2018) demonstrated this for the productive Lake Geneva, while Carstens et al. (2013) did so for the oligotrophic Lake Brienz. Consequently, we observed a decreasing trend of SOU with increasing mean lake depths, similar to  $F_{\text{red}}$  (see Fig. 2).

Elevated TOC concentrations in sediment of lakes < 100 m deep is consistent with the interpretation of limited degradation within the water column and more vigorous O<sub>2</sub> consumption in the sediment surface leading to burial of large amounts of partially mineralized OM in sediments.

Table 4 shows that sediment mass accumulation rates in > 100 m deep lakes (Maggiore, Thun, Brienz, and Walensee) unexpectedly resembled or even exceeded that observed from shallower lakes. This reflects elevated areal contributions of allochthonous material from watersheds



**Fig. 2.** Depletion rates (mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>) of the sediment-based O<sub>2</sub> sinks SOU (blue) and  $F_{\text{red}}$  (red). SOU and  $F_{\text{red}}$  decrease with increasing mean hypolimnion depth (see Table 1).



**Table 4.** Nutrient contents of the bulk sediment. TOC, TN, and TOC-MAR values were averaged over 1–10 cm sediment depth. n.d., not determined. Some data were taken from the respective literature: <sup>1</sup>Frenzel et al. (1990), <sup>2</sup>Span et al. (1990), <sup>3</sup>Schwalb et al. (1998), <sup>4</sup>Müller et al. (2007), <sup>5</sup>Zimmermann (2008), <sup>6</sup>Putyrskaya et al. (2009), <sup>7</sup>Sobek et al. (2009), <sup>8</sup>Wirth et al. (2011), <sup>9</sup>Bierlein et al. (2017), <sup>10</sup>Steinsberger et al. (2017), <sup>11</sup>Schwefel et al. (2018).

Lake	TOC (mg C g <sup>-1</sup> )	TN (mg N g <sup>-1</sup> )	O <sub>2</sub> penetr. depth (mm)	Sed. accumul. rate (mm yr <sup>-1</sup> )	TOC-MAR (g C m <sup>-2</sup> yr <sup>-1</sup> )
Maggiore	14.9 ± 6.2	1.2 ± 0.6	2.6 ± 0.1	3.4 <sup>6</sup>	33.4
Geneva	11.1 <sup>1</sup>	1.1 <sup>2</sup>	4.0 ± 0.5 <sup>11</sup>	1.0 <sup>10</sup>	11.1
Brienzen	5.4 ± 1.4	0.6 ± 0.1	6.3 ± 0.8	6.4 <sup>4</sup>	36.8
Constance	n.d.	n.d.	2.5–4.5 <sup>1,7</sup>	n.d.	n.d.
Thun	11.3 ± 2.0	1.2 ± 0.2	4.8 ± 0.2	6.8 <sup>8</sup>	67.7
Lucerne	n.d.	n.d.	4.8 ± 0.6	n.d.	n.d.
Neuchâtel	22.4 ± 0.8	2.5 ± 0.4	3.6 ± 0.4	2.5 <sup>3</sup>	23.9
Walensee	11.3 ± 1.8	1.3 ± 0.1	5.3	5.7 <sup>5</sup>	42.9
Aegeri	34.9	3.6 ± 0.7	2.5 ± 0.7	1.9 <sup>10</sup>	22.8
Baldegg	34.2 ± 3.1	3.9 ± 0.4	3.4 ± 0.8	3.3 <sup>10</sup>	45.6
Hallwil	34.4 ± 3.5	4.0 ± 0.2	3.0 ± 1.0 <sup>9</sup>	2.0 <sup>10</sup>	22.5

(Wirth et al. 2011; Thevenon et al. 2013) as well as sediment focusing by funnel- or trench-shaped lake basins (Lehman 1975; Blais and Kalff 1995). The exposure time of OM to O<sub>2</sub> falls well below 1 year. Combined with elevated MAR, these conditions bury significant amounts of OM in anoxic sediment before degradation can occur. The elevated TOC-MAR lead to anaerobic mineralization in the deeper sediment which then generates CH<sub>4</sub> and NH<sub>4</sub><sup>+</sup> observed at high concentrations in the sediment pore water. These species can delay lake recovery from eutrophication with respect to O<sub>2</sub> concentrations. However, all of the Fe(II) and Mn(II) and most of the NH<sub>4</sub><sup>+</sup> are oxidized within the sediment. These do not reach the sediment–water interface and thereby increase SOU (Supplementary Fig. S1).

The range of SOU determined from lakes > 100 m deep investigated here (11.3 ± 3.7 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>) is consistent with fluxes reported by literature sources for other large and deep lakes. Thomsen et al. (2004) reported a SOU rate of 7.7 ± 2.1 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> for Lake Michigan while Li et al. (2012) reported a SOU rate of 6.1 ± 1.4 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> for Lake Superior. Lower SOU values were reported for the ultra-deep oligotrophic Lake Baikal. These include an overall value of 3.3 ± 1.1 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> (Müller et al. 2005), 1.4 to 5.6 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> for the South Basin (Och et al. 2012), and 1.1 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> for the North Basin (Martin et al. 1998). O<sub>2</sub> penetration depths determined for the deep lakes analyzed here (2.6–6.3 mm, Table 4) resembled values previously reported for Lake Constance (1.5–6.3 mm; Frenzel et al. 1990; Sobek et al. 2009) but were distinctly lower than those reported for Lakes Michigan (21 ± 2.5 mm), Superior (40–110 mm), and Baikal (20–44 mm). This evidences the high consumption of O<sub>2</sub> within the first millimeters of sediment rendering O<sub>2</sub> penetration depths more similar to those in the fully productive and shallower Lake Baldegg

(0.9–2.0 mm). Lakes Michigan and Superior exhibit low SOU rates and corresponding higher O<sub>2</sub> penetration depths that result from low OM sedimentation rates, intense bioturbation and extensive Fe cycling with only small NH<sub>4</sub><sup>+</sup> and insignificant CH<sub>4</sub> production. For lakes analyzed in this study, excess O<sub>2</sub> that could reach deeper parts of the sediment due to the absence of reactive OM is apparently consumed by a front of upward diffusing reduced compounds produced by anaerobic mineralization of OM. Decreasing pore-water concentrations of reduced species in the uppermost layers of sediment reported here offer further evidence of rapid anaerobic turnover (Maerki et al. 2006).

Fluxes of NO<sub>3</sub><sup>-</sup> into the sediment indicating mineralization of OM by denitrification ranged from 0.32 to 2.36 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> and thereby only a small fraction of total SOU (between 2.9% in Lago Maggiore and 12.5% in Lake Lucerne). Li et al. (2012) and Thomsen et al. (2004) report similarly low estimates for Lakes Superior and Michigan (< 5%). Overall NO<sub>3</sub><sup>-</sup> removal rates (coupled nitrification–denitrification) however may be higher due to a hidden contribution from anaerobic NH<sub>4</sub><sup>+</sup> oxidation with low concentrations but high turnover rates of NO<sub>3</sub><sup>-</sup>/NO<sub>2</sub><sup>-</sup> to N<sub>2</sub> (e.g., anammox; Small et al. 2013; Crowe et al. 2017). In fact, at least in five of the lakes investigated here (Walensee, Neuchâtel, Thun, Brienzen, and Lucerne), a rapid decline in NH<sub>4</sub><sup>+</sup> concentrations (with NH<sub>4</sub><sup>+</sup> consumption rates of 0.08–0.46 mmol m<sup>-2</sup> d<sup>-1</sup>) well below the O<sub>2</sub> penetration depth was evident. These conditions clearly indicate anaerobic NH<sub>4</sub><sup>+</sup> oxidation (Zhu et al. 2013). As NO<sub>3</sub><sup>-</sup> penetrates deeper into the sediment than O<sub>2</sub> (5 mm – 45 mm), anammox bacteria might be actively oxidizing NH<sub>4</sub><sup>+</sup> (Wenk et al. 2013).


Fluxes of SO<sub>4</sub><sup>2-</sup> ranged from 0.32 to 2.70 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> (average 1.19 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>), and were well within the range of SO<sub>4</sub><sup>2-</sup> reduction rates given by Holmer and

Storkholm (2001). In general, SO<sub>4</sub><sup>2-</sup> fluxes contributed ~ 11% to SOU, while in Lake Geneva and Lake Thun, due to high loads from the catchment geology, SO<sub>4</sub><sup>2-</sup> fluxes contributed about one third to SOU. The full extent of SO<sub>4</sub><sup>2-</sup> reduction induced by OM mineralization is likely masked by (cryptic) internal sulfide re-oxidation and sulfite complexation processes (Holmkvist et al. 2011). Hence, SO<sub>4</sub><sup>2-</sup> fluxes likely underestimate actual SO<sub>4</sub><sup>2-</sup> reduction rates.

### Mineralization in the water column

In the hypolimnetic water column, OM is mineralized by a large variety of microbes that consume O<sub>2</sub> (Carstens et al. 2012). Subsequent O<sub>2</sub> consuming processes at the sediment–water interface depend on excess OM passing through the water column. Therefore, depending on the main mineralization sites of OM (hypolimnetic water, sediment surface, deep sediment) lakes show characteristic patterns in the three O<sub>2</sub> sinks (WCM, SOU, and  $F_{red}$ ).


Livingstone and Imboden (1996) proposed that WCM represents a significant proportion of AHM for large lakes with a high mean hypolimnion depth. This is true for productive lakes such as Geneva (77%) and Neuchâtel (45%) and lakes recovering from a eutrophic phase, such as Lakes Constance (58%) and Lucerne (43%) (Table 3, Fig. 3). This is in agreement with the modeled WCM value of Lake Geneva by Schwefel et al. (2018). However, lakes that never experienced high productivity periods, such as the oligotrophic Lakes Brienz, Thun, and Walensee, did not show elevated WCM rates. These lakes instead showed WCM contributions to AHM of



**Fig. 3.** Hypolimnetic O<sub>2</sub> consumption itemized to the three O<sub>2</sub> sinks:  $F_{red}$ , SOU, and WCM. Lakes are divided into two groups: very deep lakes (above dashed line), lakes biased by intense sediment focusing which causes increased SOU and  $F_{red}$  (Lago Maggiore and Walensee) and artificially aerated deep Lakes Baldegg and Hallwil (below dashed line).

only ~ 14%, ~ 29% and ~ 0%, respectively. They also exhibited low sediment TOC content but a high rate of sediment OM accumulation caused by high loads of allochthonous particles (Finger et al. 2007; Bechtel and Schubert 2009). These lakes also mineralize a relatively small amount of OM within the hypolimnetic water column compared to that deposited at the sediment surface (Fig. 3). The high SOU and  $F_{red}$  rates indicate mineralization of allochthonous OM. The very low WCM of Walensee and Lago Maggiore reflect high SOU and  $F_{red}$  caused by sediment focusing (Blais and Kalff 1995). Lake Baldegg exhibited WCM of only 5.2 mmol O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> or 12% of AHM. This likely reflects the shallower lake depth and brief mineralization window for settling autochthonous OM.

WCM primarily depends on the export of autochthonous OM from the productive zone to the hypolimnion. Because it originates from primary production, it must be therefore related to TP for concentration ranges where TP availability influences primary production. In order to compare WCM between lakes of different depths, we normalized WCM to the mean hypolimnion depth,  $z_H$  to obtain a relationship with TP (Fig. 4). In all non-aerated lakes, WCM increases in proportion to TP. For a TP range of 10–15 mg P m<sup>-3</sup>, WCM levels off indicating that additional TP exerts limited influence on primary production and thus on OM export to the hypolimnion and subsequent O<sub>2</sub> consumption. Below a TP concentration of ~ 10 mg P m<sup>-3</sup>, WCM covaries closely with TP concentration. This is in line with the observation of Müller et al. (2019) that AHM in seasonally stratified lakes increases approximately in proportion to the areal amount of bio-available P but levels off above this threshold value. O<sub>2</sub> consumption in lakes > 100 m deep may immediately respond to a decrease of TP for concentrations below this threshold. Decreasing TP concentration in Lake Geneva by half could initiate a substantial reduction in WCM, the lake's primary O<sub>2</sub> sink. By contrast, minor interventions to reduce TP levels for Lakes



**Fig. 4.** Oxygen consumption rates for the water column (WCM) normalized by mean hypolimnion depth ( $z_H$ ). WCM increases with TP up to a value of around ~ 10 mg P m<sup>-3</sup> and then levels off. Lakes Baldegg and Hallwil are not shown because they are artificially aerated.



Neuchâtel and Constance would readily influence WCM and thereby decrease total AHM.

## Conclusion

Diffusive fluxes of reduced substances from and flux of O<sub>2</sub> into the sediments were measured in lakes with depths ranging from 48 to 372 m. Areal hypolimnetic mineralization rates were calculated from (monthly) O<sub>2</sub> monitoring data and used to estimate the water column mineralization rates. This analysis showed that:

- Despite high O<sub>2</sub> concentrations (> 190 μmol O<sub>2</sub> L<sup>-1</sup>) in bottom waters of all lakes, O<sub>2</sub> penetrated sediments only to depths of 2.6 to 6.3 mm resulting in high O<sub>2</sub> fluxes. In lakes deeper than 100 m, reduced substances from anaerobic mineralization of OM in the sediment constituted a major fraction of the sediment oxygen uptake and prevent O<sub>2</sub> from penetrating deeper into the sediment.
- Most lakes deeper than 100 m exhibited similar levels of sediment O<sub>2</sub> consumption by SOU and  $F_{\text{red}}$ . This suggests that OM deposited at the sediment surface was already largely mineralized. The O<sub>2</sub> reservoir of the more extensive hypolimnia sufficed to degrade most of the settling OM. Hence, SOU and  $F_{\text{red}}$  values did not depend on trophic state for lakes deeper than 100 m.
- WCM represented a major contribution to AHM in lakes of > 100 m depth. This contribution decreased with TP when less autochthonous OM was available for O<sub>2</sub> consumption. WCM increased with increasing TP concentrations until a threshold of 10–15 mg P m<sup>-3</sup> was reached. Above this threshold, WCM leveled off. Reducing TP concentrations in productive lakes deeper than 100 m to below this threshold could therefore exert a strong influence on AHM.

## References

- Anneville, O., and J. P. Pelletier. 2000. Recovery of Lake Geneva from eutrophication: Quantitative response of phytoplankton. *Arch. Hydrobiol.* **148**: 607–624. <https://doi.org/10.1127/archiv-hydrobiol/148/2000/607>
- Bechtel, A., and C. J. Schubert. 2009. A biogeochemical study of sediments from the eutrophic Lake Lugano and the oligotrophic Lake Brienz, Switzerland. *Org. Geochem.* **40**: 1100–1114. <http://doi.org/10.1016/j.orggeochem.2009.06.009>
- Bierlein, K. A., M. Rezvani, S. A. Socolofsky, L. D. Bryant, A. Wüest, and J. C. Little. 2017. Increased sediment oxygen flux in lakes and reservoirs: The impact of hypolimnetic oxygenation. *Water Resour. Res.* **53**: 4876–4890. <http://doi.org/10.1002/2016WR019850>
- Blais, J. M., and J. Kalff. 1995. The influence of lake morphometry on sediment focusing. *Limnol. Oceanogr.* **40**: 582–588. <http://doi.org/10.4319/lo.1995.40.3.0582>
- Bloesch, J., and U. Uehlinger. 1986. Horizontal sedimentation differences in a eutrophic Swiss lake. *Limnol. Oceanogr.* **31**: 1094–1109. <https://doi.org/10.4319/lo.1986.31.5.1094>
- Bryant, L. D., D. F. McGinnis, C. Lorrai, A. Brand, J. C. Little, and A. Wüest. 2010. Evaluating oxygen fluxes using micro-profiles from both sides of the sediment-water interface. *Limnol. Oceanogr.: Methods* **8**: 610–627. <https://doi.org/10.4319/lom.2010.8.0610>
- Burns, N. M. 1995. Using hypolimnetic dissolved oxygen depletion rates for monitoring lakes. *N. Z. J. Mar. Freshwat. Res.* **29**: 1–11. <https://doi.org/10.1080/00288330.1995.9516634>
- Carey, C. C., J. P. Doubek, R. P. McClure, and P. C. Hanson. 2018. Oxygen dynamics control the burial of organic carbon in a eutrophic reservoir. *Limnol. Oceanogr. Lett.* **3**: 293–301. <https://doi.org/10.1002/lo2.10057>
- Carignan, R., and D. Lean. 1991. Regeneration of dissolved substances in a seasonally anoxic lake: The relative importance of processes occurring in the water column and in the sediments. *Limnol. Oceanogr.* **36**: 683–707. <https://doi.org/10.4319/lo.1991.36.4.0683>
- Carstens, D., K. E. Köllner, H. Bürgmann, B. Wehrli, and C. J. Schubert. 2012. Contribution of bacterial cells to lacustrine organic matter based on amino sugars and d-amino acids. *Geochim. Cosmochim. Acta* **89**: 159–172. <https://doi.org/10.1016/j.gca.2012.04.052>
- Carstens, D., M. F. Lehmann, T. B. Hofstetter, and C. J. Schubert. 2013. Amino acid nitrogen isotopic composition patterns in lacustrine sedimenting matter. *Geochim. Cosmochim. Acta* **121**: 328–338. <https://doi.org/10.1016/j.gca.2013.07.020>
- Charlton, M. N. 1980. Hypolimnion oxygen consumption in lakes: Discussion of productivity and morphometry effects. *Can. J. Fish. Aquat. Sci.* **37**: 1531–1539. <http://doi.org/10.1139/f80-198>
- Crowe, S. A., and others. 2017. Novel anammox bacteria and nitrogen loss from Lake Superior. *Sci. Rep.* **7**: 13757. <https://doi.org/10.1038/s41598-017-12270-1>
- Edwards, W. J., J. D. Conroy, and D. A. Culver. 2005. Hypolimnetic oxygen depletion dynamics in the central basin of Lake Erie. *J. Great Lakes Res.* **31**: 262–271. [https://doi.org/10.1016/S0380-1330\(05\)70319-1](https://doi.org/10.1016/S0380-1330(05)70319-1)
- Epping, E. H., and W. Helder. 1997. Oxygen budgets calculated from in situ oxygen microprofiles for Northern Adriatic sediments. *Cont. Shelf Res.* **17**: 1737–1764. [https://doi.org/10.1016/S0278-4343\(97\)00039-3](https://doi.org/10.1016/S0278-4343(97)00039-3)
- Finger, D., and others. 2007. Effects of alpine hydropower operations on primary production in a downstream lake. *Aquat. Sci.* **69**: 240–256. <http://doi.org/10.1007/s00027-007-0873-6>
- Frenzel, P., B. Thebrath, and R. Conrad. 1990. Oxidation of methane in the oxic surface layer of a deep lake sediment (Lake Constance). *FEMS Microbiol. Ecol.* **6**: 149–158. <http://doi.org/10.1111/j.1574-6968.1990.tb03935.x>

- Grasset, C., R. Mendonça, G. Villamor Saucedo, D. Bastviken, F. Roland, and S. Sobek. 2018. Large but variable methane production in anoxic freshwater sediment upon addition of allochthonous and autochthonous organic matter. *Limnol. Oceanogr.* **63**: 1488–1501. <http://doi.org/10.1002/lno.10786>
- Gudasz, C., D. Bastviken, K. Steger, K. Premke, S. Sobek, and L. J. Tranvik. 2010. Temperature-controlled organic carbon mineralization in lake sediments. *Nature* **466**: 478–481. <https://doi.org/10.1038/nature09186>
- Holmer, M., and P. Storkholm. 2001. Sulphate reduction and sulphur cycling in lake sediments: A review. *Freshwat. Biol.* **46**: 431–451. <http://doi.org/10.1046/j.1365-2427.2001.00687.x>
- Holmkvist, L., T. G. Ferdelman, and B. B. Jørgensen. 2011. A cryptic sulfur cycle driven by iron in the methane zone of marine sediment (Aarhus Bay, Denmark). *Geochim. Cosmochim. Acta* **75**: 3581–3599. <https://doi.org/10.1016/j.gca.2011.03.033>
- Kubáň, P., H. T. A. Nguyen, M. Macka, P. R. Haddad, and P. C. Hauser. 2007. New fully portable instrument for the versatile determination of cations and anions by capillary electrophoresis with contactless conductivity detection. *Electroanalysis* **19**: 2059–2065. <https://doi.org/10.1002/elan.200703908>
- Lehman, J. T. 1975. Reconstructing the rate of accumulation of lake sediment: The effect of sediment focusing. *Quatern. Res.* **5**: 541–550. [https://doi.org/10.1016/0033-5894\(75\)90015-0](https://doi.org/10.1016/0033-5894(75)90015-0)
- Li, J., S. A. Crowe, D. Miklesh, M. Kistner, D. E. Canfield, and S. Katsev. 2012. Carbon mineralization and oxygen dynamics in sediments with deep oxygen penetration, Lake Superior. *Limnol. Oceanogr.* **57**: 1634–1650. <https://doi.org/10.4319/lo.2012.57.6.1634>
- Li, Y.-H., and S. Gregory. 1974. Diffusion of ions in sea water and in deep-sea sediments. *Geochim. Cosmochim. Acta* **38**: 703–714. [http://doi.org/10.1016/0016-7037\(74\)90145-8](http://doi.org/10.1016/0016-7037(74)90145-8)
- Livingstone, D. M., and D. M. Imboden. 1996. The prediction of hypolimnetic oxygen profiles: A plea for a deductive approach. *Can. J. Fish. Aquat. Sci.* **53**: 924–932. <https://doi.org/10.1139/f95-230>
- Maerki, M., B. Müller, and B. Wehrli. 2006. Microscale mineralization pathways in surface sediments: A chemical sensor study in Lake Baikal. *Limnol. Oceanogr.* **51**: 1342–1354. <https://doi.org/10.4319/lo.2006.51.3.1342>
- Maerki, M., B. Wehrli, C. Dinkel, and B. Müller. 2004. The influence of tortuosity on molecular diffusion in freshwater sediments of high porosity. *Geochim. Cosmochim. Acta* **68**: 1519–1528. <http://doi.org/10.1016/j.gca.2003.09.019>
- Martin, P., L. Granina, K. Martens, and B. Goddeeris. 1998. Oxygen concentration profiles in sediments of two ancient lakes: Lake Baikal (Siberia, Russia) and Lake Malawi (East Africa). *Hydrobiologia* **367**: 163–174. <http://doi.org/10.1023/a:1003280101128>
- Matthews, D. A., and S. W. Effler. 2006. Long-term changes in the areal hypolimnetic oxygen deficit (AHOD) of Onondaga Lake: Evidence of sediment feedback. *Limnol. Oceanogr.* **51**: 702–714. [https://doi.org/10.4319/lo.2006.51.1\\_part\\_2.0702](https://doi.org/10.4319/lo.2006.51.1_part_2.0702)
- Matzinger, A., B. Müller, P. Niederhauser, M. Schmid, and A. Wüest. 2010. Hypolimnetic oxygen consumption by sediment-based reduced substances in former eutrophic lakes. *Limnol. Oceanogr.* **55**: 2073–2084. <https://doi.org/10.4319/lo.2010.55.5.2073>
- Müller, B., L. D. Bryant, A. Matzinger, and A. Wüest. 2012. Hypolimnetic oxygen depletion in eutrophic lakes. *Environ. Sci. Technol.* **46**: 9964–9971. <https://doi.org/10.1021/es301422r>
- Müller, B., and others. 2007. Present and past bio-available phosphorus budget in the ultra-oligotrophic Lake Brienz. *Aquat. Sci.* **69**: 227–239. <https://doi.org/10.1007/s00027-007-0871-8>
- Müller, B., and others. 2005. Internal carbon and nutrient cycling in Lake Baikal: Sedimentation, upwelling, and early diagenesis. *Global planet. Change* **46**: 101–124. <https://doi.org/10.1016/j.gloplacha.2004.11.008>
- Müller, B., T. Steinsberger, R. Schwefel, R. Gächter, M. Sturm, and A. Wüest. 2019. Oxygen consumption in seasonally stratified lakes decreases only below a marginal phosphorus threshold. *Sci. Rep.* **9**: 18054. <https://doi.org/10.1038/s41598-019-54486-3>
- Müller, B., R. Stierli, and A. Wüest. 2006. Phosphate adsorption by mineral weathering particles in oligotrophic waters of high particle content. *Water Resour. Res.* **42**: W10414. <https://doi.org/10.1029/2005wr004778>
- Och, L. M., and others. 2012. New insights into the formation and burial of Fe/Mn accumulations in Lake Baikal sediments. *Chem. Geol.* **330–331**: 244–259. <https://doi.org/10.1016/j.chemgeo.2012.09.011>
- Polerecky, L., N. Volkenborn, and P. Stief. 2006. High temporal resolution oxygen imaging in bioirrigated sediments. *Environ. Sci. Technol.* **40**: 5763–5769. <http://doi.org/10.1021/es060494l>
- Putyrskaya, V., E. Klemt, and S. Röllin. 2009. Migration of <sup>137</sup>Cs in tributaries, lake water and sediment of Lago Maggiore (Italy, Switzerland) – analysis and comparison with Lago di Lugano and other lakes. *J. Environ. Radioact.* **100**: 35–48. <https://doi.org/10.1016/j.jenvrad.2008.10.005>
- Randlett, M.-E., and others. 2015. Mineralization pathways of organic matter deposited in a river–lake transition of the Rhone River Delta, Lake Geneva. *Environ. Sci.: Processes Impacts* **17**: 370–380. <https://doi.org/10.1039/C4EM00470A>
- Remsen, C. C., E. C. Minnich, R. S. Stephens, L. Buchholz, and M. E. Lidstrom. 1989. Methane oxidation in Lake

- Superior sediments. *J. Great Lakes Res.* **15**: 141–146. [https://doi.org/10.1016/S0380-1330\(89\)71468-4](https://doi.org/10.1016/S0380-1330(89)71468-4)
- Schwalb, A., P. Hadorn, N. Thew, and F. Straub. 1998. Evidence for Late Glacial and Holocene environmental changes from subfossil assemblages in sediments of Lake Neuchâtel, Switzerland. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **140**: 307–323. [https://doi.org/10.1016/S0031-0182\(98\)00025-X](https://doi.org/10.1016/S0031-0182(98)00025-X)
- Schwefel, R., T. Steinsberger, D. Bouffard, L. D. Bryant, B. Müller, and A. Wüest. 2018. Using small-scale measurements to estimate hypolimnetic oxygen depletion in a deep lake. *Limnol. Oceanogr.* **63**: S54–S67. <https://doi.org/10.1002/lno.10723>
- Small, G. E., and others. 2013. Rates and controls of nitrification in a large oligotrophic lake. *Limnol. Oceanogr.* **58**: 276–286. <http://doi.org/10.4319/lo.2013.58.1.0276>
- Smith, D. A., and G. Matisoff. 2008. Sediment oxygen demand in the central basin of Lake Erie. *J. Great Lakes Res.* **34**: 731–744. [https://doi.org/10.1016/S0380-1330\(08\)71614-9](https://doi.org/10.1016/S0380-1330(08)71614-9)
- Sobek, S., and others. 2009. Organic carbon burial efficiency in lake sediments controlled by oxygen exposure time and sediment source. *Limnol. Oceanogr.* **54**: 2243–2254. <https://doi.org/10.4319/lo.2009.54.6.2243>
- Soomets, T., T. Kutser, A. Wüest, and D. Bouffard. 2019. Spatial and temporal changes of primary production in a deep peri-alpine lake. *Inland Waters* **9**: 49–60. <https://doi.org/10.1080/20442041.2018.1530529>
- Span, D., D. Arbouille, H. Howa, and J.-P. Vernet. 1990. Variation of nutrient stocks in the superficial sediments of Lake Geneva from 1978 to 1988. *Hydrobiologia* **207**: 161–166. <https://doi.org/10.1007/BF00041453>
- Steinsberger, T., B. Müller, C. Gerber, B. Shafei, and M. Schmid. 2019. Modeling sediment oxygen demand in a highly productive lake under various trophic scenarios. *PLoS One* **14**: e0222318. <https://doi.org/10.1371/journal.pone.0222318>
- Steinsberger, T., M. Schmid, A. Wüest, R. Schwefel, B. Wehrli, and B. Müller. 2017. Organic carbon mass accumulation rate regulates the flux of reduced substances from the sediments of deep lakes. *Biogeosciences* **14**: 3275–3285. <https://doi.org/10.5194/bg-14-3275-2017>
- Thevenon, F., S. B. Wirth, M. Fujak, J. Poté, and S. Girardclos. 2013. Human impact on the transport of terrigenous and anthropogenic elements to peri-alpine lakes (Switzerland) over the last decades. *Aquat. Sci.* **75**: 413–424. <https://doi.org/10.1007/s00027-013-0287-6>
- Thomsen, U., B. Thamdrup, D. A. Stahl, and D. E. Canfield. 2004. Pathways of organic carbon oxidation in a deep lacustrine sediment, Lake Michigan. *Limnol. Oceanogr.* **49**: 2046–2057. <https://doi.org/10.4319/lo.2004.49.6.2046>
- Torres, N. T., P. C. Hauser, G. Furrer, H. Brandl, and B. Müller. 2013. Sediment porewater extraction and analysis combining filter tube samplers and capillary electrophoresis. *Environ. Sci.: Processes Impacts* **15**: 715–720. <https://doi.org/10.1039/c3em00068k>
- Torres, N. T., and others. 2014. Early diagenetic processes generate iron and manganese oxide layers in the sediments of Lake Baikal, Siberia. *Environ. Sci.: Processes Impacts* **16**: 879–889. <https://doi.org/10.1039/C3EM00676J>
- Wehrli, B., A. Lotter, T. Schaller, and M. Sturm. 1997. High-resolution varve studies in Baldeggersee (Switzerland): Project overview and limnological background data. *Aquat. Sci.* **59**: 285–294. <http://doi.org/10.1007/BF02522359>
- Wenk, C. B., and others. 2013. Anaerobic ammonium oxidation (anammox) bacteria and sulfide-dependent denitrifiers coexist in the water column of a meromictic south-alpine lake. *Limnol. Oceanogr.* **58**: 1–12. <https://doi.org/10.4319/lo.2013.58.1.0001>
- Wirth, S. B., S. Girardclos, C. Rellstab, and F. S. Anselmetti. 2011. The sedimentary response to a pioneer geo-engineering project: Tracking the Kander River deviation in the sediments of Lake Thun (Switzerland). *Sedimentology* **58**: 1737–1761. <https://doi.org/10.1111/j.1365-3091.2011.01237.x>
- Zhu, G., and others. 2013. Hotspots of anaerobic ammonium oxidation at land–freshwater interfaces. *Nat. Geosci.* **6**: 103–107. <http://doi.org/10.1038/ngeo1683>
- Zimmermann, J. 2008. Der Walensee - eine sedimentologische Rekonstruktion seiner holozänen Ereignisgeschichte. Master thesis. Univ. Zurich & EAWAG Dübendorf.

### Acknowledgments

The authors acknowledge the Swiss Federal Office for the Environment for providing lake monitoring data. Data from Lake Geneva were provided by the *Commission Internationale pour la Protection des Eaux du Léman (CIPEL)* and the Information System of the *SOERE OLA, INRA* in Thonon-les-Bains (<http://si-ola.inra.fr>). We also gratefully acknowledge Michela Rogora, *CIPALIS*, and the *CNR Water Research Institute* in Verbania, Italy, for providing data on Lago Maggiore. We thank Martin Wessels from LUBW, Langenargen, Germany, for providing data and logistical support in investigations of Lake Constance. Further, work on Lake Constance was supported by Swiss National Science Foundation grants 200021\_146652 and 200020\_165517.

### Conflict of Interest

None declared.

Submitted 21 June 2019

Revised 11 February 2020

Accepted 16 July 2020

Associate editor: Bo Thamdrup