

This document is the accepted manuscript version of the following article:

Steinsberger, T., Schwefel, R., Wüest, A., & Müller, B. (2020). Hypolimnetic oxygen depletion rates in deep lakes: effects of trophic state and organic matter accumulation. Limnology and Oceanography. <https://doi.org/10.1002/lno.11578>

Hypolimnetic oxygen depletion rates in deep lakes:

Effects of trophic state and organic matter accumulation

Thomas Steinsberger^{1*}, Robert Schwefel^{2,3}, Alfred Wüest^{1,3}, Beat Müller¹

¹ Eawag, Swiss Federal Institute of Aquatic Science and Technology, CH-6047 Kastanienbaum, Switzerland

² UC Santa Barbara, 3015 Marine Science Building, Santa Barbara, CA 93106-6150, US.

³ Physics of Aquatic Systems Laboratory, Margaretha Kamprad Chair, École Polytechnique Fédérale de Lausanne, Institute of Environmental Engineering, Lausanne, Switzerland.

ORCID: Thomas Steinsberger: 0000-0002-0751-5424; Robert Schwefel: 0000-0003-1610-4181; Beat Müller: 0000-0003-3696-9035; Alfred Wüest: 0000-0001-7984-0368

Author Email address:

* Corresponding Author: Thomas Steinsberger: thomas.chwalek@eawag.ch

Robert Schwefel: robertschwefel@ucsb.edu

Alfred Wüest: alfred.wueest@eawag.ch

Beat Müller : beat.mueller@eawag.ch

Running Head: Hypolimnetic O₂ depletion in deep lakes

Keywords : Carbon Mineralization, Porewater, Oxygen depletion, deep lakes, lake phosphorous, water column, sediment

23 ***Abstract***

24 This study investigated the consumption of oxygen (O₂) in eleven European lakes ranging from 48
25 m to 372 m deep. In lakes less than ~100 m deep, the main pathways for O₂ consumption were
26 organic matter (OM) mineralization at the sediment surface and oxidation of reduced compounds
27 diffusing up from the sediment. In deeper lakes, mineralization of OM transported through the
28 water column to the sediment represented a greater proportion of O₂ consumption. This process
29 predominated in the most productive lakes but declined with decreasing total phosphorous (TP)
30 concentrations and hence primary production, when TP concentrations fell below a threshold value
31 of ~10 mg P m⁻³. Oxygen uptake by the sediment and the flux of reduced compounds from the
32 sediment in these deep lakes were 7.9 - 10.6 mmol m⁻² d⁻¹ and 0.6 - 3.6 mmol m⁻² d⁻¹, respectively.
33 These parameters did not depend on the lake's trophic state but did depend on sedimentation rates
34 for the primarily allochthonous or already degraded OM. These results indicate that in lakes deeper
35 than ~100 m, mineralization of autochthonous OM is mostly complete by the time of sedimentary
36 burial. This explains why hypolimnetic O₂ concentrations improve more rapidly following TP load
37 reduction in deeper lakes relative to shallower lakes, where larger sediment based O₂ consumption
38 by settled OM and release of reduced substances may inhibit the restoration of hypolimnetic O₂
39 concentrations.

40

1. *Introduction*

Sufficient O₂ concentrations in the hypolimnia of lakes are essential for maintaining habitats, supporting fish reproductive cycles, and ensuring reliable drinking water sources. Oxygen concentrations are thus a key parameter for lake management. In the middle of the 20th century, increasing phosphorous (P) concentrations contributed to drastic increases in primary production and subsequent transport of OM to deep hypolimnetic waters. Aerobic mineralization of OM and the release of anaerobically produced compounds such as methane (CH₄) and ammonium (NH₄⁺) into bottom waters are common contributors to hypolimnetic O₂ depletion. Lacustrine OM mineralization generally occurs in three different domains. These include (1) the water column during settling of organic particles (Livingstone and Imboden 1996), (2) sediment surfaces subject to intensive remineralization and biological reworking (Burns 1995), and (3) underlying, deeper sediments that host gradual but continuous production of reduced substances (Carignan and Lean 1991; Steinsberger et al. 2017). O₂ consumption rates for each of these locations have been respectively termed (1) water column mineralization rate in the hypolimnion (WCM), (2) sediment oxygen uptake (SOU) and (3) flux of reduced compounds from the sediment, F_{red}, as converted to O₂ consumption equivalents (Matzinger et al. 2010; Müller et al. 2012). The areal hypolimnetic mineralization (AHM) represents the sum of all three O₂ consumption processes over the whole hypolimnion.

For fully productive (eutrophic) lakes of less than ~100 m depth, SOU (OM mineralization on the sediment surface) and F_{red} (reduced compounds diffusing from interstitial fluids in deeper sediments) represent the primary sites of O₂ consumption (Matthews and Effler 2006; Müller et al. 2012). In deeper (>100 m) lakes such as the Great Laurentian Lakes, WCM and SOU govern O₂

depletion (Charlton 1980). Studies of Lakes Michigan (Thomsen et al. 2004), Erie (Edwards et al. 2005; Smith and Matisoff 2008) and Superior (Li et al. 2012) have further demonstrated the importance of SOU and/or WCM as O₂ sinks in these lakes. A study of the highly productive and deep Lake Geneva by Schwefel et al. (2018) showed that, as with the Great Lakes, O₂ consumption primarily occurred in the water column rather than near or in the sediment.

In general, deep lakes such as Baikal or Superior store enough O₂ in their large hypolimnia to prevent anoxia in bottom waters, assure aerobic OM remineralization and limit the production of reduced substances (Li et al. 2012; Müller et al. 2005). Even in the very productive and deep Lake Geneva, which exhibits a TP concentration of ~24 mg P m⁻³ (average from 2008 to 2018) and a carbon assimilation rate of ~250 g C m⁻² yr⁻¹ (Anneville and Pelletier 2000; Soomets et al. 2019), sediment characteristics resemble those of an oligotrophic lake with low TOC content (~1%), very small fluxes of CH₄ and NH₄⁺ from the sediment and limited O₂ consumption at the sediment-water interface (Schwefel et al. 2018; Steinsberger et al. 2017). The high fraction of WCM (~70%) on the total AHM diminishes the oxidizing capacity of the settling material on its way to the sediment. Given the mostly, already degraded OM, SOU only contributed about 27% to AHM. The O₂ made available by a large hypolimnion for extensive oxic mineralization of settling OM results in only a marginal amount of OM burial in sediments.

In shallower lakes (<100 m depth) with similar productivity, shorter settling distances may limit (oxic) mineralization in the water column. This could lead to accumulation and burial of large volumes of mostly un-mineralized OM in sediments. The subsequent slow but persistent release of reduced compounds generated by anaerobic OM degradation further consumes O₂ and often impedes later restoration measures (Wehrli et al. 1997).

In shallow lakes, this “sediment memory effect” or “sediment legacy” and associated release of reduced substances from the sediment may account for up to 80% of the O₂ depletion in the hypolimnion even with measures aimed at reestablishing oligotrophic conditions (Carey et al. 2018; Matthews and Effler 2006). Although this effect has been documented in lakes of up to ~100 m depth, its potential impact on deeper lakes is not as well understood (Matzinger et al. 2010). In most deep lakes, the contribution of different O₂ sinks to overall hypolimnion O₂ depletion remains debated or unknown.

This study aimed to determine the location and intensity of O₂ consumption in eight lakes of >100 m depth and three lakes of <100 m depth. We determined O₂ consumption rates at the sediment surface, fluxes of reduced compounds and O₂ depletion rates in the water column using lake monitoring data, O₂ microprofile measurements and simultaneous sediment porewater analysis. Oxygen consumption was characterized and interpreted in terms of spatial location, trophic state, hypolimnion depth and accumulation of OM.

2. *Material and Methods*

2.1. *Site description*

Sediment cores of the lakes characterized in Table 1 were retrieved at the deepest locations of each lake except for Lakes Maggiore, Lucerne and Constance. For Lake Maggiore, sediment cores were collected from 273 m depth at the deepest site of the Swiss part of the lake. For Lake Lucerne, sediment was collected in the deepest part of the Vitznau basin at 150 m water depth. For Lake Constance, sediment was collected at 150 m depth in the main basin. All sediment cores were

collected during the lake's stratified period (between May and October). Table S1 summarizes sampling details, location and dates. The two shallowest lakes, Hallwil and Baldegg, both experience recurring hypolimnetic hypoxia caused by high primary production. Both lakes are artificially aerated during summer stratification and mixing has been supported by bubble plumes during overturn since the 1980s. Hypolimnetic O₂ sinks are thus influenced by artificially elevated O₂ concentrations introduced by lake restoration efforts.

2.2. Sampling procedure

For each sampling site, four sediment cores were collected with a gravity corer. These cores were subjected to individual measurements of sediment porewater, CH₄, O₂ micro-optode profiles and bulk analysis of sediment chemical and physical properties.

Sediment cores subjected to bulk analysis were stored in a cold room (4°C) until processing. The cores were sampled as 0.5 to 1 cm sections, weighed and freeze-dried. Water content was calculated from the weight difference before and after drying. Sediment porosity was estimated from water content, sediment density and TOC content (Och et al. 2012). After grinding with an agate mortar, the sediment was further analyzed for TOC/TN with a Euro EA 3000 (HEKAtech). Total organic carbon mass accumulation rates (TOC-MAR) were determined for each site at depths between 1-10 cm sediment depths in 5 mm spatial resolution. Estimation of TOC-MAR followed detailed methods outlined in Steinsberger et al. (2017). Sedimentation rates were derived from literature sources or estimated from ²¹⁰Pb and ¹³⁷Cs measurements as reported from prior analyses of the respective lake sediments.

For CH₄ and sediment porewater analysis, 60 cm long PVC liners with predrilled and taped holes (diameter 1.2 cm and 2 mm, respectively) were used. Syringes with clipped tips were used to

sample 2 cm³ of sediment aliquots for CH₄ analysis. Aliquots were directly transferred to 113.7 mL flasks containing 2 mL of 7 M NaOH, then capped, stored and prepared for GC-MS headspace analysis. The sampling resolution was 1 cm for the upper 10 cm, 2 cm for the middle 10 cm and 3 cm from 20 to 35 cm sediment depth.

Sediment porewater was extracted using MicroRhizon filter tubes of 1.5 mm diameter (Torres et al. 2013). Punching through the taped pre-drilled holes allowed for withdrawal of 10 – 50 µL of sediment porewater. This porewater was transferred to a flip tube and immediately injected onto a capillary electrophoresis CE-C⁴D column (Section 2.3). Relative to the sediment-water interface, porewater was sampled every 5 mm for the upper 5 cm, every 1 cm for the following 5 cm and every 2 cm from 10 to 20 cm sediment depth.

The fourth core was used to sample bottom water for O₂ concentration using the Winkler method. Vertical concentration profiles of O₂ in the sediment cores were performed with a PreSens O₂ micro-optode mounted onto an automated micromanipulator (Section 2.3).

2.3. Oxygen and sediment porewater analysis

Sediment porewater analysis and O₂ concentration profiles were measured directly on site at each lake. Sediment porewater samples were analyzed for anions and cations with two portable capillary electrophoresis (CE) devices, each with a capacitively coupled contactless conductivity detector (C⁴D) calibrated directly on site (Kubáň et al. 2007). These devices were set at 15 kV and 0.5 µA to achieve complete separation of cations (NH₄⁺, K⁺, Ca²⁺, Na⁺, Mg²⁺, Fe²⁺, Mn²⁺) and anions (Cl⁻, NO₃⁻, NO₂⁻, SO₄²⁻, H₂PO₄⁻) within eight minutes (Torres et al. 2013). All calibration standards and the background electrolyte buffer were prepared from corresponding salts with ultrapure water

(Merck) immediately prior to each sampling campaign. These were checked using a multi-cation standard (Fluka) and a multi-anion standard (Roth). Standard deviations of all calibration measurements were <5%.

The diffusive fluxes of reduced substances (J_{CH_4} , J_{NH_4} , $J_{Mn(II)}$, $J_{Fe(II)}$) from the sediments were calculated from porewater concentration gradients (Supporting Information Fig. S1). The one-dimensional steady-state diffusion model by Epping and Helder (1997) was adapted to calculate fluxes:

$$J_i = -\frac{D}{G} * \frac{dC_i}{dx} \quad (1)$$

For J in $mmol\ m^{-2}\ d^{-1}$, D represents the diffusion coefficient based on temperature corrected values from Li and Gregory (1974), G represents porosity corrected formation factor ($1.02 * porosity^{-1.81}$) for clay-silt sediment and $\frac{dC_i}{dx}$ represents the concentration gradient of species i (Maerki et al. 2004). Based on the redox stoichiometry (Eq. 2), the estimated fluxes were converted to O_2 equivalents ($g\ O_2\ m^{-2}\ d^{-1}$) (Matzinger et al. 2010; Müller et al. 2012) as follows:

$$F_{red} = 2 * J_{CH_4} + 2 * J_{NH_4} + 0.5 * J_{Mn(II)} + 0.25 J_{Fe(II)} \quad (2)$$

O_2 measurements were conducted with a PreSens O_2 micro-optode with a flat tip size of 230 μm , a detection limit of 0.64 $\mu mol\ O_2\ L^{-1}$ and a response time <10 s. The micro-optode was mounted to an automated micromanipulator. O_2 profiles were taken between 2 cm above and 3 cm below the sediment-water interface at 100 μm intervals and a penetration speed of 100 $\mu m\ s^{-1}$ stopping for each measurement, completing a full profile in <9 minutes. The O_2 penetration depth was defined as the sediment depth where O_2 concentrations dropped below 3 $\mu mol\ O_2\ L^{-1}$. The O_2 micro-optode was calibrated in the laboratory with a two-point calibration. During the on-site measurements,

standard deviations for O₂ measurements was typically ~11%. Up to nine replicate measurements were taken at different positions for each core.

Simultaneous O₂ measurements were not performed in Lakes Hallwil, Geneva or Constance. Data from Bierlein et al. (2017) (Lake Hallwil), Schwefel et al. (2018) (Lake Geneva) and Sobek et al. (2009) (Lake Constance) were used for comparison instead.

The SOU flux was calculated similar to the flux of reduced substances (Eq. 1). Fluxes of the other two main electron acceptors, sulfate (SO₄²⁻) and nitrate (NO₃⁻), were converted to O₂ equivalents and added to SOU (Müller et al. 2007):

$$SOU = J_{O_2} + 1.25 * J_{NO_3} + 2 * J_{SO_4} \quad (3)$$

2.4. Areal hypolimnetic and water column mineralization

Based on monthly monitoring data, average AHMs for the past 10 years (2007-2017) were calculated from differences in hypolimnetic O₂ content (>15 m water depth) as measured between maximum values after winter mixing (February to April) and minimum values from October to November.

If reduced substances such as NH₄⁺ and/or CH₄ occurred in bottom waters, they were included in the budget with their O₂ equivalents (Matzinger et al. 2010). WCM was calculated by subtracting SOU and F_{red} from AHM as:

$$AHM = SOU + F_{red} + WCM \quad (4)$$

$$WCM = AHM - (2 * J_{CH_4} + 2 * J_{NH_4} + 0.5 * J_{Mn(II)} + 0.25 * J_{Fe(II)}) - (J_{O_2} + 1.25 * J_{NO_3} + 2 * J_{SO_4}) \quad (5)$$

2.5. Comparability and limitations of SOU , F_{red} and WCM parameters

Hypolimnetic O_2 sinks fundamentally depend on aerobic or anaerobic OM mineralization. Although the scale and biogeochemical processes may differ, these sinks are comparable because all fluxes are normalized to the hypolimnion area and expressed as O_2 equivalents ($mmol\ O_2\ m^{-2}\ d^{-1}$). However, spatial and temporal variation may influence the individual size of the O_2 sinks.

Sediment-based O_2 sinks, F_{red} and SOU are controlled by the amount of settled OM and are therefore sensitive to spatial effects such as sediment focusing. Gravitational slumping or wind forcing can resuspend fine-grained, organic-rich material and transport it from littoral areas to deeper parts of lakes (Bloesch and Uehlinger 1986; Carignan and Lean 1991; Lehman 1975). Seasonal variations may also influence SOU since hypolimnetic O_2 concentrations tended to decrease until the end of the stratified season. A single O_2 measurement may therefore over- or underestimate annual average SOU values. Seasonal variation in F_{red} is less relevant since it depends only indirectly on fluctuating OM input or overlying O_2 concentrations (Frenzel et al. 1990; Schwefel et al. 2018; Steinsberger et al. 2019). By contrast, seasonal variation in primary production and associated settling and mineralization throughout the hypolimnetic water column strongly influence WCM but this variation is imbedded in yearly average AHM calculations.

3. Results

3.1. Fluxes of reduced substances from the sediment

Typical sediment concentration profiles of CH_4 , NH_4^+ , Fe(II) , Mn(II) , O_2 , NO_3^- and SO_4^{2-} are shown in the Supporting Fig. S1. Table 2 summarizes fluxes (J) of all measured species converted to O_2 equivalents from the eleven lakes.

In lakes deeper than 100 m, F_{red} showed similar low values of between 0.6 and 3.6 $\text{mmol O}_2 \text{ m}^{-2} \text{ d}^{-1}$. Shallower Lakes Hallwil, Baldegg and Aegeri exhibited higher F_{red} values of between 4.0 $\text{mmol O}_2 \text{ m}^{-2} \text{ d}^{-1}$ (Lake Aegeri) and 13.8 $\text{mmol O}_2 \text{ m}^{-2} \text{ d}^{-1}$ (Lake Baldegg).

CH_4 and NH_4^+ represented the largest contributions to F_{red} . $J_{\text{Fe(II)}}$ and $J_{\text{Mn(II)}}$ were comparatively negligible and contributed on average only 1% and 2%, respectively, even though deeper parts of the sediment (>10 cm depth) exhibited high sediment porewater concentrations. Lago Maggiore sediment porewater for example contained 0.7 $\text{mmol Mn(II) L}^{-1}$ and Lake Walensee sediment porewater contained 0.8 $\text{mmol Fe(II) L}^{-1}$. In lakes of >100 m depth, NH_4^+ and CH_4 porewater concentrations were at or below detection limits in the uppermost sediment layers (sometimes down to 5 cm). These values increased to 0.3 – 0.6 mmol L^{-1} (NH_4^+) and 2 – 7 mmol L^{-1} (CH_4) downcore (see Supporting Information Fig. S1). Except for Lago Maggiore, NH_4^+ did not reach the sediment surface but was oxidized within the uppermost centimeter. Here, J_{CH_4} represented up to 99% of F_{red} (Fig. 1). In the shallower Lakes Hallwil, Baldegg and Aegeri (<100 m depth), in addition to J_{CH_4} (54% to 62%) J_{NH_4} was responsible for the remaining fraction of F_{red} (Fig. 1; ~40%).

3.2. Sediment oxygen uptake and penetration depths

Among lakes deeper than 100 m, SOU rates ranged only from 7.9 to 11.0 $\text{mmol O}_2 \text{ m}^{-2} \text{ d}^{-1}$. However, SOU values in Lakes Maggiore (16.4 $\text{mmol O}_2 \text{ m}^{-2} \text{ d}^{-1}$) and Walensee (17.6 mmol O_2

233 $\text{m}^{-2} \text{d}^{-1}$) were exceptionally high and resembled the SOU average of $19.0 \text{ mmol O}_2 \text{ m}^{-2} \text{d}^{-1}$ measured
 234 from the shallower and more productive Lakes Hallwil and Baldegg.

235 All lakes showed high O_2 concentrations in the hypolimnion ($>0.19 \text{ mmol O}_2 \text{ L}^{-1}$, Supporting
 236 Information Fig. S2). However, penetration depths into the sediment were low (between 2.0 ± 1.0
 237 mm in Lake Baldegg and 6.3 ± 0.8 mm in Lake Brienz, see Table 4) and did not correlate with
 238 bottom water O_2 concentrations. J_{O_2} ranged from $4.4 \text{ mmol O}_2 \text{ m}^{-2} \text{d}^{-1}$ in Lake Geneva to 18.8 mmol
 239 $\text{O}_2 \text{ m}^{-2} \text{d}^{-1}$ in Lake Baldegg (Table 2). J_{O_2} contributed an average of $78 \pm 13\%$ to total SOU. J_{NO_3}
 240 and J_{SO_4} varied between $0.3 - 2.4 \text{ mmol O}_2 \text{ m}^{-2} \text{d}^{-1}$ and $0.3 - 2.0 \text{ mmol O}_2 \text{ m}^{-2} \text{d}^{-1}$ and respectively
 241 represented $6.7 \pm 3.2 \%$ and $11.2 \pm 9.1 \%$ of SOU. For lakes with high hypolimnetic NO_3^- and/or
 242 SO_4^{2-} concentrations ($>0.4 \text{ mmol L}^{-1}$; Lakes Geneva and Thun), combined J_{NO_3} and J_{SO_4} represented
 243 up to 45% of SOU (Fig. 1).

244

245 **3.3. Mineralization in the water column**

246 WCM rates varied between 2.1 and $27.4 \text{ mmol O}_2 \text{ m}^{-2} \text{d}^{-1}$. As a measure of OM mineralization
 247 within the hypolimnetic water column, WCM depends on the lakes' volume and primary
 248 production and thus its TP concentration. Oligotrophic lakes ($\text{TP} < 10 \text{ mg P m}^{-3}$) that have not
 249 experienced eutrophication in the past (Walensee, Lakes Thun, Brienz and Aegeri) exhibited low
 250 WCM values of $2.1 \text{ mmol O}_2 \text{ m}^{-2} \text{d}^{-1}$ (Lake Thun) and $4.2 \text{ mmol O}_2 \text{ m}^{-2} \text{d}^{-1}$ (Lake Aegeri). WCM
 251 could not be estimated from Walensee because the sum of F_{red} ($3.6 \text{ mmol O}_2 \text{ m}^{-2} \text{d}^{-1}$) and SOU (17.6
 252 $\text{mmol O}_2 \text{ m}^{-2} \text{d}^{-1}$) already equaled measured AHM ($20.0 \text{ mmol O}_2 \text{ m}^{-2} \text{d}^{-1}$; Table 3). Lago Maggiore
 253 similarly showed a very low WCM value ($3.9 \text{ mmol O}_2 \text{ m}^{-2} \text{d}^{-1}$). Both lakes have exceptionally
 254 steep walls which cause substantial sediment focusing leading to elevated SOU and F_{red} at the

central coring sites. In the oligotrophic but recovering Lake Constance, WCM values resembled those measured from the mesotrophic Lake Neuchâtel (14.4 and 11.0 mmol O₂ m⁻² d⁻¹, respectively). The highest WCM rate was observed from the most productive very deep Lake Geneva (27.4 mmol O₂ m⁻² d⁻¹). The shallower but similarly productive Lakes Baldegg and Hallwil exhibited lower respective WCM rates of 5.2 and 10.5 mmol O₂ m⁻² d⁻¹ (Table 3).

4. Discussion

4.1. Fluxes of reduced substances from the sediment

Very low F_{red} values (<0.2 mmol O₂ m⁻² d⁻¹) were previously reported in lakes >100 m deep such as Lakes Superior (Li et al. 2012; Remsen et al. 1989), Michigan (Thomsen et al. 2004) and Baikal (Och et al. 2012; Torres et al. 2014). These lakes are characterized by low OM accumulation rates and high O₂ penetration depths resulting in prolonged (up to several decades) aerobic mineralization of the settled OM. Consequently, fluxes of CH₄, NH₄⁺, Fe(II) Mn(II) and hence, F_{red}, are close to zero.

The deep lakes investigated by this study (>100 m) exhibited distinctly higher F_{red} values (average 2.6 ± 1.6 mmol O₂ m⁻² d⁻¹) and showed a consistent dependence on mean lake depth (Fig. 2) (Steinsberger et al. 2017). These were dominated by CH₄ fluxes which constituted up to 99% of F_{red} (Table 2, Figure 1). NH₄⁺ contributions were negligible in all lakes that have never surpassed mesotrophic production in the past. Deeper sedimentary horizons (>10 cm) however exhibited elevated CH₄ and NH₄⁺ concentration of up to 7.0 mmol L⁻¹ and 0.6 mmol L⁻¹, respectively. These resembled values measured from river deltas or from shallower, more productive lakes (Randlett

et al. 2015; Sobek et al. 2009; Steinsberger et al. 2017). In contrast to the Great Lakes, the deep peri-alpine lakes investigated by this study accumulated inorganic matter from weathering of rocks and OM from their watersheds. This gradually degrading allochthonous OM is diluted with additional mineral deposits resulting in small concentrations of TOC (Supporting Information Fig. S3). High sedimentation rates lead to rapid OM burial in the anoxic sediments. A laboratory study by Grasset et al. (2018) showed that the mineralization of allochthonous OM by methanogenesis, although slower than autochthonous OM degradation, can yield similar overall CH_4 production. This finding helps to explain high porewater accumulation of CH_4 and NH_4^+ in deeper sediments collected from Walensee and Lago Maggiore.

Consequently, upward diffusing CH_4 and NH_4^+ comes into contact with the main electron acceptors O_2 , NO_3^- and SO_4^{2-} , and is (partially) oxidized within the sediment. In consequence, a substantial fraction of O_2 , NO_3^- or SO_4^{2-} consumption could be attributed to F_{red} oxidation rather than by direct aerobic mineralization of OM.

In productive lakes <100 m deep (Lakes Baldegg and Hallwil), porewater concentration gradients of CH_4 , NH_4^+ , Fe(II) and Mn(II) extended to the sediment-water interface. F_{red} attained high rates of $5.6 \text{ mmol O}_2 \text{ m}^{-2} \text{ d}^{-1}$ (Lake Hallwil) and $13.8 \text{ mmol O}_2 \text{ m}^{-2} \text{ d}^{-1}$ (Lake Baldegg) fueled by the high primary production and mineralization of large amounts of TOC deposited during past anoxic periods (Steinsberger et al. 2019).

4.2. Sediment oxygen uptake dependence on organic matter

O_2 is depleted by the initial aerobic degradation of OM at the sediment surface and by oxidation of reduced compounds produced during anaerobic mineralization within the sediment (Steinsberger

et al. 2017). Lakes >100 m deep exhibited SOU in the narrow 7.9 – 11.0 mmol O₂ m⁻² d⁻¹ range. Walensee and Lago Maggiore exhibited exceptionally high SOU values of 17.6 and 16.4 mmol O₂ m⁻² d⁻¹ (respectively), which resemble values observed in the shallower and more productive Lakes Hallwil and Baldegg (14.8 and 23.2 mmol O₂ m⁻² d⁻¹; Table 3). The steep slopes of Walensee and Lago Maggiore may increase OM deposition rates by focusing sediment into the central parts of the basin which in turn increases sediment-related O₂ consumption. The sediment area relevant for SOU and F_{red} is thus substantially smaller than the hypolimnion area used for AHM estimation. Equation 4 above may thus systematically underestimate WCM values for lakes having steep walls. Sediment related O₂ consumption depends on several key parameters such as bottom water O₂ concentrations, O₂ penetration depth, TOC mass accumulation rate and OM susceptibility to mineralization (Bryant et al. 2010; Gudas et al. 2010; Polerecky et al. 2006; Sobek et al. 2009). All lakes >100 m deep exhibited sediment TOC contents of 0.5 - 1.5%, whereas shallower lakes exhibited sediment TOC concentrations of 3.4 - 3.5 % (Supporting Information Fig. S3, Tab. 4). The more rapid mineralization rates for autochthonous OM relative to the more recalcitrant terrestrial OM (Sobek et al. 2009) suggest that the former is preferentially mineralized in the water column and then deposited mostly as already degraded OM onto the sediment for lakes deeper than 100 m. Schwefel et al. (2018) demonstrated this for the productive Lake Geneva while Carstens et al. (2013) did so for the oligotrophic Lake Brienz. Consequently, we observed a decreasing trend of SOU with increasing mean lake depths, similar to F_{red} (see Fig. 2). Elevated TOC concentrations in sediment of lakes <100 m deep is consistent with the interpretation of limited degradation within the water column and more vigorous O₂ consumption in the sediment surface leading to burial of large amounts of partially mineralized OM in sediments.

Table 4 shows that sediment mass accumulation rates in >100 m deep lakes (Maggiore, Thun, Brienz and Walensee) unexpectedly resembled or even exceeded that observed from shallower lakes. This reflects elevated areal contributions of allochthonous material from watersheds (Thevenon et al. 2013; Wirth et al. 2011) as well as sediment focusing by funnel- or trench-shaped lake basins (Blais and Kalff 1995; Lehman 1975). The exposure time of OM to O₂ falls well below one year. Combined with elevated MAR, these conditions bury significant amounts of OM in anoxic sediment before degradation can occur. The elevated TOC-MAR lead to anaerobic mineralization in the deeper sediment which then generates CH₄ and NH₄⁺ observed at high concentrations in the sediment porewater. These species can delay lake recovery from eutrophication with respect to O₂ concentrations. However, all of the Fe(II) and Mn(II) and most of the NH₄⁺ are oxidized within the sediment. These do not reach the sediment-water interface and thereby increase SOU (Supplementary Fig. S1).

The range of SOU determined from lakes >100 m deep investigated here (11.3 ± 3.7 mmol O₂ m⁻² d⁻¹) is consistent with fluxes reported by literature sources for other large and deep lakes. Thomsen et al. (2004) reported a SOU rate of 7.7 ± 2.1 mmol O₂ m⁻² d⁻¹ for Lake Michigan while Li et al. (2012) reported a SOU rate of 6.1 ± 1.4 mmol O₂ m⁻² d⁻¹ for Lake Superior. Lower SOU values were reported for the ultra-deep oligotrophic Lake Baikal. These include an overall value of 3.3 ± 1.1 mmol O₂ m⁻² d⁻¹ (Müller et al. 2005), 1.4 to 5.6 mmol O₂ m⁻² d⁻¹ for the South Basin (Och et al. 2012) and 1.1 mmol O₂ m⁻² d⁻¹ for the North Basin (Martin et al. 1998). O₂ penetration depths determined for the deep lakes analyzed here (2.6 – 6.3 mm, Table 4) resembled values previously reported for Lake Constance (1.5 – 6.3 mm; Sobek et al. (2009) and Frenzel et al. (1990) but were distinctly lower than those reported for Lakes Michigan (21 ± 2.5 mm), Superior (40 – 110 mm) and Baikal (20 – 44 mm). This evidences the high consumption of O₂ within the first millimeters

of sediment rendering O_2 penetration depths more similar to those in the fully productive and shallower Lake Baldegg (0.9 – 2.0 mm). Lakes Michigan and Superior exhibit low SOU rates and corresponding higher O_2 penetration depths that result from low OM sedimentation rates, intense bioturbation and extensive Fe cycling with only small NH_4^+ and insignificant CH_4 production. For lakes analyzed in this study, excess O_2 that could reach deeper parts of the sediment due to the absence of reactive OM is apparently consumed by a front of upward diffusing reduced compounds produced by anaerobic mineralization of OM. Decreasing porewater concentrations of reduced species in the uppermost layers of sediment reported here offer further evidence of rapid anaerobic turnover (Maerki et al. 2006).

Fluxes of NO_3^- into the sediment indicating mineralization of OM by denitrification ranged from 0.32 to 2.36 $mmol\ O_2\ m^{-2}\ d^{-1}$ and thereby only a small fraction of total SOU (between 2.9% in Lago Maggiore and 12.5% in Lake Lucerne). Li et al. (2012) and Thomsen et al. (2004) report similarly low estimates for Lakes Superior and Michigan (<5%). Overall NO_3^- removal rates (coupled nitrification-denitrification) however may be higher due to a hidden contribution from anaerobic NH_4^+ oxidation with low concentrations but high turnover rates of NO_3^-/NO_2^- to N_2 (e.g., anammox; (Crowe et al. 2017; Small et al. 2013). In fact, at least in five of the lakes investigated here (Walensee, Neuchâtel, Thun, Brienz and Lucerne) a rapid decline in NH_4^+ concentrations (with NH_4^+ consumption rates of 0.08 - 0.46 $mmol\ m^{-2}\ d^{-1}$) well below the O_2 penetration depth was evident. These conditions clearly indicate anaerobic NH_4^+ oxidation (Zhu et al. 2013). As NO_3^- penetrates deeper into the sediment than O_2 (5 mm – 45 mm), anammox bacteria might be actively oxidizing NH_4^+ (Wenk et al. 2013).

Fluxes of SO_4^{2-} ranged from 0.32 to 2.70 $mmol\ O_2\ m^{-2}\ d^{-1}$ (average 1.19 $mmol\ O_2\ m^{-2}\ d^{-1}$), and were well within the range of SO_4^{2-} reduction rates given by Holmer and Storkholm (2001). In

general, SO_4^{2-} fluxes contributed ~11% to SOU, while in Lake Geneva and Lake Thun, due to high loads from the catchment geology, SO_4^{2-} fluxes contributed about one third to SOU. The full extent of SO_4^{2-} reduction induced by OM mineralization is likely masked by (cryptic) internal sulfide re-oxidation and sulfite complexation processes (Holmkvist et al. 2011). Hence, SO_4^{2-} fluxes likely underestimate actual SO_4^{2-} reduction rates.

4.3. Mineralization in the water column

In the hypolimnetic water column, OM is mineralized by a large variety of microbes that consume O_2 (Carstens et al. 2012). Subsequent O_2 consuming processes at the sediment-water interface depend on excess OM passing through the water column. Therefore, depending on the main mineralization sites of OM (hypolimnetic water, sediment surface, deep sediment) lakes show characteristic patterns in the three O_2 sinks (WCM, SOU and F_{red}).

Livingstone and Imboden (1996) proposed that WCM represents a significant proportion of AHM for large lakes with a high mean hypolimnion depth. This is true for productive lakes such as Geneva (77%) and Neuchâtel (45%) and lakes recovering from a eutrophic phase, such as Lakes Constance (58%) and Lucerne (43%) (Tab. 3 and Fig. 3). This is in agreement with the modeled WCM value of Lake Geneva by Schwefel et al. (2018). However, lakes that never experienced high productivity periods, such as the oligotrophic Lakes Brienz, Thun and Walensee, did not show elevated WCM rates. These lakes instead showed WCM contributions to AHM of only ~14%, ~29% and ~0%, respectively. They also exhibited low sediment TOC content but a high rate of sediment OM accumulation caused by high loads of allochthonous particles (Bechtel and Schubert 2009; Finger et al. 2007). These lakes also mineralize a relatively small amount of OM within the

hypolimnetic water column compared to that deposited at the sediment surface (Fig. 3). The high SOU and F_{red} rates indicate mineralization of allochthonous OM. The very low WCM of Walensee and Lago Maggiore reflect high SOU and F_{red} caused by sediment focusing (Blais and Kalff 1995). Lake Baldegg exhibited WCM of only $5.2 \text{ mmol O}_2 \text{ m}^{-2} \text{ d}^{-1}$ or 12% of AHM. This likely reflects the shallower lake depth and brief mineralization window for settling autochthonous OM.

WCM primarily depends on the export of autochthonous OM from the productive zone to the hypolimnion. Because it originates from primary production, it must be therefore related to TP for concentration ranges where TP availability influences primary production. In order to compare WCM between lakes of different depths, we normalized WCM to the mean hypolimnion depth, z_H to obtain a relationship with TP (Fig. 4). In all non-aerated lakes, WCM increases in proportion to TP. For a TP range of $10 - 15 \text{ mg P m}^{-3}$, WCM levels off indicating that additional TP exerts limited influence on primary production and thus on OM export to the hypolimnion and subsequent O_2 consumption. Below a TP concentration of $\sim 10 \text{ mg P m}^{-3}$, WCM covaries closely with TP concentration. This is in line with the observation of Müller et al. (2019) that AHM in seasonally stratified lakes increases approximately in proportion to the areal amount of bio-available P but levels off above this threshold value. O_2 consumption in lakes $>100 \text{ m}$ deep may immediately respond to a decrease of TP for concentrations below this threshold. Decreasing TP concentration in Lake Geneva by half could initiate a substantial reduction in WCM, the lake's primary O_2 sink. By contrast, minor interventions to reduce TP levels for Lakes Neuchâtel and Constance would readily influence WCM and thereby decrease total AHM.

5. Conclusion

Diffusive fluxes of reduced substances from and flux of O_2 into the sediments were measured in lakes with depths ranging from 48 to 372 m. Areal hypolimnetic mineralization rates were calculated from (monthly) O_2 monitoring data and used to estimate the water column mineralization rates. This analysis showed that:

- despite high O_2 concentrations ($>190 \mu\text{mol } O_2 \text{ L}^{-1}$) in bottom waters of all lakes, O_2 penetrated sediments only to depths of 2.6 to 6.3 mm resulting in high O_2 fluxes. In lakes deeper than 100 m, reduced substances from anaerobic mineralization of OM in the sediment constituted a major fraction of the sediment oxygen uptake and prevent O_2 from penetrating deeper into the sediment.
- Most lakes deeper than 100 m exhibited similar levels of sediment O_2 consumption by SOU and F_{red} . This suggests that OM deposited at the sediment surface was already largely mineralized. The O_2 reservoir of the more extensive hypolimnia sufficed to degrade most of the settling OM. Hence, SOU and F_{red} values did not depend on trophic state for lakes deeper than 100 m.
- WCM represented a major contribution to AHM in lakes of >100 m depth. This contribution decreased with TP when less autochthonous OM was available for O_2 consumption. WCM increased with increasing TP concentrations until a threshold of 10 - 15 mg P m^{-3} was reached. Above this threshold, WCM levelled off. Reducing TP concentrations in productive lakes deeper than 100 m to below this threshold could therefore exert a strong influence on AHM.

6. *Acknowledgements*

The authors acknowledge the Swiss Federal Office for the Environment for providing lake monitoring data. Data from Lake Geneva were provided by the *Commission Internationale pour la Protection des Eaux du Léman (CIPEL)* and the Information System of the *SOERE OLA, INRA* in Thonon-les-Bains (<http://si-ola.inra.fr>). We also gratefully acknowledge Michela Rogora, *CIPAI*S and the *CNR Water Research Institute* in Verbania, Italy for providing data on Lago Maggiore. We thank Martin Wessels from LUBW, Langenargen, Germany for providing data and logistical support in investigations of Lake Constance. Further, work on Lake Constance was supported by Swiss National Science Foundation grants 200021_146652 and 200020_165517. The authors declare no conflict of interests.

7. References

- Anneville, O., and J. P. Pelletier. 2000. Recovery of Lake Geneva from eutrophication: quantitative response of phytoplankton. *Archiv für Hydrobiologie* **148**: 607-624, <http://dx.doi.org/10.1127/archiv-hydrobiol/148/2000/607>.
- Bechtel, A., and C. J. Schubert. 2009. A biogeochemical study of sediments from the eutrophic Lake Lugano and the oligotrophic Lake Brienz, Switzerland. *Org. Geochem.* **40**: 1100-1114, <http://doi.org/10.1016/j.orggeochem.2009.06.009>.
- Bierlein, K. A., M. Rezvani, S. A. Socolofsky, L. D. Bryant, A. Wüest, and J. C. Little. 2017. Increased sediment oxygen flux in lakes and reservoirs: The impact of hypolimnetic oxygenation. *Water Resour. Res.* **53**: 4876-4890, <http://doi.org/10.1002/2016WR019850>.

- 454 Blais, J. M., and J. Kalff. 1995. The influence of lake morphometry on sediment focusing. *Limnol.*
455 *Oceanogr.* **40**: 582-588, <http://doi.org/10.4319/lo.1995.40.3.0582>.
- 456 Bloesch, J., and U. Uehlinger. 1986. Horizontal sedimentation differences in a eutrophic Swiss
457 lake. *Limnol. Oceanogr.* **31**: 1094-1109, <http://dx.doi.org/10.4319/lo.1986.31.5.1094>.
- 458 Bryant, L. D., D. F. McGinnis, C. Lorrai, A. Brand, J. C. Little, and A. Wüest. 2010. Evaluating
459 oxygen fluxes using microprofiles from both sides of the sediment-water interface. *Limnol.*
460 *Oceanogr. Methods* **8**: 610-627, <http://dx.doi.org/10.4319/lom.2010.8.0610>.
- 461 Burns, N. M. 1995. Using hypolimnetic dissolved oxygen depletion rates for monitoring lakes. *N.*
462 *Z. J. Mar. Freshwat. Res.* **29**: 1-11, <http://dx.doi.org/10.1080/00288330.1995.9516634>.
- 463 Carey, C. C., J. P. Doubek, R. P. McClure, and P. C. Hanson. 2018. Oxygen dynamics control the
464 burial of organic carbon in a eutrophic reservoir. *Limnology and Oceanography Letters*,
465 <http://dx.doi.org/10.1002/lol2.10057>.
- 466 Carignan, R., and D. Lean. 1991. Regeneration of dissolved substances in a seasonally anoxic lake:
467 The relative importance of processes occurring in the water column and in the sediments.
468 *Limnol. Oceanogr.* **36**: 683-707, <http://dx.doi.org/10.4319/lo.1991.36.4.0683>.
- 469 Carstens, D., K. E. Köllner, H. Bürgmann, B. Wehrli, and C. J. Schubert. 2012. Contribution of
470 bacterial cells to lacustrine organic matter based on amino sugars and d-amino acids.
471 *Geochim. Cosmochim. Acta* **89**: 159-172, <https://doi.org/10.1016/j.gca.2012.04.052>.
- 472 Carstens, D., M. F. Lehmann, T. B. Hofstetter, and C. J. Schubert. 2013. Amino acid nitrogen
473 isotopic composition patterns in lacustrine sedimenting matter. *Geochim. Cosmochim.*
474 *Acta* **121**: 328-338, <https://doi.org/10.1016/j.gca.2013.07.020>.

- 475 Charlton, M. N. 1980. Hypolimnion Oxygen Consumption in Lakes: Discussion of Productivity
476 and Morphometry Effects. *Can. J. Fish. Aquat. Sci.* **37**: 1531-1539,
477 <http://doi.org/10.1139/f80-198>.
- 478 Crowe, S. A. and others 2017. Novel anammox bacteria and nitrogen loss from Lake Superior.
479 *Scientific Reports* **7**: 13757, <https://doi.org/10.1038/s41598-017-12270-1>.
- 480 Edwards, W. J., J. D. Conroy, and D. A. Culver. 2005. Hypolimnetic Oxygen Depletion Dynamics
481 in the Central Basin of Lake Erie. *J. Great Lakes Res.* **31**: 262-271,
482 [https://doi.org/10.1016/S0380-1330\(05\)70319-1](https://doi.org/10.1016/S0380-1330(05)70319-1).
- 483 Epping, E. H., and W. Helder. 1997. Oxygen budgets calculated from in situ oxygen microprofiles
484 for Northern Adriatic sediments. *Cont. Shelf Res.* **17**: 1737-1764,
485 [http://dx.doi.org/10.1016/S0278-4343\(97\)00039-3](http://dx.doi.org/10.1016/S0278-4343(97)00039-3).
- 486 Finger, D. and others 2007. Effects of alpine hydropower operations on primary production in a
487 downstream lake. *Aquat. Sci.* **69**: 240-256, <http://doi.org/10.1007/s00027-007-0873-6>.
- 488 Frenzel, P., B. Thebrath, and R. Conrad. 1990. Oxidation of methane in the oxic surface layer of a
489 deep lake sediment (Lake Constance). *FEMS Microbiol. Ecol.* **6**: 149-158,
490 <http://doi.org/10.1111/j.1574-6968.1990.tb03935.x>.
- 491 Grasset, C., R. Mendonça, G. Villamor Saucedo, D. Bastviken, F. Roland, and S. Sobek. 2018.
492 Large but variable methane production in anoxic freshwater sediment upon addition of
493 allochthonous and autochthonous organic matter. *Limnol. Oceanogr.* **63**: 1488-1501,
494 <http://doi.org/10.1002/lno.10786>.
- 495 Gudas, C., D. Bastviken, K. Steger, K. Premke, S. Sobek, and L. J. Tranvik. 2010. Temperature-
496 controlled organic carbon mineralization in lake sediments. *Nature* **466**: 478-481,
497 <http://dx.doi.org/10.1038/nature09186>.

- 498 Holmer, M., and P. Storkholm. 2001. Sulphate reduction and sulphur cycling in lake sediments: a
499 review. *Freshwat. Biol.* **46**: 431-451, <http://doi.org/10.1046/j.1365-2427.2001.00687.x>.
- 500 Holmkvist, L., T. G. Ferdelman, and B. B. Jørgensen. 2011. A cryptic sulfur cycle driven by iron
501 in the methane zone of marine sediment (Aarhus Bay, Denmark). *Geochim. Cosmochim.*
502 *Acta* **75**: 3581-3599, <http://dx.doi.org/10.1016/j.gca.2011.03.033>.
- 503 Kubáň, P., H. T. A. Nguyen, M. Macka, P. R. Haddad, and P. C. Hauser. 2007. New Fully Portable
504 Instrument for the Versatile Determination of Cations and Anions by Capillary
505 Electrophoresis with Contactless Conductivity Detection. *Electroanalysis* **19**: 2059-2065,
506 <http://dx.doi.org/10.1002/elan.200703908>.
- 507 Lehman, J. T. 1975. Reconstructing the rate of accumulation of lake sediment: The effect of
508 sediment focusing. *Quatern. Res.* **5**: 541-550, [http://dx.doi.org/10.1016/0033-](http://dx.doi.org/10.1016/0033-5894(75)90015-0)
509 [5894\(75\)90015-0](http://dx.doi.org/10.1016/0033-5894(75)90015-0).
- 510 Li, J., S. A. Crowe, D. Miklesh, M. Kistner, D. E. Canfield, and S. Katsev. 2012. Carbon
511 mineralization and oxygen dynamics in sediments with deep oxygen penetration, Lake
512 Superior. *Limnol. Oceanogr.* **57**: 1634-1650, <http://dx.doi.org/10.4319/lo.2012.57.6.1634>.
- 513 Li, Y.-H., and S. Gregory. 1974. Diffusion of ions in sea water and in deep-sea sediments.
514 *Geochim. Cosmochim. Acta* **38**: 703-714, [http://doi.org/10.1016/0016-7037\(74\)90145-8](http://doi.org/10.1016/0016-7037(74)90145-8).
- 515 Livingstone, D. M., and D. M. Imboden. 1996. The prediction of hypolimnetic oxygen profiles: a
516 plea for a deductive approach. *Can. J. Fish. Aquat. Sci.* **53**: 924-932,
517 <http://dx.doi.org/10.1139/f95-230>.
- 518 Maerki, M., B. Müller, and B. Wehrli. 2006. Microscale mineralization pathways in surface
519 sediments: A chemical sensor study in Lake Baikal. *Limnol. Oceanogr.* **51**: 1342-1354,
520 <http://dx.doi.org/10.4319/lo.2006.51.3.1342>.

- 521 Maerki, M., B. Wehrli, C. Dinkel, and B. Müller. 2004. The influence of tortuosity on molecular
522 diffusion in freshwater sediments of high porosity. *Geochim. Cosmochim. Acta* **68**: 1519-
523 1528, <http://doi.org/10.1016/j.gca.2003.09.019>.
- 524 Martin, P., L. Granina, K. Martens, and B. Goddeeris. 1998. Oxygen concentration profiles in
525 sediments of two ancient lakes: Lake Baikal (Siberia, Russia) and Lake Malawi (East
526 Africa). *Hydrobiologia* **367**: 163-174, <http://doi.org/10.1023/a:1003280101128>.
- 527 Matthews, D. A., and S. W. Effler. 2006. Long-term changes in the areal hypolimnetic oxygen
528 deficit (AHOD) of Onondaga Lake: Evidence of sediment feedback. *Limnol. Oceanogr.*
529 **51**: 702-714, http://dx.doi.org/10.4319/lo.2006.51.1_part_2.0702.
- 530 Matzinger, A., B. Müller, P. Niederhauser, M. Schmid, and A. Wüest. 2010. Hypolimnetic oxygen
531 consumption by sediment-based reduced substances in former eutrophic lakes. *Limnol.*
532 *Oceanogr.* **55**: 2073-2084, <http://dx.doi.org/10.4319/lo.2010.55.5.2073>.
- 533 Müller, B., L. D. Bryant, A. Matzinger, and A. Wüest. 2012. Hypolimnetic oxygen depletion in
534 eutrophic lakes. *Environ. Sci. Technol.* **46**: 9964-9971,
535 <http://dx.doi.org/10.1021/es301422r>.
- 536 Müller, B. and others 2007. Present and past bio-available phosphorus budget in the ultra-
537 oligotrophic Lake Brienz. *Aquat. Sci.* **69**: 227-239, [https://doi.org/10.1007/s00027-007-](https://doi.org/10.1007/s00027-007-0871-8)
538 [0871-8](https://doi.org/10.1007/s00027-007-0871-8).
- 539 Müller, B. and others 2005. Internal carbon and nutrient cycling in Lake Baikal: sedimentation,
540 upwelling, and early diagenesis. *Global Planet. Change* **46**: 101-124,
541 <http://dx.doi.org/10.1016/j.gloplacha.2004.11.008>.

- 542 Müller, B., T. Steinsberger, R. Schwefel, R. Gächter, M. Sturm, and A. Wüest. 2019. Oxygen
543 consumption in seasonally stratified lakes decreases only below a marginal phosphorus
544 threshold. *Scientific Reports* **9**: 18054, <https://doi.org/10.1038/s41598-019-54486-3>.
- 545 Müller, B., R. Stierli, and A. Wüest. 2006. Phosphate adsorption by mineral weathering particles
546 in oligotrophic waters of high particle content. *Water Resour. Res.* **42**,
547 <https://doi.org/10.1029/2005wr004778>.
- 548 Och, L. M. and others 2012. New insights into the formation and burial of Fe/Mn accumulations
549 in Lake Baikal sediments. *Chem. Geol.* **330–331**: 244-259,
550 <http://dx.doi.org/10.1016/j.chemgeo.2012.09.011>.
- 551 Polerecky, L., N. Volkenborn, and P. Stief. 2006. High temporal resolution oxygen imaging in
552 bioirrigated sediments. *Environ. Sci. Technol.* **40**: 5763-5769,
553 <http://doi.org/10.1021/es060494l>.
- 554 Putyrskaya, V., E. Klemm, and S. Röhlir. 2009. Migration of ¹³⁷Cs in tributaries, lake water and
555 sediment of Lago Maggiore (Italy, Switzerland) – analysis and comparison with Lago di
556 Lugano and other lakes. *J. Environ. Radioact.* **100**: 35-48,
557 <https://doi.org/10.1016/j.jenvrad.2008.10.005>.
- 558 Randlett, M.-E. and others 2015. Mineralization pathways of organic matter deposited in a river–
559 lake transition of the Rhone River Delta, Lake Geneva. *Environmental Science: Processes*
560 *& Impacts* **17**: 370-380, <http://dx.doi.org/10.1039/C4EM00470A>.
- 561 Remsen, C. C., E. C. Minnich, R. S. Stephens, L. Buchholz, and M. E. Lidstrom. 1989. Methane
562 oxidation in Lake Superior sediments. *J. Great Lakes Res.* **15**: 141-146,
563 [http://dx.doi.org/10.1016/S0380-1330\(89\)71468-4](http://dx.doi.org/10.1016/S0380-1330(89)71468-4).

- Schwalb, A., P. Hadorn, N. Thew, and F. Straub. 1998. Evidence for Late Glacial and Holocene environmental changes from subfossil assemblages in sediments of Lake Neuchâtel, Switzerland. *Palaeogeogr., Palaeoclimatol., Palaeoecol.* **140**: 307-323, [https://doi.org/10.1016/S0031-0182\(98\)00025-X](https://doi.org/10.1016/S0031-0182(98)00025-X).
- Schwefel, R., T. Steinsberger, D. Bouffard, L. D. Bryant, B. Müller, and A. Wüest. 2018. Using small-scale measurements to estimate hypolimnetic oxygen depletion in a deep lake. *Limnol. Oceanogr.* **63**: S54-S67, <http://dx.doi.org/10.1002/lno.10723>.
- Small, G. E. and others 2013. Rates and controls of nitrification in a large oligotrophic lake. *Limnol. Oceanogr.* **58**: 276-286, <http://doi.org/10.4319/lo.2013.58.1.0276>.
- Smith, D. A., and G. Matisoff. 2008. Sediment oxygen demand in the central basin of Lake Erie. *J. Great Lakes Res.* **34**: 731-744, [http://dx.doi.org/10.1016/S0380-1330\(08\)71614-9](http://dx.doi.org/10.1016/S0380-1330(08)71614-9).
- Sobek, S. and others 2009. Organic carbon burial efficiency in lake sediments controlled by oxygen exposure time and sediment source. *Limnol. Oceanogr.* **54**: 2243-2254, <http://dx.doi.org/10.4319/lo.2009.54.6.2243>.
- Soomets, T., T. Kutser, A. Wüest, and D. Bouffard. 2019. Spatial and temporal changes of primary production in a deep peri-alpine lake. *Inland Waters* **9**: 49-60, <https://doi.org/10.1080/20442041.2018.1530529>.
- Span, D., D. Arbouille, H. Howa, and J.-P. Vernet. 1990. Variation of nutrient stocks in the superficial sediments of Lake Geneva from 1978 to 1988. *Hydrobiologia* **207**: 161-166, <http://dx.doi.org/10.1007/BF00041453>.
- Steinsberger, T., B. Müller, C. Gerber, B. Shafei, and M. Schmid. 2019. Modeling sediment oxygen demand in a highly productive lake under various trophic scenarios. *PLOS ONE* **14**: e0222318, <https://doi.org/10.1371/journal.pone.0222318>.

- Steinsberger, T., M. Schmid, A. Wüest, R. Schwefel, B. Wehrli, and B. Müller. 2017. Organic carbon mass accumulation rate regulates the flux of reduced substances from the sediments of deep lakes. *Biogeosciences* **14**: 3275-3285, <http://dx.doi.org/10.5194/bg-14-3275-2017>.
- Thevenon, F., S. B. Wirth, M. Fujak, J. Poté, and S. Girardclos. 2013. Human impact on the transport of terrigenous and anthropogenic elements to peri-alpine lakes (Switzerland) over the last decades. *Aquat. Sci.* **75**: 413-424, <https://doi.org/10.1007/s00027-013-0287-6>.
- Thomsen, U., B. Thamdrup, D. A. Stahl, and D. E. Canfield. 2004. Pathways of organic carbon oxidation in a deep lacustrine sediment, Lake Michigan. *Limnol. Oceanogr.* **49**: 2046-2057, <https://doi.org/10.4319/lo.2004.49.6.2046>.
- Torres, N. T., P. C. Hauser, G. Furrer, H. Brandl, and B. Müller. 2013. Sediment porewater extraction and analysis combining filter tube samplers and capillary electrophoresis. *Environmental Science: Processes & Impacts* **15**: 715-720, <http://dx.doi.org/10.1039/c3em00068k>.
- Torres, N. T. and others 2014. Early diagenetic processes generate iron and manganese oxide layers in the sediments of Lake Baikal, Siberia. *Environmental Science: Processes & Impacts* **16**: 879-889, <https://doi.org/10.1039/C3EM00676J>
- Wehrli, B., A. Lotter, T. Schaller, and M. Sturm. 1997. High-resolution varve studies in Baldeggersee (Switzerland): project overview and limnological background data. *Aquat. Sci.* **59**: 285-294, <http://doi.org/10.1007/BF02522359>.
- Wenk, C. B. and others 2013. Anaerobic ammonium oxidation (anammox) bacteria and sulfide-dependent denitrifiers coexist in the water column of a meromictic south-alpine lake. *Limnol. Oceanogr.* **58**: 1-12, <https://doi.org/10.4319/lo.2013.58.1.0001>.

- 609 Wirth, S. B., S. Girardclos, C. Rellstab, and F. S. Anselmetti. 2011. The sedimentary response to a
610 pioneer geo-engineering project: Tracking the Kander River deviation in the sediments of
611 Lake Thun (Switzerland). *Sedimentology* **58**: 1737-1761, [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-3091.2011.01237.x)
612 [3091.2011.01237.x](https://doi.org/10.1111/j.1365-3091.2011.01237.x)
- 613
- 614 Zhu, G. and others 2013. Hotspots of anaerobic ammonium oxidation at land–freshwater interfaces.
615 *Nature Geoscience* **6**: 103, <http://doi.org/10.1038/ngeo1683>.
- 616 Zimmermann, J. 2008. Der Walensee - eine sedimentologische Rekonstruktion seiner holozänen
617 Ereignisgeschichte. Master Thesis University Zurich & EAWAG Dübendorf,
- 618

Table 1. Physiographic and limnological characteristics of the lakes investigated by this study. A_H and V_H represent the hypolimnion area and volume, respectively. z_H is the mean hypolimnion depth (V_H/A_H). The hypolimnion was defined as the water body below 15 m water depth.

Lake	Max. depth	Volume	Surface	A_H	V_H	z_H
	<i>m</i>	<i>km³</i>	<i>km²</i>	<i>km²</i>	<i>km³</i>	<i>m</i>
Maggiore (MAG)	372	37.6	213	196	34.5	176
Geneva (GEN)	310	89.1	582	534	80.8	151
Brienzi (BRI)	260	5.0	29.8	27.8	4.59	165
Constance (CON)	253	47.8	482	403	41.4	103
Thun (THU)	217	6.46	47.9	43.9	5.78	132
Lucerne (LUC)	214	2.30	22	21.1	1.98	93.9
Neuchâtel (NEU)	154	13.8	218	156	11.1	71.3
Walensee (WAL)	151	2.51	24.1	22.8	2.15	94.3
Aegeri (AEG)	81	0.35	7.21	6.4	0.26	46.6
Baldegg (BAL)	66	0.17	5.22	4.53	0.13	27.6
Hallwil (HAL)	48	0.29	9.95	8.58	0.19	22.6

Table 2: Sediment porewater fluxes of reduced and oxidized species expressed as O_2 equivalents in $mmol\ O_2\ m^{-2}\ d^{-1}$. Reduced species are summed as F_{red} and oxidants as SOU . Values of Lake Constance are averages from three sediment cores. Lakes are sorted according to decreasing mean hypolimnion depth.

Lake	J_{CH_4}	$J_{NH_4^+}$	$J_{Fe(II)}$	$J_{Mn(II)}$	F_{red}	J_{O_2}	$J_{NO_3^-}$	$J_{SO_4^{2-}}$	SOU
Maggiore	3.85	1.96	0.04	0.13	5.97	14.0	0.47	1.95	16.4
Geneva	1.74	0.30	0.01	0.03	2.07	4.48	0.86	2.7	8.04
Brienzen	1.40	0	0	0.02	1.42	7.06	0.48	0.33	7.87
Constance	1.31	0.54	0.04	0.11	2.02	6.78	0.53	1.28	8.59
Thun	2.25	0	0	0.05	2.30	8.24	0.32	2.05	10.6
Lucerne	0.57	0	0.03	0.03	0.62	9.08	1.37	0.50	11.0
Neuchâtel	3.06	0	0.01	0.01	3.08	9.52	0.60	0.50	10.6
Walensee	3.32	0	0.07	0.18	3.57	16.2	0.76	0.66	17.6
Aegeri	2.15	1.59	0.11	0.11	3.96	10.6	0.46	0.48	11.6
Baldegg	7.71	5.68	0.25	0.11	13.8	18.8	2.36	2.04	23.2
Hallwil	3.48	1.96	0.13	0.03	5.60	12.4	1.10	1.32	14.8

Table 3: Estimation of the water column mineralization rate (WCM) calculated as the difference between AHM and the sum of SOU and F_{red} . TP is the average concentration after winter mixing over the past 10 years. For Lake Brienzen, the value of 0.6 mg P m⁻³ is calculated as the bio-available TP (Müller et al. 2006).

Lake	F_{red}	SOU	AHM	WCM	TP
$mmol O_2 m^{-2} d^{-1}$				$mg P m^{-3}$	
Maggiore	6.0	16.4	26.3	3.9	11.0
Geneva	2.1	8.0	37.5	27.4	24.3
Brienzen	1.4	7.9	13.1	3.8	0.6
Constance	2.0	8.6	25.0	14.4	7.5
Thun	2.3	10.6	15.0	2.1	1.8
Lucerne	0.6	11.0	20.3	8.7	4.8
Neuchâtel	3.1	10.6	24.7	11.0	9.8

Walensee	3.6	17.6	20.0	~0	3.2
Aegeri	4.0	11.6	19.7	4.1	5.4
Baldegg	13.8	23.2	42.2	5.2	24.1
Hallwil	5.6	14.8	30.9	10.5	15.8

Table 4: Nutrient contents of the bulk sediment. TOC, TN and TOC-MAR values were averaged over 1-10 cm sediment depth. n.d.: not determined. Some data was taken from the respective literature: ¹ Frenzel et al. (1990), ² Span et al. (1990), ³ Schwalb et al. (1998), ⁴ Müller et al. (2007), ⁵ Zimmermann (2008), ⁶ Putyrskaya et al. (2009), ⁷ Sobek et al. (2009), ⁸ Wirth et al. (2011), ⁹ Bierlein et al. (2017),¹⁰ Steinsberger et al. (2017), ¹¹ Schwefel et al. (2018).

Lake	TOC mg C g ⁻¹	TN mg N g ⁻¹	O ₂ penetr. depth mm	Sed. accumul. rate mm yr ⁻¹	TOC-MAR g C m ⁻² yr ⁻¹
Maggiore	14.9±6.2	1.2±0.6	2.6±0.1	3.4 ⁶	33.4
Geneva	11.1 ¹	1.1 ²	4.0±0.5 ¹¹	1.0 ¹⁰	11.1
Brienzen	5.4±1.4	0.6±0.1	6.3±0.8	6.4 ⁴	36.8
Constance	n.d	n.d	2.5 – 4.5 ^{1,7}	n.d	n.d
Thun	11.3±2.0	1.2±0.2	4.8±0.2	6.8 ⁸	67.7
Lucerne	n.d	n.d	4.8±0.6	n.D	n.d
Neuchâtel	22.4±0.8	2.5±0.4	3.6±0.4	2.5 ³	23.9
Walensee	11.3±1.8	1.3±0.1	5.3	5.7 ⁵	42.9
Aegeri	34.9	3.6±0.7	2.5±0.7	1.9 ¹⁰	22.8
Baldegg	34.2±3.1	3.9±0.4	3.4±0.8	3.3 ¹⁰	45.6
Hallwil	34.4±3.5	4.0±0.2	3.0±1.0 ⁹	2.0 ¹⁰	22.5




Figure 1. Flux at the sediment-water-interface (SWI) expressed in O₂ equivalents [mmol O₂ m⁻² d⁻¹] of the main reduced substances, NH₄⁺ and CH₄ (shown negatively scaled as black and red bars), and oxidants, O₂, SO₄²⁻ and NO₃⁻ (positively scaled as light blue, green and purple bars, respectively). Lakes were sorted according to F_{red}. Fluxes of Mn(II) and Fe(II) were negligible and are not shown.




Figure 2. Depletion rates [mmol O₂ m⁻² d⁻¹] of the sediment-based O₂ sinks SOU (blue) and F_{red} (red). SOU and F_{red} decrease with increasing mean hypolimnion depth (see Table 1).




Figure 3: Hypolimnetic O₂ consumption itemized to the three O₂ sinks: F_{red}, SOU and WCM. Lakes are divided into two groups: very deep lakes (above dashed line), lakes biased by intense sediment focusing which causes increased SOU and F_{red} (Lago Maggiore and Walensee) and artificially aerated deep Lakes Baldegg and Hallwil (below dashed line).




Figure 4: Oxygen consumption rates for the water column (WCM) normalized by mean hypolimnion depth (z_H). WCM increases with TP up to a value of around $\sim 10 \text{ mg P m}^{-3}$ and then levels off. Lakes Baldegg and Hallwil are not shown because they are artificially aerated.

Supporting Information

Hypolimnetic oxygen depletion in deep lakes: effects of trophic state and organic matter accumulation

Thomas Steinsberger¹, Robert Schwefel^{2,3}, Alfred Wüest^{1,3}, Beat Müller¹

¹ Eawag, Swiss Federal Institute of Aquatic Science and Technology, CH-6047 Kastanienbaum, Switzerland

² UC Santa Barbara, 3015 Marine Science Building, Santa Barbara, CA 93106-6150, US.

³ Physics of Aquatic Systems Laboratory, Margaretha Kamprad Chair, École Polytechnique Fédérale de Lausanne, Institute of Environmental Engineering, Lausanne, Switzerland.

ORCID: Thomas Steinsberger: 0000-0002-0751-5424; Robert Schwefel: 0000-0003-1610-4181; Beat Müller: 0000-0003-3696-9035; Alfred Wüest: 0000-0001-7984-0368




Figure S1 : Sediment porewater concentration profiles of CH_4 (black), NH_4^+ (purple), Mn(II) (green) and Fe(II) (red) for Lakes : Maggiore (MAG), Geneva (GEN), Brienz (BRI), Constance Site A and B (CON), Thun (THU), Lucerne (LUC), Neuchâtel (NEU), Walensee (WAL), Aegeri (AEG), Baldegg (BAL) and Hallwil (HAL).




Figure S2: Sediment porewater profiles of O_2 (black), SO_4^{2-} (red) and NO_3^- (purple). Note the separate y axes for O_2 (right side) and NO_3^-/SO_4^{2-} (left side). Lake abbreviations : Maggiore (MAG), Geneva (GEN), Brienz (BRI), Constance Site A and B (CON), Thun (THU), Lucerne (LUC), Neuchâtel (NEU), Walensee (WAL), Aegeri (AEG), Baldegg (BAL) and Hallwil (HAL).




Figure S3: Sediment TOC (red) and TN (purple) concentration profiles. Lakes Aegeri (AEG), Baldegg (BAL) and Halwill (HAL) are shown with different TN scale. Lake abbreviations : Maggiore (MAG), Brienz (BRI), Thun (THU), Neuchâtel (NEU), Walensee (WAL), Aegeri (AEG), Baldegg (BAL) and Hallwil (HAL).

Table S1. Additional core sampling info

Lake	Sampling Depth (m)	Nr. of Sampling sites	Coordinates	Number of cores at each site	Sampling date	Number of O ₂ profiles
Maggiore	273	1	698758 106354	4	10.07.2018	2
Brien	259	1	639226 174179	4	23.07.2018	5
Geneva	175	1	535600 140900	4	13.05.2017	5 (Schwefel et al. 2018)
Thun	214	1	622502 170139	4	27.07.2018	4
Constance	81, 144	2	749287 274886 743978 274105	4	01-04.08 2017	5 and data from Sobek et al. (2009)
Walen	151	1	735380 220704	4	04.07.2018	1
Lucerne	150	1	675762 207210	4	24.04.2018	3
Neuchatel	152	1	559310 198615	4	19.09.2018	6
Aegeri	79	1	690601 218797	16	March-October 2014	8
Baldeg	65	1	662219 227762	16	March-October 2014	9
Hallwil	44	1	658862 236313	8	08.04 – 30.07 2015	2 (Bierlein et al. 2017)

Bierlein, K. A., M. Rezvani, S. A. Socolofsky, L. D. Bryant, A. Wüest, and J. C. Little. 2017. Increased sediment oxygen flux in lakes and reservoirs: The impact of hypolimnetic oxygenation. *Water Resour. Res.* **53**: 4876-4890.

Schwefel, R., T. Steinsberger, D. Bouffard, L. D. Bryant, B. Müller, and A. Wüest. 2018. Using small-scale measurements to estimate hypolimnetic oxygen depletion in a deep lake. *Limnol. Oceanogr.* **63**: S54-S67.

Sobek, S. and others 2009. Organic carbon burial efficiency in lake sediments controlled by oxygen exposure time and sediment source. *Limnol. Oceanogr.* **54**: 2243-2254.




Figure 1. Flux at the sediment-water-interface (SWI) expressed in O₂ equivalents [mmol O₂ m⁻² d⁻¹] of the main reduced substances, NH₄⁺ and CH₄ (shown negatively scaled as black and red bars), and oxidants, O₂, SO₄²⁻ and NO₃⁻ (positively scaled as light blue, green and purple bars, respectively). Lakes were sorted according to Fred. Fluxes of Mn(II) and Fe(II) were negligible and are not shown.

187x162mm (600 x 600 DPI)




Figure 2. Depletion rates [mmol O₂ m⁻² d⁻¹] of the sediment-based O₂ sinks SOU (blue) and F_{red} (red). SOU and F_{red} decrease with increasing mean hypolimnion depth (see Table 1).

157x157mm (300 x 300 DPI)




Figure 3: Hypolimnetic O₂ consumption itemized to the three O₂ sinks: F_{red}, SOU and WCM. Lakes are divided into two groups: very deep lakes (above dashed line), lakes biased by intense sediment focusing which causes increased SOU and F_{red} (Lago Maggiore and Walensee) and artificially aerated deep Lakes Baldegg and Hallwil (below dashed line).

188x163mm (300 x 300 DPI)




Figure 4: Oxygen consumption rates for the water column (WCM) normalized by mean hypolimnion depth (z_H). WCM increases with TP up to a value of around ~10 mg P m⁻³ and then levels off. Lakes Baldegg and Hallwil are not shown because they are artificially aerated.

171x119mm (300 x 300 DPI)




Figure S1 : Sediment porewater concentration profiles of CH₄ (black), NH₄⁺ (purple), Mn(II) (green) and Fe(II) (red) for Lakes : Maggiore (MAG), Geneva (GEN), Brienz (BRI), Constance Site A and B (CON), Thun (THU), Lucerne (LUC), Neuchâtel (NEU), Walensee (WAL), Aegeri (AEG), Baldegg (BAL) and Hallwil (HAL).

203x146mm (300 x 300 DPI)




Figure S2: Sediment porewater profiles of O_2 (black), SO_4^{2-} (red) and NO_3^- (purple). Note the separate y axes for O_2 (right side) and NO_3^-/SO_4^{2-} (left side). Lake abbreviations : Maggiore (MAG), Geneva (GEN), Brienz (BRI), Constance Site A and B (CON), Thun (THU), Lucerne (LUC), Neuchâtel (NEU), Walensee (WAL), Aegeri (AEG), Baldegg (BAL) and Hallwil (HAL).

203x134mm (300 x 300 DPI)




Figure S3: Sediment TOC (red) and TN (purple) concentration profiles. Lakes Aegeri (AEG), Baldegg (BAL) and Halwill (HAL) are shown with different TN scale. Lake abbreviations : Maggiore (MAG), Brienz (BRI), Thun (THU), Neuchâtel (NEU), Walensee (WAL), Aegeri (AEG), Baldegg (BAL) and Hallwil (HAL).