



Contents lists available at ScienceDirect

Environmental Pollution

journal homepage: www.elsevier.com/locate/envpol



Erratum

Erratum to “Anaerobic testosterone degradation in *Steroidobacter denitrificans* – Identification of transformation products” [Environ. Pollut. 158 (2010) 2572–2581]

Michael Fahrbach ^a, Martin Krauss ^a, Alfred Preiss ^b, Hans-Peter E. Kohler ^a, Juliane Hollender ^{a,*}

^aEawag, Swiss Federal Institute of Aquatic Science and Technology, Überlandstrasse 133, P.O. Box 611, CH-8600 Dübendorf, Switzerland

^bFraunhofer Institute of Toxicology and Experimental Medicine (ITEM), Nikolai-Fuchs-Strasse 1, D-30625 Hannover, Germany

The publisher regrets that there was an error in the footnote ^b of Table 1 in the above paper. The correct footnote is shown below.

^bDifference between testosterone consumed totally (ΔS) and testosterone assimilated (ΔS_B). The assimilated amount of testosterone was calculated by the equation: $9C_{19}H_{28}O_2 + 50NH_4^+ + 29CO_2 + 24H_2O \rightarrow 50 <C_4H_9O_2N> + 50H^+$ Thus, 1 mg of cell dry mass requires 0.00175 mmol of testosterone. Dissimilated testosterone (ΔS_E) was calculated according to $\Delta S = \Delta S_E + \Delta S_B$.

There was also an error in section 3.1. Complete degradation of testosterone with nitrate as the electron acceptor, FS^T should have been expressed as $AcBE2-1^T$.

DOI of original article: 10.1016/j.envpol.2010.05.017.

* Corresponding author.

E-mail addresses: michael.fahrbach@web.de (M. Fahrbach), martin.krauss@eawag.ch (M. Krauss), alfred.preiss@item.fraunhofer.de (A. Preiss), hkohler@eawag.ch (H.-P.E. Kohler), juliane.hollender@eawag.ch (J. Hollender).