

1

2   **A set of new and cross-amplifying microsatellite loci for**  
3   **conservation genetics of the endangered stone crayfish**  
4   **(*Austropotamobius torrentium*)**

5

6   Christoph Vorburger<sup>1,2</sup>, Nicola Rhyner<sup>2</sup>, Hanna Hartikainen<sup>2,3</sup>, Jukka Jokela<sup>1,2</sup>

7

8   <sup>1</sup>*Institute of Integrative Biology, ETH Zürich, Universitätstrasse 16, 8092 Zürich, Switzerland*

9   <sup>2</sup>*EAWAG, Swiss Federal Institute of Aquatic Science and Technology, Überlandstrasse 133;*

10   *8600 Dübendorf, Switzerland*

11   <sup>3</sup>*Department of Life Sciences, The Natural History Museum, Cromwell Road, SW7 5BD,*

12   *London, United Kingdom*

13

14

15

16

17

18

19

20   **correspondence:**   Christoph Vorburger  
21                           Institute of Integrative Biology, ETH Zürich & EAWAG  
22                           Überlandstrasse 133  
23                           8600 Dübendorf  
24                           Switzerland  
25                           phone: +41 58 765 5196  
26                           fax: +41 58 765 5315  
27                           e-mail: christoph.vorburger@eawag.ch  
28

## Abstract

Like other indigenous crayfish species in Europe, the stone crayfish (*Austropotamobius torrentium*) is endangered due to habitat degradation, competition with invasive American crayfish and a deadly disease (crayfish plague) vectored by these invaders. To study the genetic diversity and connectivity of the remnant stone crayfish populations, eight new microsatellite loci were developed and supplemented by four cross-amplifying loci developed for related species. These twelve loci were tested for polymorphism in 130 stone crayfish from five streams in Switzerland. Variability was low with 1–7 alleles per locus, but the markers revealed strong genetic differentiation among streams. Use of these microsatellites will provide important information for the conservation and management of this endangered species.

**Keywords:** *Austropotamobius torrentium*, Habitat fragmentation, Microsatellites, Stone crayfish

45 Microsatellite sequences were isolated by ecogenics GmbH (Switzerland) using the high-  
46 throughput genomic sequencing approach detailed in Abdelkrim et al. (2009). Genomic DNA  
47 (11 µg) of a single stone crayfish was analyzed on a Roche 454 GS-FLX platform (Roche,  
48 Switzerland) using a 1/16th run and the GS FLX Titanium reagents. The 50'529 reads  
49 obtained had an average length of 257 bp. Microsatellite inserts suitable for primer design  
50 with a tetra- or a trinucleotide motif of at least 6 units or a dinucleotide motif of at least 11  
51 units were present in 137 sequences. Primers were designed for 30 loci and tested in 15  
52 unrelated stone crayfish. Only four primer pairs produced interpretable and polymorphic  
53 amplicons of the expected size. The remaining sequences obtained from ecogenics GmbH  
54 were therefore subjected to another search using the QDD software (Megléczy et al. 2010) and  
55 we designed and tested an additional 23 primer pairs on the same individuals. Of these, again  
56 four produced scorable products of variable size. We also tested 13 primer pairs developed  
57 for the congeneric species *A. pallipes* and *A. italicus* (Pedraza-Lara et al. 2011; Gouin et al.  
58 2000), of which four yielded interpretable amplification products. The 12 usable  
59 microsatellite loci so obtained were then grouped into three multiplex reactions according to  
60 annealing temperatures and product sizes to genotype 130 stone crayfish from five streams in  
61 northeastern Switzerland (Table 1). Three streams were in close proximity of each other (<1  
62 km) but separated by man-made migration barriers, the other two streams were more distant  
63 (20 - 50 km) and located in different drainages. DNA was extracted from clipped crayfish  
64 pleopods using a high salt protocol (Sunnucks and Hales 1996). PCRs were performed in 11  
65 µl final volumes, using the Qiagen multiplex kit and the following thermal profile: 95°C for  
66 15 min, 35 cycles at 94°C for 30 s, 1:30 min annealing at 52°C, 61°C or 60°C (for multiplex  
67 I, II or III, respectively) and 1:30 min at 72°C, and a final extension at 60°C for 30 min. The  
68 forward primer of each pair was labeled with a fluorophore as detailed in Table 1, PCR

products were visualized on an ABI 3130 Genetic Analyzer and alleles were sized using GeneMapper<sup>®</sup> (Applied Biosystems). Polymorphism was relatively low in these samples. Two cross-amplifying loci were not polymorphic, the mean allele number was 2.75 and the mean observed heterozygosity 0.321 (Table 1). No significant deviations from Hardy-Weinberg and linkage equilibria were detected. Genetic differentiation among stone crayfish populations from different drainages was strong and highly significant ( $F_{ST}$  values between 0.171 and 0.364). Even the populations from the three closely adjacent but artificially fragmented streams were significantly differentiated (pairwise  $F_{ST}$  between 0.054 and 0.101, all  $P < 0.005$ ). These markers will thus be useful to reconstruct historical population structure as well as recent fragmentation effects, despite their limited variability in these remnant populations at the edge of the species' distribution. The microsatellites will also be used to inform management of this endangered crayfish and to identify suitable source populations for reintroduction attempts.

## Acknowledgements

We thank ecogenics GmbH for help with marker development and Michael Kugler from the Office for Nature, Hunt and Fisheries of Canton St. Gallen for catching permits and support.

## References

- Abdelkrim J, Robertson BC, Stanton JAL, Gemmell NJ (2009) Fast, cost-effective development of species-specific microsatellite markers by genomic sequencing. *BioTechniques* 46 (3):185-191
- Gouin N, Grandjean F, Souty-Grosset C (2000) Characterization of microsatellite loci in the endangered freshwater crayfish *Austropotamobius pallipes* (Astacidae) and their potential use in other decapods. *Mol Ecol* 9 (5):636-637
- Megl  cz E, Costedoat C, Dubut V, Gilles A, Malausa T, Pech N, Martin J-F (2010) QDD: a user-friendly program to select microsatellite markers and design primers from large sequencing projects. *Bioinformatics* 26 (3):403-404. doi:10.1093/bioinformatics/btp670
- Pedraza-Lara C, Gonzalez EG, Bloor P, Doadrio I (2011) Isolation and characterization of novel polymorphic microsatellite loci for the white-clawed crayfish, *Austropotamobius italicus* (Faxon, 1914). *Mol Ecol Res* 11 (3):586-589. doi:10.1111/j.1755-0998.2011.03004.x
- Sunnucks P, Hales DF (1996) Numerous transposed sequences of mitochondrial cytochrome oxidase I-II in aphids of the genus *Sitobion* (Hemiptera: Aphididae). *Mol Biol Evol* 13 (3):510-524

**Table 1.** Characterization of eight novel microsatellite loci for stone crayfish (*Austropotamobius torrentium*) and four cross-amplifying loci from congeneric species in a sample of 130 individuals from five streams in northeastern Switzerland.  $N_A$ : number of different alleles;  $H_O$ : observed heterozygosity;  $H_E$ : expected heterozygosity.

Locus	GenBank accession no./ Reference	Primer sequence (5' – 3') [fluorophore]	Motif	Multiplex	$N_A$	Allele size range (bp)	$H_O$	$H_E$
Newly developed loci:								
AT1	KJ150252	F: GAGGTCTAAGGCGACGAGG [ATTO550] R: CAAGTAAGGGCCGGGTGAG	(AGG) <sub>8</sub> ...(AGC) <sub>6</sub>	II	3	198-210	0.503	0.506
AT9	KJ150253	F: ACGTGTGTCTGTCTCTATAAATTC [ATTO565] R: ATTAATTACGAGGAGGAACAAGAG	(CTGT) <sub>7</sub>	III	2	204-217	0.210	0.213
AT22	KJ150254	F: GCACCCCGTTTTCTATGCG [YYE] R: AGCAAGGTACACCTCTCGG	(AAT) <sub>8</sub>	II	2	241-244	0.225	0.226
AT23	KJ150255	F: CTGTTTCTTGGGTCTCGGC [ATTO565] R: CTCACCGACACCATAACCG	(TGC) <sub>8</sub>	II	3	154-169	0.422	0.423
AT37	KJ150256	F: ACTATCCGACCGAACGAACC [FAM] R: ACAGAACCGATTCTTGGCAT	(TAACC) <sub>12</sub>	III	7	287-338	0.489	0.558
AT40	KJ150257	F: ATGGGTGTAGACCACGAACG [ATTO550] R: CGTGCTAGAATCATTGATCTTACCTT	(AT) <sub>13</sub>	II	2	113-115	0.273	0.258
AT43	KJ150258	F: TTTCCGAATTTCAATCTGCTT [ATTO565] R: CTGCCTTCTCCCTTAACGTG	(AAAC) <sub>5</sub> ...(AC) <sub>15</sub>	I	2	119-134	0.381	0.371
AT52	KJ150259	F: AACGAAGGTTTCAGAACGCC [YYE] R: CGCTGTTTCTCGGTTCTTGT	(AG) <sub>9</sub>	I	2	104-108	0.254	0.240
Cross-amplifying loci:								
AP1	Gouin et al. (2000)	F: TCTTGGGGATTGGCTAGTTG [YYE] R: CCTGAACATAAAAGGTGCTTTGG		I	2	145-147	0.516	0.448
AP6	Gouin et al. (2000)	F: GCTGTGTGGGATGGAGGT [FAM] R: CACTAGCGTATTCAAGCAACT		II	1	342	0.000	0.000
Aitali3	Pedraza-Lara et al. (2011)	F: ATATCATGCGCGTCATCTCC [ATTO565] R: CACACGAGCACCAGACATTG		II	6	249-308	0.579	0.582
Aitali4	Pedraza-Lara et al. (2011)	F: CGTTGATGGTAGAGGGGAAG [FAM] R: CGTCACCGACCATATAAAGTG		I	1	196	0.000	0.000