

## Clarification

---

In our article “Infection dynamics in coexisting sexual and asexual host populations: support for the Red Queen hypothesis” (*American Naturalist* 184:S22–S30), we failed to mention in the methods section that, in addition to scoring snails as uninfected or *Microphallus*-infected, we also scored infections by all other sterilizing trematodes. We were also not clear that the data in figure 3 were based on the proportions of uninfected females in the sexual population versus the asexual population, where “uninfected” means not infected by any species of sterilizing trematodes. Hence the total number of females in these calculations includes uninfected females, *Microphallus*-infected females, and females infected by other sterilizing trematodes. We felt that infections by all species of sterilizing trematodes should be included in our fitness estimates, but we did not make that clear. We thank Ben Bolker and Sang Woo Park for pointing out the oversight.

DANIELA VERGARA,<sup>1</sup> JUKKA JOKELA,<sup>2</sup> and CURTIS M. LIVELY<sup>1</sup>

1. Department of Biology, Indiana University, 1001 East Third Street, Bloomington, Indiana 47405; 2. EAWAG, Swiss Federal Institute of Aquatic Science and Technology, Überlandstrasse 133, CH-8600 Dübendorf, Switzerland; and ETH Zürich, Swiss Federal Institute of Technology, Institute of Integrative Biology, 8092 Zurich, Switzerland

Submitted August 17, 2017

---