

Adaptive plasticity and genetic divergence in feeding efficiency during parallel adaptive radiation of whitefish (*Coregonus* spp.)

BÄNZ LUNDSGAARD-HANSEN ^{1,2}, BLAKE MATTHEWS ³, PASCAL VONLANTHEN ², ANDREAS TAVERNA ²,
OLE SEEHAUSEN ^{1,2}

¹ Division of Aquatic Ecology & Evolution, Institute of Ecology and Evolution, University of Bern, Baltzerstrasse
6, CH-3012 Bern, Switzerland

² Eawag Swiss Federal Institute of Aquatic Science and Technology, Department of Fish Ecology & Evolution,
Centre of Ecology, Evolution and Biogeochemistry, Seestrasse 79, CH-6047 Kastanienbaum, Switzerland

³ Eawag Swiss Federal Institute of Aquatic Science and Technology, Department of Aquatic Ecology, Centre of
Ecology, Evolution and Biogeochemistry, Seestrasse 79, CH-6047 Kastanienbaum, Switzerland

*Corresponding author: Bänz Lundsgaard-Hansen, Tel.: +41 (0)58 765 22 02 Fax: +41 (0)58 765 21 68, E-mail:
baenz.lundsgaard@eawag.ch

Running title: Divergence in feeding efficiency of whitefish

Abstract

Parallel phenotypic divergence in replicated adaptive radiations could either result from parallel genetic divergence in response to similar divergent selection regimes, or from equivalent phenotypically plastic response to the repeated occurrence of contrasting environments. In postglacial fish replicated divergence in phenotypes along the benthic-limnetic habitat axis is commonly observed. Here we use two benthic-limnetic species pairs of whitefish from two Swiss lakes, raised in a common garden design, with reciprocal food treatments in one species pair, to experimentally measure whether feeding efficiency on benthic prey has a genetic basis or whether it underlies phenotypic plasticity (or both). To do so we offered experimental fish mosquito larvae, partially buried in sand, and measured multiple feeding efficiency variables. Our results reveal both, genetic divergence as well as phenotypically plastic divergence in feeding efficiency, with the phenotypically benthic species raised on benthic food being the most efficient forager on benthic prey. This indicates that both, divergent natural selection on genetically heritable traits and adaptive phenotypic plasticity, are likely important mechanisms driving phenotypic divergence in adaptive radiation.

Key Words: speciation, limnetic, benthic, natural selection, heritable, trophic adaptation, size

Introduction

Parallel adaptive radiations of closely related taxa often exhibit a repeated occurrence of similar ecotypes in similar niches (Schluter, 2000). Such parallel ecotypic differentiation is often attributed to similar evolutionary responses to divergent selection between contrasting environments (Schluter & Nagel, 1995; Schluter, 2000; Barrett & Schluter, 2008), which assumes a genetically heritable basis of the traits characterizing the adaptive radiation. On the other hand, adaptive radiation in general and replicated radiation in particular can be facilitated by phenotypic plasticity (Pfenning et al., 2010). The evolution of similar solutions to the same problems (the repeated evolution of similar phenotypes in different radiations) can be explained by ancestral developmental plasticity (Pfenning et al., 2010). Importantly, the ancestral plasticity hypothesis does not negate the importance of natural selection for the fixation of phenotypic differences, rather it proposes that plasticity explains the origin of those differences (West-Eberhard 2003). Neither are adaptation through divergent evolution and adaptation through phenotypic plasticity mutually exclusive and both could act in concert during the origins of adaptive radiations.

There is growing evidence for fitness trade-offs between differentiated morphs or species inhabiting distinct ecological environments across such a broad range of taxa as plants, snails, insects and fish (Boulding & Van Alstyne, 1993; Schluter, 1995; Schluter, 2000; Via et al., 2000; Rundle, 2002; Nosil, 2004). Such trade-offs suggest that the genetically heritable divergence is a result of divergent natural selection and support the idea that natural selection plays an important role in species formation (Schluter, 2000; Via et al., 2000; Rundle, 2002; Nosil, 2004). But some of these reciprocal transplant experiments were not designed to determine whether genetically heritable or phenotypically plastic divergence in early development caused differential fitness in contrasting environments (Boulding & Van Alstyne, 1993; Schluter, 1995). Indeed, there is considerable empirical support for the importance of phenotypic plasticity in diversification of various taxa: Adaptive radiations such as that of Darwin's finches, cichlid fish, stickleback and Anolis lizards all display

variable levels of phenotypic plasticity in traits characterizing these radiations (Grant, 1986; Day et al., 1994; Losos et al., 2000; Bouton et al., 2002; West-Eberhard, 2003; Wund et al., 2008). Furthermore it has been suggested that phenotypic plasticity increases species richness of a clade, most likely by facilitating adaptive diversification and by reducing the risk of extinction (Pfennig & McGee, 2010).

Northern postglacial fish provide striking examples of adaptive radiations, but the mechanisms of inheritance in these radiations, in particular the relative importance of phenotypic plasticity and genetic predisposition in behavior, are not fully understood. Adaptation to alternative trophic niches has been repeatedly observed in these taxa and has been proposed to be an important driver in their diversification (Skúlason & Smith, 1995; Schluter, 2000). Typically, a split along the benthic (lake bottom) to limnetic (open water) habitat axis is observed, which is accompanied by divergence in morphology and trophic ecology: Limnetic morphs/species are usually planktivorous, rather slender, smaller, with a narrower mouth and longer and more numerous gill rakers, whereas benthic morphs/species are more benthivorous, more deep bodied, larger, with a larger mouth and fewer and shorter gill rakers (Robinson & Wilson 1994; Smith & Skúlason, 1996; Schluter, 2000). Taxa displaying this benthic-limnetic-split can be found e.g. in threespine stickleback, rainbow smelt, brown trout, Arctic charr, *Prosopium* and *Coregonus* (Smith & Skúlason, 1996; Taylor, 1999; Schluter, 2000). A genetic basis for shape divergence and differences in feeding efficiency and swimming behavior has been shown in some of these morphs/species (Robinson, 2000; Adams & Huntingford, 2002; Rogers et al. 2002; Klemetsen et al. 2006). In other cases it has been shown that plasticity can affect morphological divergence (Robinson & Parsons 2002), but plasticity in feeding behavior and efficiency have only rarely been measured (but see Day & McPhail, 1996). Experiments specifically designed to measure phenotypically plastic and genetically heritable components in morphology of benthic vs. limnetic ecotypes, found evidence for the presence of both (Day et al., 1994; Adams & Huntingford, 2004; Proulx & Magnan, 2004). However, to our knowledge no study has

yet measured the effects of plasticity and of genetic divergence on morphology as well as on feeding behavior in one and the same experiment, although this is important to identify the traits that affect feeding efficiency.

Whitefish species complexes might fulfill the four criteria that define an adaptive radiation (Schluter 2000), i.e. common ancestry (Bernatchez & Dodson 1994; Pigeon et al. 1997 ; Ostbye et al. 2005a ; Hudson et al. 2011), fast speciation (Bernatchez et al. 1999; Ostbye et al. 2006, Hudson et al. 2011), phenotype-environment correlation (Harrod et al. 2010) and trait utility (Bernatchez 2004; Kahilainen et al. 2007; 2011); and thus represent a good model system to study mechanisms of diversification in adaptive radiations. Sympatric whitefish morphs/species (we adopt species hereafter) are morphologically most strongly divergent in number of gill rakers and in adult body size (Steinmann, 1950; Svärdson 1979; Lindsey 1981; Vonlanthen et al. 2012)), traits likely involved in foraging, which have also been shown to probably be under divergent selection (Bernatchez, 2004; Ostbye et al., 2005b; Rogers & Bernatchez, 2007). Speciation involves divergence along the benthic-limnetic habitat axis as described above (Bernatchez et al., 1996; Lu & Bernatchez, 1999; Ostbye et al., 2006; Landry et al., 2007), but adaptive radiations with more than two species frequently occurred in European whitefish (Steinmann 1950; Svärdson 1979; Hudson et al., 2007; Siwertsson et al. 2010; Vonlanthen et al. 2012). Phenotype-environment correlations between traits involved in foraging and niche utilization in sympatric whitefish has been well documented and suggests an important role of trophic adaptation in the commonly observed benthic-limnetic split of these fish (Bernatchez et al. 1999; Amundsen et al. 2004; Harrod et al. 2010). Similarly, suggestive evidence for trait utility has been observed in sympatric whitefish, indicating that a higher number of gill rakers likely facilitates feeding on smaller zooplankton (Kahilainen et al. 2007; 2011). However, experimental evidence for divergence in feeding efficiency between whitefish species as well as for trait utility is lacking and remains to be tested.

In the large subalpine lakes of Switzerland, multiple whitefish radiations constituting more than 40 different species originated after the last glacial maximum 15000 years ago (Steinmann, 1950; Hudson et al., 2011; Vonlanthen et al. 2012). This impressive whitefish species diversity consists of at least five different adaptive radiations that evolved in parallel (Hudson et al., 2011). Such young and replicated radiations offer excellent opportunities to test for the importance of driving forces and mechanisms of diversification. Here, we raised two species pair of benthic-limnetic whitefish from two Swiss lakes in a common garden design, with reciprocal food treatment in one species pair, to experimentally measure whether feeding efficiency divergence between them has a genetic basis, if phenotypic plasticity can modify feeding efficiency and how feeding efficiency is affected by variation in phenotypes (fish body size and shape). If variation in feeding efficiency was entirely genetically determined we expected to not find any differentiation between the same species raised on different food, but differentiation between the different species independent of the food they were raised on (Fig. 1a). If, on the other hand, variation in feeding efficiency was entirely the result of phenotypic plasticity we expected to find no differentiation between the different species when raised on the same food but differentiation between the same species raised on different food (Fig. 1b). If feeding efficiency was affected by both, genetic divergence and phenotypic plasticity, we expected to find the strongest difference between the benthic species raised on benthic food and the limnetic species raised on limnetic food, while the other treatments would be expected to be intermediate (Fig. 1c). In scenario 1b and 1c we assumed plasticity to be adaptive such that feeding efficiency on benthic food would be higher for fish raised on benthic food than for fish raised on limnetic food. For all hypothetical scenarios outlined above, we for simplicity further assumed that the strength of plasticity does not differ between species. Based on the above outlined existence of empirical evidence for the importance of both, plasticity and genetic divergence in morphology and feeding efficiency of north temperate fish, we predicted that variation in feeding efficiency would have both, a genetic and an environmentally induced component.

139

140 **Materials & Methods**

141 *Study species*

142 Whitefish from two subalpine lakes, Lake Thun and Lake Lucerne, were used. In each of these lakes
143 at least five different whitefish species have been documented based on phenotypic and genetic data
144 (Svarvar & Müller 1982; Bittner, 2009; Vonlanthen et al. 2012), and these represent two
145 independently evolved radiations (Hudson et al. 2011). We studied two species, a benthic and a
146 limnetic ecotype, from each of the lakes, namely *C. sp.* “Bodenbalchen” and *C. zugensis* from Lake
147 Lucerne and *C. sp.* „Balchen” and *C. albellus* from Lake Thun. We chose to focus on these species,
148 because they phenotypically correspond to the commonly observed benthic-limnetic split of north
149 temperate fish (Schluter 2000). *C. sp.* “Bodenbalchen” and *C. sp.* “Balchen” correspond to the benthic
150 phenotypes, they grow fast and reach maximum sizes of 600 and 450 mm, respectively, their mean
151 gill raker number is 29.4 (22-34) and 30.5 (22-33) and both of them spawn in very shallow water of
152 approximately 2-5 m depth (Steinmann 1950; Vonlanthen et al. 2012). *C. zugensis* and *C. albellus*
153 correspond to the limnetic phenotype, they grow slow and reach a maximum size of 300 mm, their
154 mean gill raker number is 38.8 (34-43) and 38.1 (35-44) respectively and both spawn in 25 m and
155 deeper (Steinmann 1950; Vonlanthen et al. 2012). Both species pairs are genetically clearly
156 differentiated from each other (Vonlanthen et al. 2012). Gut content analysis of Lake Lucerne species
157 showed that *C. zugensis* almost exclusively feeds on zooplankton, while the diet of *C. sp.*
158 “Bodenbalchen” is more benthic (Michel, 1996; Mookerji et al. 1998). Although gut content data for
159 fish from Lake Thun is lacking, evidence for divergence in resource use in between whitefish species
160 with different gill raker numbers is abundant (i. e. Bernatchez et al. 1999; Amundsen et al. 2004;
161 Harrod et al. 2010). This suggests that the studied whitefish species from Lake Thun also differ in
162 resource use in nature.

163 *Breeding and raising of fish*

164 Parental fish were caught in winter 2006, during their spawning time on their respective spawning
165 grounds, to breed experimental fish. The benthic species from the two lakes were caught in
166 approximately 2-5 m depth with gill nets having 38-45 mm mesh sizes. The limnetic species were
167 caught in 30-50 m depth using gill nets of 25-28 mm mesh sizes. By doing target fishing on the
168 extreme ends of whitefish spawning depth gradients and by visual inspection of the catches, we
169 made sure that pure individuals belonging to a particular species and no hybrids were caught,
170 although hybridization has not been uncommon during eutrophication of Swiss lakes (Bittner et al.
171 2010; Vonlanthen et al. 2012). From the catches five females and five males were randomly selected
172 from each species. Eggs and sperm were striped in the lab and eggs of all five females were mixed.
173 The eggs were fertilized simultaneously with sperm from the five males, ideally resulting in 25 half-
174 sib families per species. All fish were fed ad libidum once a day, except on Sundays. All juvenile fish
175 were fed with zooplankton for approximately one year. Zooplankton was collected daily from Lake
176 Lucerne by trawling a plankton-net with a mesh size of 250 µm in a depth of around 8 m. Most
177 common zooplankton taxa were *Daphnia*, Copepods, *Chydorus* and *Bosmina*, which ranged from a
178 size of 250 µm to approximately 5 mm. As soon as fish were large enough to be fed with mosquito
179 larvae (*Chironomus plumosus*), food of all juveniles from Lake Thun was switched to mosquito larvae
180 and the juveniles from Lake Lucerne were subsequently raised in a split family design with reciprocal
181 food treatments. Frozen mosquito larvae were used to simulate a benthic feeding environment and
182 zooplankton was used to simulate a limnetic feeding environment. This resulted in four different
183 treatments for fish from Lake Lucerne: Fish belonging to the benthic species raised on benthic food
184 *BB* and raised on limnetic food *BL*; fish belonging to the limnetic species raised on benthic food *LB*
185 and raised on limnetic food *LL*; and two treatments for Lake Thun, *BB* and *LB*. Each treatment was
186 distributed over two raising aquaria, each with a volume of 120*71*50 cm for fish from Lake Lucerne
187 and of 120*142*50 cm for fish from Lake Thun. A flow through system (~ 2.5 l/min) with lake water

was used. Water temperature during raising varied over the seasons and ranged from 6 to 15 ° C (temperature fluctuations were much less pronounced in experiments, as no experiments were done in winter, see below). Illumination was provided with a Cool White T 8 light tube with 5200 LM and with 12 h day and 12 h night rhythm. Initially each aquarium contained 100 individuals. One raising aquarium of the LB treatment from Lake Thun was lost due to a technical accident. As a consequence, the limnetic species of Lake Thun was raised in one aquarium only. Mortalities in aquaria of Lake Thun fish were: $BB_{AQ1}=0.03$; $BB_{AQ2}=0.07$; $LB_{AQ1}=0.1$. In aquaria of Lake Lucerne fish they were: $BB_{AQ1}=0.06$; $BB_{AQ2}=0.07$; $BL_{AQ1}=0.06$; $BL_{AQ2}=0.2$; $LB_{AQ1}=0.05$; $LB_{AQ2}=0.06$; $LL_{AQ1}=0.02$; $LL_{AQ2}=0.09$. When densities of fish diverged through time between raising aquaria, food provisioning was adjusted by eye. At the end of the raising time, fish from the same treatment, which were raised in different aquaria, were consequently never significantly different in size (t-test: the smallest observed p-value = 0.07 for Lake Lucerne fish of the LL treatment).

Two months before the trials started and for the duration of the trials we switched the food environments in the holding tanks once every week to allow all fish to familiarize with both food-types and avoid food recognition or other short-term learning effects to affect our results. The switching of food was paused from October 2009 to May 2010, as no experiments were done in this time period.

Experimental set-up

Experimental aquaria, each with a size of 55*142*40 cm, were divided lengthwise into two compartments using a Plexiglas wall, resulting in one compartment with a size of 33*142*40 cm and the other compartment with a size of 22*142*40 cm. Water temperature varied between 12 – 15 ° C over the entire experimental phase, and was similar between experimental aquaria at each day. The water flow in the aquaria was paused from the moment the fish was introduced into the tank until the experiment was finished. Illumination was the same as during rearing of the fish. The front

window of the aquaria was covered with a reflecting mirror foil to prevent fish from seeing the observer, to avoid observer-induced behavioral changes. The bottom of the aquaria was covered with a layer of quartz sand. The trial was conducted in the larger front compartment, where one fish was tested at one time. Additionally two fish were put in the smaller back compartment and were left there for the entire duration of the trials to stimulate natural behavior of the single experimental fish in the front compartment (single isolated individuals did not display natural behavior).

Trials were performed from July 2009 to August 2010. Which treatment was tested at which day was randomized for all fish tested in 2009 and for all fish from Lake Thun. Additionally a low number of fish from Lake Lucerne (7 individuals from the LB and 1 from the BL treatment, see Table 1) was tested in 2010 to increase the sample size in these treatments. Despite this, the effect of time was unlikely to bias our findings, as time (in days after the first trial was done) was overall not different between any treatment comparison in any of the lakes. To make sure time did not affect our results, we also included the factor year in generalized linear model analysis to control for potential time effects.

Approximately 48 hours before a trial, the experimental fish was introduced into the experimental aquarium and was not fed until the trial started, to increase its motivation to feed. When an experiment started, two petri dishes filled with quartz sand, each containing 10 partially buried but well visible mosquito larvae, were deposited on the bottom using threads to let them down. As soon as the petri dishes were placed on the bottom, the experimental fish was videotaped until all the 20 mosquito larvae were eaten. Fish that did not start feeding within an hour were removed and were not re-used in this experiment (in total 5, all from the limnetic species from Lake Lucerne: 4 LL and 1 LB). All fish that started feeding ate all the larvae within less than one hour after first feeding. After each trial the fish was removed from the experimental aquaria, was anesthetized, total length and weight were measured and a picture from the left side of the body was taken for shape analysis.

Photos were not available for four fish used in the experiments due to a technical problem with a storage device (see Table 1).

Behavioral measurements

Three variables related to feeding efficiency were measured from the video tapes, time to first feeding, time to food depletion, and the number of unsuccessful attacks. Time to first feeding was the time until a fish started feeding after the petri dishes were placed at the bottom of the experimental aquaria. Time to food depletion was the time a fish needed to eat all twenty larvae, measured from the moment it started feeding. As all fish that started feeding ate all larvae, time to food depletion was equivalent to a feeding rate. The number of unsuccessful attacks was the number of targeted attacks a fish made that did not yield a mosquito larva (because it couldn't grab it/lost it immediately after grabbing it). Because all fish were given the same number of larvae, this measure was equivalent to food capture efficiency. Time to first feeding was related to the ability to detect food, the motivation to feed on it and maybe also searching efficiency, while time to food depletion was related to a combination of searching efficiency, food capture efficiency and handling time and the number of unsuccessful attacks represents food capture efficiency. For all feeding efficiency variables, a lower value indicates a higher efficiency.

Shape measurement

Overall body shape variation was quantified using geometric morphometrics methods (Bookstein, 1991). Fourteen homologous landmarks distributed over the whole fish body, were selected based on standard landmark description and previous analysis of *Coregonus* body shape variation (Zelditch et al., 2004; Vonlanthen et al., 2009). Landmarks were set using the software TPSDIG (Rohlf; 2006). Nonshape variation, such as variation in location and orientation, was removed using Generalized Procrustes superimposition (Rohlf & Slice, 1990). Shape variables (x-y-coordinates of individual

landmarks) for each individual were then generated using the thin-plate-spline equation (Bookstein, 1991).

Size correction was done by regression of each shape variable against fish size to remove variation due to allometry (Loy et al., 1998). Residuals were then used for further analysis. As the allometric relationships differed between lakes but not between treatments within lakes, size correction and further analysis of morphometric data was done separately for the two lakes, but pooled for the treatments within lakes. A Principal Component Analysis was performed to display the major axes of shape variation. All morphometric analyses, including size corrections, were performed as implemented in MORPHOJ v.1.02H (Klingenberg, 2011).

Data analysis

Differentiation in Shape and Growth

To test whether size or shape differed significantly between two treatments a Wilcoxon Rank Sum Test was used, because traits were not always normally distributed (Lehmann, 1975). If four treatments were compared, a Kruskal Wallis ANOVA was used (Kruskal & Wallis, 1952). Additionally fish length was compared between treatments and years in two ANOVAs (one per lake) including treatment and year as explanatory variables (residuals of the ANOVAs were normally distributed indicating that assumptions were met). These statistical tests were performed using R v. 2.13.0 (R Development Core Team 2010).

Differentiation in feeding efficiency

Generalized linear models were used to test for associations of feeding efficiency variables with species identity of a fish (referred to as species), with food environment (referred to as environment), PC1 and PC2 of body shape (referred to as PC1 and PC2, respectively), total length of a fish (referred to as length) and the year the experiment was performed (referred to as year). One

GLM was calculated for each lake and for each response variable. The error distribution with the best structural fit of the data to the model was chosen (Burnham & Anderson, 1998), which was a gaussian distribution for time to food depletion (after a log transformation for Lake Lucerne and a square root transformation for Lake Thun), a negative binomial distribution for time to first feeding and a quasi poisson distribution for number of unsuccessful attacks. For gaussian error distribution we used the identity link function, for the quasi poisson distribution we used the log link function and for negative binomial error distribution we used the logit link function as implemented in R (Bolker et al. 2008; R Development Core Team 2010). The initial model included all potential explanatory variables as well as an interaction of species and environment. A backward elimination model selection approach based on AIC was then used to find the model that best explained the variance in the data (Burnham & Anderson, 1998), while always retaining the main effects (species and environment). If necessary, an AICc instead of an AIC was calculated to correct for low sample sizes ($n < 40$, Burnham & Anderson, 1998). QAIC, which is an approximation to AIC, was calculated, when a quasi poisson error distribution was used, because AIC cannot be calculated when using this error distribution (see Table 2) (Lebreton, 1992; Burnham & Anderson, 1998). We compared models using AIC, Akaike Weights (w_i) and evidence ratios (L ratio) (Burnham & Anderson, 1998). In the final model the relationship between residuals and the fitted values was visually checked to ensure normal residuals and similar variance over the fitted values (Zuur et al., 2009). As differences in AIC between the most likely and the second most likely model were sometimes small (< 2 , Burnham & Anderson, 1998), the results of the second most likely model were also examined, but they are not reported as these models yielded similar results and interpretation.

In multivariate analysis two or more explanatory variables influence each other, when they share information. This can lead to different estimated effects of a variable depending on which co-variables are included. Thus univariate post hoc comparisons of the feeding efficiency measure and the variables retained in the final models were performed, if the most likely model contained more

than one explanatory variable. As Post hoc tests for species, environment and year we calculated Wilcoxon Rank Sum Tests, while Spearman Rank Tests were calculated as post hoc tests for PC2 and length (Lehmann, 1975; Lehmann & D'Abrera, 1998). For post hoc tests for species and environment for Lake Lucerne we applied a random sampling approach, because to compare the two species without confounding the comparison by effects of the raising environment (or vice versa), it was necessary that both species contained the same numbers of fish raised in each of the two environments. Therefore an equal number of fish from all four treatments was needed for these comparisons (N_{ph} ranges from 22 to 26, see Table 1). Equal numbers per treatment were achieved by randomly sub-sampling the number of fish in a particular treatment 1000 times to the same sample size in the treatment with the smallest sample size. Subsequently fish from the same species but different rearing environment were pooled to calculate differences between species (or vice versa). For each pooled random sample, Wilcoxon Rank Sum Tests were performed and test statistics were averaged.

All generalized linear model statistics and post hoc comparisons were performed using R v. 2.13.0 (R Development Core Team, 2010). Analyses based on a negative binomial distribution were performed using the packages MASS in R v. 2.13.0 (Venables & Ripley, 2002). All graphs visualizing the models were created using the package Gplots in R v. 2.13.0.

Results

We compared fish feeding efficiency (time to first feeding, time to food depletion and the number of unsuccessful attacks) and fish morphology (length and shape) between different raising aquaria within treatment. As only one out of 24 comparisons was significant (less than expected by chance) and it was further no more significant after Bonferroni correction (the lowest p-value=0.014; critical

p-value after Bonferroni correction = 0.002), we pooled aquaria of the same treatments for all analyses.

Differentiation in size and shape

Individual fish sizes ranged from 95 mm to 186 mm for Lake Lucerne: The BB fish were largest, the BL fish second largest, the LB were second smallest and the LL fish were smallest (Table 1). These between treatment differences in size were significant in an ANOVA including treatment (n=99; F-ratio=15.9, p<0.001) and year (n=99, F-ratio=0.39, p=0.53) as explanatory variables. In Lake Thun fish sizes ranged from 112 to 187 and there was a trend for increased size of the benthic species (n=34, F-ratio=2.89, p=0.09), while fish from the different years did not differ significantly in size (n=34, F-ratio=0.23, p=0.64). Pairwise post-hoc tests for size differences between the treatments reveal plasticity and heritable differences in size (Supplementary Table 1a). PC1 of shape accounted for 31 % of shape variation in Lake Lucerne and for 38% in Lake Thun. PC2 accounted for 24% of shape variation in Lake Lucerne and 15% in Lake Thun. Other PC scores are not included as they neither differentiated between treatments nor were associated with any measured feeding efficiency variable, and the percentage of explained variance was rather low (< 11 % in Lake Thun, < 8 % in Lake Lucerne). PC1 did not differ between treatments in either of the lakes (not shown). But the four treatments of Lake Lucerne fish did significantly differ in PC2 (Kruskal-Wallis chi-squared=8.7, d.f.=3, p=0.03). Pairwise post hoc tests between treatments indicate that shape divergence mainly arises as a consequence of genetic differences between species and not as a result of phenotypic plasticity (Supplementary Table 1b). PC2 was lower in the benthic species, corresponding to more sub-terminal mouths in this species (see Fig. 4d). In Lake Thun the two species did not differ in PC2 (Wilcoxon Rank Sum Test: n=34, W=135, p=0.76, Supplementary Table 1b).

Differentiation in Feeding Efficiency

i) Lake Lucerne

All three measures of feeding efficiency revealed that the BB fish were most efficient and the LL fish were least efficient in feeding on benthic insect larvae (Fig. 2). The BL and the LB fish were intermediate between fish from the BB and LL treatments in all feeding efficiency variables. Results from Kruskal-Wallis ANOVA show that these between treatment differences were significant for time to first feeding and time to food depletion; while there was a trend for the number of unsuccessful attacks (Supplementary Table 2). Pairwise post-hoc tests for feeding efficiency differences between treatments indicate both, a genetic basis as well as phenotypic plasticity, in feeding efficiency divergence (Supplementary Table 2). All feeding efficiency variables were negatively correlated with fish length (Fig. 2), indicating that larger fish were generally more efficient. However, these correlations were only significant over all four treatments and except from one exception not significant within treatment (Foraging Time in the BB treatment, Supplementary Table 3).

The observation of plasticity and species divergence in feeding efficiency as well as in fish length (Supplementary Table 1 and 2), combined with the observation of effects of fish length on feeding efficiency (Supplementary Table 3), suggest that species and plasticity effects on feeding efficiency can be twofold: We referred to *direct* species /environmental effects on feeding efficiency in subsequent paragraphs, if length is included in a model as a co-variable and the measured species/environmental effect is therefore independent of effects of length on feeding efficiency. Additionally the effects of length on feeding efficiency can be considered as *indirect* plasticity or species effects, because the more benthic a treatment is the larger its fish are and the larger fish are, the more efficient they feed on benthic food.

Using generalized linear modeling, time to food depletion in Lake Lucerne was best explained by a model including species, environment and length (Table 2). The effects of the environment and of species were both significant and there was a trend for an effect of length (Table 3). If we controlled

for the effect of length on time to food depletion, fish raised on benthic food and those from the benthic species were more efficient than fish raised on limnetic food and belonging to the limnetic species (Figure 3a). If we controlled for the effects of species and of the environment on time to food depletion, larger fish depleted food in less time (Figure 4a). Post hoc tests revealed that all of the variables retained in the most likely model (Length, species, environment) were significantly associated with time to food depletion (Supplementary Table 2 and 3). The differences in significance levels between multivariate modeling and univariate post hoc tests arose as a result of shared information between different explanatory variables affecting their significance levels in the GLM.

Time to first feeding was best explained by a model including species, environment, length and year (Table 2), whereas only the effect of the environment was significant and there was a trend for the effect of length (Table 3). If we controlled for the effect of length and year on time to first feeding, fish from the benthic species and raised on benthic food were more efficient than fish from the limnetic species and raised on limnetic food (Fig. 3b). Plasticity effects seemed to be stronger in the limnetic species, although there was no statistical support for this, as the interaction between genetics and environment was not significant. If we controlled for the effects of species, the environment and year, larger fish had a lower time to first feeding than smaller fish (Fig. 4b). Univariate post hoc tests revealed that there was a significant association of time to first feeding with the environment, with species and with length, but not with year (Supplementary Table 2 and 3).

The number of unsuccessful attacks was best explained by a model including species, environment, length, PC2 and year (Table 2), whereas length was the only variable with a significant effect (Table 3). Larger fish displayed fewer unsuccessful attacks, independent of the effects of species, of the environment and other co-variables retained in the most likely model (Fig. 4c). There was a trend for the effect of year, with fish tested in the second year failing less often in grabbing larvae. PC2 was non-significant, but there might be a weak trend. Controlling for species, the environment, length and year, illustrated that fish with a more sub-terminal mouth tended to display less unsuccessful

attacks (Fig. 4d). Species and the environment were non-significant, but the benthic species seemed to be slightly more efficient than the limnetic species, when controlling for the effects of length, PC2 and year (Fig. 3c). Univariate post hoc tests revealed a similar pattern as the GLM and were only significant for length (Supplementary Table 2 and 3; and PC2: $S=11972$, $\rho=0.07$, $p=0.46$).

ii) Lake Thun

In Lake Thun, the benthic species was more efficient than the limnetic species by means of time to food depletion and the number of unsuccessful attacks (Fig. 2). Time to first feeding on the other hand was lower in limnetic than in benthic fish (Fig. 2). Wilcoxon Rank Sum Tests show that species differences in time to food depletion were significant, while other efficiency variables were not significantly different between species (Supplementary Table 2). Feeding efficiency was generally higher for larger fish (Fig. 2), however these correlations were neither significant over both species nor within species (Supplementary Table 3).

Using generalized linear modeling, time to food depletion was best explained by a model including species only (Table 2), where the benthic species depleted the food in significantly shorter time (Table 3, Fig. 2). Time to first feeding was also best explained by a model including species only (Table 2), but in this case the effect of species was non-significant (Table 3, Fig. 2). The number of unsuccessful attacks was best explained by a model including species, length, PC2 and year (Table 2), where PC2 was the only variable with a significant effect (Table 3). Fish having a more sub-terminal mouth failed less often in grabbing larvae independent of their species identity, of their length and of the year they were tested (Fig. 4f). There was a trend for length; as larger fish, independent of their genetic background, their shape (PC2), and the year when they were tested, displayed fewer unsuccessful attacks (Fig. 4e). And there was a trend for year, with fish tested in the second year failing to grab larvae less often. The effect of species was not significant, but the benthic species had fewer failed attacks than the limnetic species (Fig. 2e). However, this difference between the two

species disappeared when we controlled for the effects of length, PC2 and year on the number of unsuccessful attacks (Fig. 3d). Univariate post hoc tests were non-significant for an association of species and length with the number of unsuccessful attacks, while they were significant for PC2 and year (Supplementary Table 2 and 3; PC2: $n=34$, $\rho=-0.4$, $p=0.02$).

Fish tested in the second year generally tended to be slightly more efficient than fish tested in the first year, although the effect of year was never significant in any model (see results above). Size differences between the years cannot explain this pattern, because effects of year remained similar if one controlled for the effects of size on efficiency by including it as a co-variable and fish tested in the second year were not generally larger. We lack a testable explanation for this observation. But independent of the reason, the effect of time was unlikely to bias our findings, as treatments were generally randomly assigned to experimental days and time (in days after the first trial was done) was not different between treatments in neither of the lakes. Further the inclusion of year in the GLM analysis controls for year effects and the effects inferred from the GLMs are thus independent of potentially confounding year effects.

Discussion

Our results show that the sympatric benthic-limnetic species pairs of whitefish differ in their feeding efficiency on benthic food, with the benthic species being more efficient than the limnetic species when raised on the same food in both lakes, suggesting a genetic basis of feeding efficiency divergence. These results are in agreement with field studies reporting that sympatric whitefish species often exhibit differences in resource use along the between benthic-limnetic resource axis (Bernatchez et al., 1999; Amundsen et al., 2004; Harrod et al., 2010) and they add more evidence that adaptation to different trophic niches is likely involved in diversification of north temperate fish. In Lake Lucerne, we further found effects of phenotypic plasticity on feeding efficiency, while we did not quantify plasticity effects in Lake Thun. These findings of a genetic basis and of phenotypic

plasticity in feeding efficiency are consistent with the suggested importance of both divergent natural selection on heritable traits as well as adaptive phenotypic plasticity in the evolutionary diversification of traits related to trophic ecology in whitefish (Wimberger, 1994; Rogers & Bernatchez, 2007), and more generally in the build-up of diversity in adaptive radiation (Schluter, 2000; Pfennig et al., 2010).

Species divergence in growth

Independent of the food the fish were raised on, the benthic species grew bigger than the limnetic species, indicating heritable species divergence in growth. This was found for both lakes albeit it was marginally non-significant in fish from Lake Thun. These findings with faster growth in the benthic species (*C. sp. "Balchen"* and *C. sp. "Bodenbalchen"*) are in the same direction as species divergence in nature (Vonlanthen et al. 2012) and are consistent with previous work reporting a genetic basis in species divergence in growth of various fishes, including many salmonids and whitefish (Hatfield, 1997; Garant et al., 2003; Rogers & Bernatchez, 2007). Among fish from Lake Lucerne, we also observed effects of the rearing environment on growth. Fish raised on benthic food generally grew larger than fish raised on limnetic food. This may be explained by differences in energetic profitabilities between our food treatments (zooplankton vs. mosquito larvae), which were not standardized to equal energetic content.

The observed heritable species divergence in growth might have accumulated as a result of divergent selection favoring different growth patterns in the benthic and the limnetic habitat. Slower growth in the limnetic habitat is probably associated with high bioenergetic costs of living in this habitat, with small, spatially widely distributed prey (Mookerji et al. 1998; Trudel et al., 2001; Kahilainen et al. 2007). The benthic habitat with larger and more spatially clustered prey requires less swimming effort and attacks, what allows faster growth (Kahilainen et al. 2003). In this experiment we showed

that increased size is associated with increased feeding efficiency on benthic food and might constitute an adaptation to exploit benthic resources. It might additionally constitute a different predator escape strategy, namely through accelerated growth to reach a size above the predation window of piscivore fish instead of adaptations in predator avoidance through swimming behavior (Kahilainen and Lehtonen 2002; Roger et al., 2002). Because the studied species are young, having emerged after the last glacial maximum (Hudson et al., 2011), our findings of heritable growth divergence between species are consistent with a role of divergent selection on growth early in the speciation process, as it has been shown for other whitefish systems (Rogers and Bernatchez 2007). Taken together evidence for divergent selection on growth and the predominant role of size as a mate-choice signal in fish (Foote and Larkin 1988; Sigurjonsdottir and Gunnarsson 1989; McKinnon et al. 2004), indicates that size might potentially be a magic trait of speciation in whitefish (magic-trait model of speciation: Gavrillets 2004).

In Lake Lucerne, our results further show weak but significant species divergence in shape: The benthic species has a more sub-terminal mouth. In Lake Thun, the two species were not significantly divergent in the shape components we measured. The measured shape components were non-labile in respect to our divergent raising environments for Lake Lucerne fish, indicated by the lack of plasticity effects on shape. Many studies have reported critical effects of the timing of environmental induction on the strength of the plastic response to it (West-Eberhard, 2003). In our experiment all fish had to be raised on zooplankton in the first year (whitefish larvae cannot effectively be raised on benthic food), which could explain why we did not find strong plasticity in morphology induced by divergent feeding regimes while other authors, studying other fish taxa, did find such effects (Day & McPhail, 1996; Bouton et al., 2002; Robinson & Parson, 2002; Muschick et al., 2011). Alternatively it could reflect real differences in canalization of morphology between whitefish and other fish species, however, other studies reported strong plasticity in whitefish morphology (Lindsey 1981).

Evidence for inherited species differences and phenotypic plasticity in feeding efficiency

We found that both benthic species were generally more efficient in foraging on benthic food than their limnetic sister species, suggesting heritable divergence in feeding efficiency. A genetic component of feeding efficiency between benthic-limnetic sister species is consistent with previous experiments using north temperate fish (Robinson, 2000; Adams & Huntingford, 2002). GLM analyses indicate that the effects of this feeding efficiency divergence between species are twofold. On one hand they are manifested as direct behavioral effects, independent of morphological differences (fish length and shape) between species. On the other hand they can be manifested as indirect effects due to inherited differences in length and shape, which themselves influences feeding efficiency. In our experiment fish of the benthic species from both lakes grew larger, and larger fish were generally more efficient foragers on benthic food, independent of their genetic background. This observation of increased efficiency with increasing size is consistent with empirical observation that the more benthic species are usually larger (Schluter 2000; Vonlanthen et al. 2012) and it is not inconsistent with ontogenetic diet shifts to more benthic prey with increasing size in whitefish (Sandlund et al. 1992; Pothoven & Nalepa 2006). Further, in Lake Lucerne we found the benthic species to have a more sub-terminal mouth than the limnetic species and individuals with a sub-terminal mouth displayed fewer failed attacks in our experiments than those with a more terminal mouth. This is consistent predictions from functional morphology and with the empirical observation on many fish taxa, including whitefish, that the position of the mouth relative to the body is associated with benthic vs. limnetic feeding, with benthic feeders having a more sub-terminal mouth (Steinmann, 1950; McCart, 1970; Caldecutt & Adams, 1998; Bernatchez 1999; Clabaut et al., 2007; Harrod et al. 2010).

The observed heritable divergence in feeding behavior between the benthic and limnetic whitefish species is consistent with a role for divergent natural selection favoring different trophic strategies in contrasting foraging environments in north temperate fish (Schluter 1995; Rogers et al., 2002; Klemetsen et al. 2006). Speciation in Lake Thun and Lake Lucerne whitefish was proposed to be intra-

lacustrine (Hudson et al., 2011), therefore ecological character displacement after allopatric speciation and secondary contact seems very unlikely. It remains uncertain whether species divergence in feeding efficiency was a driving force of speciation at the very beginning of the process, or whether it could have occurred as a by-product after speciation was initialized (speciation could have been initialized e.g. by physiological adaptation to different thermal regimes in the contrasting environments and divergence in feeding efficiency would have accumulated afterwards). The species differences in feeding efficiency could also have evolved through genetic assimilation of initially plastic differences in feeding efficiency between whitefish growing up in different habitats (West-Eberhard, 2003).

Our results of species differences in exploiting benthic resources can be considered as evidence for trait utility (Schluter 2000). Trait utility means that a trait associated with a particular environment enhances performance there (Schluter 2000). This feature of adaptive radiations has so far not experimentally been demonstrated for whitefish (Bernatchez 2004), though indirect evidence from comparative approaches suggest that a high number of gill rakers increases fitness in the limnetic environment (Kahilainen et al. 2007; 2011). We showed that the overall phenotype (including behavior), which can be seen as a multi-dimensional trait, of the benthic species increases its ability to exploit the benthic environment. Our results also show that increased size increases feeding efficiency on benthic prey independent of a fish's genetic background. And this is consistent with trait utility of size in regard to fitness in the benthic niche. However, it remains to be tested whether increased size would also increase feeding efficiency on limnetic prey, before we want to draw strong conclusions about trait utility of size in the benthic niche. Other potential traits are, among others, the number of gill rakers (which was not quantified here, as fish were kept alive), the position of the mouth as well as behavioral traits (for example swimming behavior).

Consistent with earlier work (Day and McPhail, 1996), we also observed significant effects of environmental plasticity on feeding efficiency. Fish raised on benthic food became more efficient

foragers on benthic food than fish raised on zooplankton, indicating that the observed plasticity in feeding efficiency was adaptive. This is consistent with the suggested importance of phenotypic plasticity for species diversification in adaptive radiation (West-Eberhard, 1989, 2003; Pfennig et al. 2010). We have two lines of evidence that the effects of plasticity are not simply due to plastic components of size and shape, but primarily the result of plasticity in feeding behavior itself. First, fish size (length) was included in general linear models with significant environmental effects, indicating that plasticity effects are not just due to plasticity in length. Second, there was no plasticity in shape. Earlier work on sticklebacks suggested that behavioral plasticity mainly influenced searching efficiency (Day & McPhail 1996). Consistent with this, the two efficiency variables that showed plasticity in our experiments, time to food depletion and time to first feeding, are more related to detection ability and searching efficiency; whereas the number of unsuccessful attacks, which did not reveal plasticity, is more related to prey capture efficiency. Phenotypic plasticity was suggested to explain why some taxa are more diverse than others, with plasticity increasing species diversity (Pfennig & McGee, 2010). Whitefish and Arctic charr are of the most diverse taxa within the order of the Salmoniformes (Kottelat & Freyhof 2007). Maybe their ability to display strong phenotypic plasticity in feeding behavior and morphology might be one explanation for their high species diversity.

Conclusions

Natural selection is thought to be the most important mechanism behind the diversification of species in adaptive radiations (Schluter, 2000). Our findings of heritable feeding efficiency differences between whitefish species of two parallel adaptive radiations are consistent with this. Additionally, our observation of strong phenotypic plasticity in feeding efficiency indicates an important role of adaptive phenotypic plasticity in diversification of north temperate fish. In conclusion, our data suggest that both, phenotypic plasticity and evolutionary divergence resulting from divergent natural selection, are likely important mechanisms of adaptive radiation.

576 **Acknowledgments**

577 We would like to thank Erwin Schäffer, Lucie Greuter, Jakob Brodersen, Kay Lucek, Alan Hudson,
578 Rudolf Müller, Oliver Selz and all other members of the Fish Ecology and Evolution lab for assistance
579 and valuable comments and suggestions on the manuscript. We also thank the veterinary office of
580 the canton of Lucerne for authorization to conduct this experiment (License number: 04/07). We
581 acknowledge financial support by the Eawag Action Field Grant AquaDiverse – understanding and
582 predicting changes in aquatic biodiversity.

583 **References**

- 584 Adams, C. E. & Huntingford, F. A. 2002. The functional significance of inherited differences in feeding
585 morphology in a sympatric polymorphic population of Arctic charr. *Evol. Ecol.* 16: 15-25.
- 586 Adams, C. E. & Huntingford, F. A. 2004. Incipient speciation driven by phenotypic plasticity? Evidence
587 from sympatric populations of Arctic charr. *Biol. J. Linn. Soc.* 81: 611-618.
- 588 Amundsen, P. A., Knudsen, R., Klemetsen, A. & Kristoffersen, R. 2004. Resource competition and
589 interactive segregation between sympatric whitefish morphs. *Ann. Zool. Fenn.* 41: 301-307.
- 590 Barrett, R. D. H. & Schluter, D. 2008. Adaptation from standing genetic variation. *Trends Ecol. Evol.*
591 23: 38-44.
- 592 Bernatchez, L. 2004. Ecological Theory of Adaptive Radiation: An Empirical Assessment from
593 Coregonine Fishes (Salmoniformes). In: *Evolution Illuminated* (Hendry, A. P. & Stearns, S. C., eds.). pp.
594 175-207. Oxford University Press, Oxford.
- 595 Bernatchez, L., Chouinard, A. & Lu, G. Q. 1999. Integrating molecular genetics and ecology in studies
596 of adaptive radiation: whitefish, *Coregonus sp.*, as a case study. *Biol. J. Linn. Soc.* 68: 173-194.
- 597 Bernatchez, L., & Dodson, J. J. 1994. Phylogenetic relationships among palearctic and nearctic
598 whitefish (*Coregonus sp.*) populations as revealed by mitochondrial DNA variation. *Can. J. Fish.*
599 *Aquat. Sci.* 51: 240-251.
- 600 Bernatchez, L., Vuorinen, J. A., Bodaly, R. A & Dodson, J. J. 1996. Genetic evidence for reproductive
601 isolation and multiple origins of sympatric trophic ecotypes of whitefish (*Coregonus*). *Evolution* 50:
602 624-635.
- 603 Bittner, D. 2009. Gonad deformations in whitefish (*Coregonus spp.*) from Lake Thun, Switzerland - A
604 population genetic and transcriptomic approach. In: *CMPG, Vol. PhD.* pp. University of Bern, Bern.

605 Bittner, D., Excoffier, L. & Largiadèr, C. R. 2010. Patterns of morphological changes and hybridization
 606 between sympatric whitefish morphs (*Coregonus spp.*) in a Swiss lake: a role for eutrophication? Mol.
 607 Ecol. 19: 2152-2167.

608 Bolker, B. M., Brooks, M. E., Clark, C. J., Geange, S. W., Poulsen, J. R., Stevens, M. H. H., White, J.-S. S.
 609 2008. Generalized linear mixed models: a practical guide for ecology and evolution. Trends Ecol. Evol.
 610 24: 127-135.

611 Bookstein, F.L. 1991. Morphometric Tools for Landmark Data, Geometry and Biology. Cambridge
 612 University Press, Cambridge.

613 Boulding, E. G. & Van Alstyne, K. L. 1993. Mechanisms of differential survival and growth of two
 614 species of *Littorina* on wave-exposed and on protected shores. J. Exp. Mar. Biol. Ecol. 169: 139-166.

615 Bouton, N., Witte, F. & Van Alphen, J.J.M. 2002. Experimental evidence for adaptive phenotypic
 616 plasticity in a rock-dwelling cichlid fish from Lake Victoria. Biol. J. Linn. Soc. 77: 185-192.

617 Burnham, K. P. & Anderson, D. R. 1998. Model selection and multimodel inference, a practical
 618 information-theoretic approach. Second edition. Springer, New York.

619 Caldecutt, W. J. & Adams, D. C. 1998. Morphometrics of trophic osteology in the threespine
 620 stickleback, *Gasterosteus aculeatus*. Copeia 4: 827-838.

621 Clabaut, C., Bunje, P. M. E., Salzburger, W. & Meyer, A. 2007. Geometric morphometric analyses
 622 provide evidence for the adaptive character of the tanganyikan cichlid fish radiations. Evolution 61:
 623 560-518.

624 Day, T. & McPhail, J. D. 1996. The effect of behavioural and morphological plasticity on foraging
 625 efficiency in the threespine stickleback (*Gasterosteus sp.*). Oecologia 108: 380–388.

626 Day, T., Pritchard, J. & Schluter, D. 1994. A comparison of two stickleback. Evolution 48: 1723-1734.

627 Foote, C.J., & Larkin P. A. 1988. The role of male choice in the assortative mating of anadromous and
628 non-anadromous sockeye salmon, *Oncorhynchus nerka*. Behaviour 106: 43-62.

629 Garant, D., Dodson, J.J. & Bernatchez, L. 2003. Differential reproductive success and heritability of
630 alternative reproductive tactics in wild Atlantic Salmon (*Salmo salar* L.). Evolution 57: 1133–1141.

631 Gavrillets, S. 2004. Fitness Landscapes and the Origin of Species. Princeton Univ. Press, Princeton.

632 Grant, P. R. 1986. Ecology and evolution of Darwin's finches. Princeton University Press, Princeton.

633 Harrod, C., Mallela, J. & Kahilainen, K. 2010. Phenotype-environment correlations in a putative
634 whitefish adaptive radiation. J. Anim. Ecol. 79: 1057-1068.

635 Hatfield, T. 1997. Genetic divergence in adaptive characters between sympatric species of
636 stickleback. Am. Nat. 149: 1009-1029.

637 Hudson, A. G., Vonlanthen, P. & Seehausen, O. 2011. Rapid parallel adaptive radiations from a single
638 hybridogenic ancestral population. P. R. Soc. B 278: 58-66.

639 Hudson, A. G., Vonlanthen, P., Müller, R., & Seehausen, O. 2007. Review: The geography of
640 speciation and adaptive radiation in coregonines. Adv. Limnol. 60: 111-146.

641 Kahilainen, K. K., & Lehtonen, H. 2002. Brown trout (*Salmo trutta* (L)) and Arctic charr (*Salvelinus*
642 *alpinus* (L)) as predators of three sympatric whitefish (*Coregonus lavaretus* (L)) in the subarctic Lake
643 Muddusjärvi. Ecol. Freshw. Fish 11: 158-167.

644 Kahilainen, K. K., Lehtonen, H., & Könönen, K. 2003. Consequences of habitat segregation to growth
645 rate of two sparsely rakered whitefish (*Coregonus lavaretus* (L.)) in a subarctic lake. Ecol. Freshw. Fish
646 12: 275-285.

647 Kahilainen, K. K., Malinen, T., Tuomaala, A., Alajärvi, E., Tolonen, A., & Lehtonen, A. 2007. Empirical
648 evaluation of phenotype-environment correlation and trait utility with allopatric and sympatric
649 whitefish (*Coregonus lavaretus* (L.)) in subarctic lakes. Biol. J. Linn. Soc. 92: 561-572.

650 Kahilainen, K. K., Siwertsson, A., Gjelland, K. O., Knudsen, R., Bohn, T., & Amundsen, P.-A. 2011. The
651 role of gill raker variability in adaptive radiation of coregonid fish. Evol. Ecol. 25: 573-588.

652 Klemetsen, A., Knudsen, R., Primicerio, R. & Amundsen, P. A. 2006. Divergent, genetically based
653 feeding behaviour of two sympatric Arctic charr, *Salvelinus alpinus* (L.), morphs. Ecol. Freshw. Fish 15:
654 350-355.

655 Klingenberg, C. P. 2011. MORPHOJ: an integrated software package for geometric morphometrics.
656 Mol. Ecol. Resour. 11: 353-357.

657 Kottelat, M. & Freyhof, J. 2007. Handbook of European freshwater fishes. Kottelat, Cornol,
658 Switzerland and Freyhof, Berlin, Germany.

659 Kruskal, W. & Wallis, W. A. 1952. Use of ranks in one-criterion variance analysis. J. Am. Stat. Assoc.
660 47: 583–621.

661 Landry, L., Vincent, W. F. & Bernatchez, L. 2007. Parallelism between limnological features and
662 phenotypic evolution of lake whitefish dwarf ecotypes. J. Evolution Biol. 20: 971-984.

663 Lindsey, C. C. 1981. Stocks are chameleons: Plasticity in gill rakers of coregonid fishes. Can. J. Fish.
664 Aquat. Sci. 38: 1497-1506.

665 Lebreton, J.D., Burnham, K. P., Clobert, J. & Anderson, D. R. 1992. Modeling survival and testing
666 biological hypotheses using marked animals: a unified approach with case studies. Ecol. Monogr. 62:
667 67-118.

668 Lehmann, E. L. 1975. Nonparametric statistical methods based on ranks. McGraw-Hill, New York.

669 Lehmann, E. L. & D'Abrera, H. J. M. 2006. Nonparametrics: Statistical methods based on ranks.
 670 Springer, New York.

671 Losos, J. B., Douglas, A. C., Glossip, D., Goellner, R., Hampton, A., Roberts, G. et al. 2000. Evolutionary
 672 implications of phenotypic plasticity in the hindlimb of the lizard *Anolis sagrei*. *Evolution* 54: 301-305.

673 Loy, A., Mariani, L., Bertelletti, M. & Tunesi, L. 1998. Visualizing allometry: geometric morphometrics
 674 in the study of shape changes in the early stages of the two-banded sea bream, *Diplodus vulgaris*
 675 (Perciformes, Sparidae). *J. Morphol.* 237: 137-146.

676 Lu, G. & Bernatchez, L. 1999. Correlated trophic specialization and genetic divergence in sympatric
 677 lake whitefish ecotypes (*Coregonus clupeaformis*): support for the ecological speciation hypothesis.
 678 *Evolution* 53: 1491-1505.

679 McCart, P. 1970. Evidence for the existence of sibling species of pygmy whitefish (*Prosopium*
 680 *coulteri*) in three Alaskan lakes. In *Biology of Coregonid Fishes*, ed. CC Lindsey, CS Woods, pp. 81-98.
 681 Univ. Manitoba Press, Winnipeg.

682 McKinnon J. S., Mori, S., Blackman, B. K., David, L., Kingsley, D. M. , Jamieson, L., Chou, J., & Schluter,
 683 D. 2004. Evidence for ecology's role in speciation. *Nature* 429: 294-298.

684 Michel, M. 1996. Untersuchungen zur Nahrungsökologie von Grossfelchen im Vierwaldstättersee
 685 während des Sommerhalbjahres 1996. Master thesis EAWAG, supervised by R. Müller.

686 Muschick, M., Barluenga, M., Salzburger, W. & Meyer, A. 2011. Adaptive phenotypic plasticity in the
 687 Midas cichlid fish pharyngeal jaw and its relevance in adaptive radiation. *BMC Evol. Biol.* 11: 116.

688 Mookerji, N., Heller, C., Meng, H. J., Bürgi, R., & Müller, R. 1998. Diel and seasonal patterns of food
 689 uptake and prey selection by *Coregonus sp.* in re-oligotrophicated Lake Lucerne, Switzerland. *J. Fish.*
 690 *Biol.* 52: 443-457.

691 Nosil, P. 2004. Reproductive isolation caused by visual predation on migrants between divergent
692 environments. P. Roy. Soc. Lond. B Bio. 271: 1521-1528.

693 Ostbye, K., Amundsen, P. A., Bernatchez, L., Klemetsen, A., Knudsen, R., Kristoffersen, R. *et al.* 2006.
694 Parallel evolution of ecomorphological traits in the European whitefish *Coregonus lavaretus* (L.)
695 species complex during postglacial times. Mol. Ecol. 15: 3983-4001.

696 Ostbye, K., Bernatchez L., Naesje, T. F., Himberg, M., & Hindar, K. 2005a. Evolutionary history of
697 European whitefish (*Coregonus lavaretus*) as inferred from mtDNA phylogeography and gill-raker
698 numbers. Mol. Ecol. 14: 4371-4388.

699 Ostbye, K., Naesje, T. F., Bernatchez, L., Sandlund, O. T., & Hindar, K. 2005b. Morphological
700 divergence and origin of sympatric populations of European whitefish (*Coregonus lavaretus* (L) in
701 Lake Femud, Norway. J. Evol. Biol. 18: 683-702.

702 Pfennig, D. W. & McGee, M. 2010. Resource polyphenism increases species richness: a test of the
703 hypothesis. Philos. T. R. Soc. B 365: 577-591.

704 Pfennig, D. W., Wund, M. A., Snell-Rood, E. C., Cruickshank, T., Schlichting, C. D. & Moczek, A. P.
705 2010. Phenotypic plasticity's impacts on diversification and speciation. Trends Ecol. Evol. 25: 459–
706 467.

707 Pigeon, D., Chouinard, A., & Bernatchez, L. 1997. Multiple modes of speciation involved in the
708 parallel evolution of sympatric morphotypes of lake whitefish (*Coregonus clupeaformis*). Evolution
709 51: 196-205.

710 Pothoven, S. A., & Nalepa, T. F. 2006. Feeding ecology of lake whitefish in Lake Huron. J. Great Lakes
711 Res. 32: 489-501.

712 Proulx, R. & Magnan, P. 2004. Contribution of phenotypic plasticity and heredity to the trophic
713 polymorphism of lacustrine brook charr (*Salvelinus fontinalis* M.). Evol. Ecol. Res. 6: 503-522.

714 R Development Core Team. 2010. R: A language and environment for Statistical Computing. Vienna,
 715 Austria. <http://www.R-project.org>.

716 Robinson, B. W. 2000. Trade offs in habitat-specific foraging efficiency and the nascent adaptive
 717 divergence of sticklebacks in lakes. Behavior 137: 865-888.

718 Robinson, B.W. & Wilson, D.S. 1994. Character release and displacement in fishes: A neglected
 719 literature. Am. Nat. 144: 596-627.

720 Robinson, B.W. & Parsons, K. J. 2002. Changing times, spaces and faces: tests and implications of
 721 adaptive morphological plasticity in the fishes of northern postglacial lakes. Can. J. Fish. Aquat. Sci.
 722 59: 1819-1833.

723 Rogers, S. M. & Bernatchez, L. 2005. Integrating QTL mapping and genome scans towards
 724 characterization of candidate loci of parallel selection in the lake whitefish (*Coregonus clupeaformis*).
 725 Mol. Ecol. 14: 351-361.

726 Rogers, S. M. & Bernatchez, L. 2007. The genetic architecture of ecological speciation and the
 727 association with signatures of selection in natural lake whitefish (*Coregonus sp.*, Salmonidae) species
 728 pairs. Mol. Biol. Evol. 24: 1423-1438.

729 Rogers, S. M., Gagnon, V. & Bernatchez, L. 2002. Genetically based phenotype-environment
 730 association for swimming behavior in lake whitefish ecotypes (*Coregonus Clupeaformis* Mitchill).
 731 Evolution 56: 2322-2329.

732 Rohlf, F. J. 2006. TPSDig Version 2.1. State University of New York at Stony Brook, Stony Brook, NY,
 733 USA.

734 Rohlf, F. J. & Slice, D. 1990. Extensions of the Procrustes method for the optimal superimposition of
 735 landmarks. Syst. Zool., 39:40-59.

- 736 Rundle, H. D. 2002. A test of ecologically dependent postmating isolation between sympatric
737 sticklebacks. *Evolution* 56: 322–329.
- 738 Sandlund, O. T., Naesje, T. F., & Jonson, B. 1992. Ontogenetic changes in habitat use by whitefish,
739 *Coregonus lavaretus*. *Environ. Biol. Fish.* 33: 341-349.
- 740 Schluter, D. 1995. Adaptive radiation in sticklebacks: trade-offs in feeding performance and growth.
741 *Ecology* 76: 82–90.
- 742 Schluter, D. 2000. *The Ecology of Adaptive Radiation*. Oxford University Press, Oxford.
- 743 Schluter, D. & Nagel, L. 1995. Parallel speciation by natural selection. *Am. Nat.* 146: 292–301.
- 744 Sigurjonsdottir H., & Gunnarsson, K. 1989. Alternative mating tactics of Arctic charr, *Salvelinus*
745 *alpinus*, in Thingvallavatn, Iceland. *Environ. Biol. Fish.* 26: 159-176.
- 746 Siwertsson, A., Knudsen, R., Kahilainen, K., Praebel, K., Primicerio, & R. Amundsen, P. A. 2010.
747 Sympatric diversification as influenced by ecological opportunity and historical contingency in a
748 young species lineage of whitefish. *Evol. Ecol. Res.* 12: 929-948.
- 749 Skúlason, S. & Smith, T. B. 1995. Resource polymorphism in vertebrates. *Trends Ecol. Evol.* 10: 366-
750 370.
- 751 Smith, T. B. & Skúlason, S. 1996. Evolutionary significance of resource polymorphisms in fishes,
752 amphibians, and birds. *Annu. Rev. Ecol. Syst.* 27: 111–133.
- 753 Steinmann, P. 1950. Monographie der schweizerischen Koregonen. Beitrag zum Problem der
754 Entstehung neuer Arten. Spezieller Teil. *Schweiz. Z. Hydrol.* 12: 340-491.
- 755 Svärdson, G. 1979. Speciation of Scandinavian *Coregonus*. *Rep. Inst. Freshw. Res. Drott.* 57: 1-95.
- 756 Svarvar, P. O. & Müller, R. 1982. Die Felchen des Alpnersees. *Schweiz. Z. Hydrol.* 44: 295-314.

757 Taylor, E. B. 1999. Species pairs of north temperate freshwater fishes: evolution, taxonomy, and
 758 conservation. *Rev. Fish Biol. Fisher.* 9: 299-324.

759 Trudel, M., Tremblay, A., Schetagne, R., & Rasmussen, J. B. 2001. Why are dwarf fish so small? An
 760 energetic analysis of polymorphism in lake whitefish (*Coregonus clupeaformis*). *Can. J. Fish. Aquat.*
 761 *Sci.* 58: 394–405.

762 Venables, W. N. & Ripley, B. D. 2002. *Modern Applied Statistics with S*. Fourth Edition. Springer, New
 763 York.

764 Via, S., Bouck, A., & Skillman, S. 2000. Reproductive isolation between sympatric races of pea aphids.
 765 II. Selection against migrants and hybrids in the parental environments. *Evolution* 54: 1626-1637.

766 Vonlanthen, P., Bittner, D., Hudson, A. G., Young, K. A, Müller, R., Lundsgaard-Hansen, B. *et al.* 2012.
 767 Eutrophication causes speciation reversal in whitefish adaptive radiations. *Nature* 482: 357-363.

768 Vonlanthen, P., Roy, D., Hudson, A. G., Largiader, C. R., Bittner, D. & Seehausen, O. 2009. Divergence
 769 along a steep ecological gradient in lake whitefish (*Coregonus sp.*). *J. Evolution Biol.* 22: 498-514.

770 West Eberhard, M. J. 1989. Phenotypic plasticity and the origins of diversity. *Annu. Rev. Ecol. Syst.*
 771 20: 249-278.

772 West Eberhard, M. J. 2003. *Developmental plasticity and evolution*. Oxford University Press, Oxford.

773 Wimberger, P.H. 1994. Trophic polymorphisms, plasticity, and speciation in vertebrates. In *Theory*
 774 *and application in fish feeding ecology* (eds D. J. Stouder, K. L. Fresh & R. J. Feller). University of South
 775 Carolina Press, Columbia, SC.

776 Wund, M. A., Baker, J. A., Clancy, B., Golub, J. L. & Foster, S. A. 2008. A test of the flexible stem model
 777 of evolution: ancestral plasticity, genetic accommodation, and morphological divergence in the
 778 threespine stickleback radiation. *Am. Nat.* 172: 449-462.

779 Zelditch, M.L., Swiderski, D.L., Sheets, H.D. & Fink, W.L. 2004. Geometric Morphometrics for
780 Biologists: A Primer. Elsevier Academic Press, San Diego.

781 Zuur, A.F., Ieno, E.N., Walker, N., Saveliev, A.A. & Smith, G.M. 2009. Mixed effects models and
782 extensions in ecology with R. First edition. Springer, New York.

783

784

785

FIGURE CAPTIONS

Figure 1: Hypothetical scenarios for a) variation in feeding efficiency, if it was entirely under genetic control, b) variation in feeding efficiency, if it was entirely the results of phenotypic plasticity, c) variation in feeding efficiency, if it would have similar genetic and plastic components. Plasticity effects in panel b) and c) are adaptive. In all panels, plasticity is assumed to be equally strong in both species. Shown are the treatments on the x-axis and a hypothetical feeding efficiency value on the y-axis. Error bars are hypothetical standard deviations. In the top line of the legend to the x-axis a large fish corresponds to the benthic species and a small fish to the limnetic species; in the line below a mosquito larvae corresponds to a benthic raising environment and a zooplankton item to a limnetic raising environment; in the lowest line the first letter stands for the species (B=benthic, L=limnetic) and the second letter for the raising environment (B=benthic, L=limnetic).

Figure 2: Feeding efficiency vs. treatments and feeding efficiency vs. length. Figures on the left show the treatments (for both lakes separately) on the x-axis (see legend of Figure 1 for more detail), and figures on the right show total fish length on the x-axis. The y-axis shows time to food depletion [sec] in figures a) and b), time to first feeding [sec] in figures c) and d), and the number of unsuccessful attacks in figures e) and f). Error bars are the treatment specific standard deviations. In the figures on the left, LU stands for Lake Lucerne and Thun stands for Lake Thun. In the figures on the right, empty dots represent fish from Lake Lucerne (circles=BB; triangles=BL; squares=LB; crosses=LL) and filled dots represent fish from Lake Thun (circles=BB, squares=LB). Solid lines correspond to a linear regression line for Lake Lucerne and dashed lines to a linear regression line for Lake Thun.

Figure 3: Effects of species and the environment on feeding efficiency. Shown are the treatments on the x-axis (see legend of Figure 1 for more detail) and the residuals of the most likely model




excluding species (and in Lake Lucerne also the environment) from that model. This illustrates the effects of species and of the environment corrected for the effects of co-variables in the most likely model (residuals). Positive residuals indicate lower efficiency than predicted based on co-variables alone, while negative residuals predict higher efficiency than predicted based on co-variables alone.







a) Time to food depletion of fish from lake Lucerne: Residuals of the model “Time to food depletion = length” on the y-axis. b) Time to first feeding of fish from Lake Lucerne: Residuals of the model “Time to first feeding = length + year” on the y-axis. c) Number of unsuccessful attacks of fish from Lake Lucerne: Residuals of the model “Number of unsuccessful attacks = length + PC2 + year” on the y-axis. d) Number of unsuccessful attacks of fish from Lake Thun: Residuals of the model “Number of unsuccessful attacks = length + PC2 + year” on the y-axis. Error bars are the standard deviations of the residuals per treatment. As species was the only variable retained in the most likely model of Time to food depletion and Time to first feeding of fish from Lake Thun, no residuals could be generated and plotted against species (but see Fig. 2).

Figure 4: Effects of fish length and shape on feeding efficiency. Shown are either length (panel a-c and e) or PC2 (panel d and f) on the x-axis and the residuals of the corresponding most likely model excluding either length or PC2. This illustrates the effect of length and PC2 corrected for the effects of their co-variables in the most likely model (residuals). Positive residuals indicate lower efficiency than predicted based on co-variables, while negative residuals predict higher efficiency than predicted based on co-variables. a) Time to food depletion of fish from Lake Lucerne: Residuals of the model “Time to food depletion = species + environment” on the y-axis. b) Time to first feeding of fish from Lake Lucerne: Residuals of the model “Time to first feeding = species + environment + year” on the y-axis. c) Number of unsuccessful attacks of fish from Lake Lucerne: Residuals of the model “Number of unsuccessful attacks = species + environment + PC2 + year” on the y-axis. d) Number of unsuccessful attacks of fish from Lake Lucerne: Residuals of the model “Number of unsuccessful attacks = species + environment + length + year” on the y-axis. e) Number of unsuccessful attacks of


835 fish from Lake Thun: Residuals of the model “Number of unsuccessful attacks = species + PC2 + year”
836 on the y-axis. f) Number of unsuccessful attacks of fish from Lake Thun: Residuals of the model
837 “Number of unsuccessful attacks = species + length + year” on the y-axis. Fish head shapes (drawn
838 from a subset of landmarks) in panel d) and f) correspond to head shapes at the extremes of the PC2
839 axis and differences are threefold overdrawn. As the most likely models of time to food depletion
840 and time to first feeding of fish from Lake Thun did not contain length or a shape PC, morphological
841 effects on these efficiency measures are not illustrated.

842







a)




b)




c)




d)




a)




b)




c)



d)



e)



f)




Table 1: Sample sizes and body size variation per treatment. Four treatments were available for fish from Lake Lucerne and two treatments were available for fish from Lake Thun. The first letter of the treatment refers to genetic background of the fish and the second letter of the treatment refers to their food during raising. B stands for benthic and L stands for limnetic. The first number corresponds to fish tested in 2009 and the second number to fish tested in 2010. N_{total} includes all fish. These fish were used to test for divergence between Treatments in length. N_{shape} includes all fish for which shape data was available. These fish were used to test for divergence between treatments in shape. N_{ph} (ph= post hoc) includes all fish that started feeding, including those for which shape data was missing. These fish were used in post-hoc tests for associations of feeding efficiency with each of the explanatory variables except shape. N_{GLM} includes all fish that started feeding and for which shape data was available. These fish were used in the GLMs. Fish without shape data had to be excluded from the GLMs even when no shape variable was kept for the most likely model, because AIC is only comparable between models with the same number of observations. In the last column we report mean length of fish (mm) from a particular treatment with the respective standard deviations (with years separated using “/”).

Lake	Treatment	Genetics	Environment	N _{total}	N _{shape}	N _{np}	N _{GLM}	Mean length
Lucerne	BB	Benthic	Benthic	23/0	22/0	23/0	22/0	160 (13)
	BL	Benthic	Limnetic	21/1	20/1	21/1	20/1	151 (17)/186 (0)
	LB	Limnetic	Benthic	17/7	17/7	16/7	16/7	141 (14)/139(24)
	LL	Limnetic	Limnetic	30/0	28/0	26/0	24/0	133 (12)
	Total			91/8	87/8	86/8	82/8	
Thun	BB	Benthic	Benthic	10/7	10/7	10/7	10/7	144 (21)/154(14)
	LB	Limnetic	Benthic	10/7	10/7	10/7	10/7	139 (17)/135(9)
	Total			20/14	20/14	20/14	20/14	

843 **Table 2: Generalized linear model selection.** Models of Lake Lucerne are reported first, models of Lake Thun are reported below. Given for each model are
844 its AIC, delta AIC to the most likely model (Delta i), the likelihood of each model (Likelihood), Akaike weights (w_i) and the evidence ratio (L ratio). The
845 evidence ratio indicates how much less likely a particular model is compared to the most likely model. The model likelihood decreases for each model from
846 the top to the bottom and the most likely model is highlighted in bold. $N_{\text{unsuccessful}}$ = number of unsuccessful attacks. SP = species, ENV = raising environment,
847 L = length, PC1/PC2=principal components fish body shape variation and Y = year.

Backward model selection Lucerne	AIC	Delta i	Likelihood	w_i	L ratio
Time to food depletion = (SP×ENV)+SP+ENV+L+PC1+PC2+Y	232.67	6.39	0.04	0.02	24.41
Time to food depletion = SP+ENV+L+PC2+Y	230.68	4.4	0.11	0.06	9.03
Time to food depletion = SP+ENV+L+PC2+Y	228.88	2.6	0.27	0.14	3.67
Time to food depletion = SP+ENV+L+Y	227.44	1.16	0.56	0.28	1.79
Time to food depletion = SP+ENV+L	226.28	0	1	0.5	
Time to first feeding = (SP×ENV)+SP+ENV+L+PC1+PC2+Y	1206.6	4.4	0.11	0.07	9.03
Time to first feeding = SP+ENV+L+PC1+PC2+Y	1206.1	3.9	0.14	0.09	7.03
Time to first feeding = SP+ENV+L+PC1+Y	1204.1	1.9	0.39	0.24	2.59
Time to first feeding = SP+ENV+L+Y	1202.2	0	1	0.61	

$N_{\text{unsuccessful}} = (\text{SP} \times \text{ENV}) + \text{SP} + \text{ENV} + \text{L} + \text{PC1} + \text{PC2} + \text{Y}$	181.79	3.52	0.17	0.11	5.81
$N_{\text{unsuccessful}} = \text{SP} + \text{ENV} + \text{L} + \text{PC1} + \text{PC2} + \text{Y}$	180.26	1.99	0.37	0.24	2.7
$N_{\text{unsuccessful}} = \text{SP} + \text{ENV} + \text{L} + \text{PC2} + \text{Y}$	178.27	0	1	0.65	
Backward model selection Lake Thun	AIC	Delta i	Likelihood	Weigths	L ratio
Time to food depletion = SP+L+PC1+PC2+Y	223.13	4.48	0.11	0.04	9.39
Time to food depletion = SP+L+PC1+PC2	221.18	2.53	0.28	0.11	3.54
Time to food depletion = SP+PC1+PC2	220.07	1.42	0.49	0.19	2.03
Time to food depletion = SP+PC1	219.35	0.7	0.7	0.27	1.42
Time to food depletion = SP	218.65	0	1	0.39	
Time to first feeding = SP+ L+ PC1+PC2+Y	458.19	6.87	0.03	0.02	31.03
Time to first feeding = SP+L+ PC2+Y	456.2	4.88	0.09	0.04	11.47
Time to first feeding = SP+L+Y	454.22	2.9	0.23	0.12	4.26
Time to first feeding = SP+L	452.39	1.07	0.59	0.3	1.71
Time to first feeding = SP	451.32	0	1	0.52	
$N_{\text{unsuccessful}} = \text{SP} + \text{L} + \text{PC1} + \text{PC2} + \text{Y}$	73.2	2.97	0.23	0.18	4.41
$N_{\text{unsuccessful}} = \text{SP} + \text{L} + \text{PC2} + \text{Y}$	70.23	0	1	0.82	

849 **Table 3: Generalized linear model coefficients of the most likely models.** The different models are
850 listed in rows, the different variables are listed in columns. Abbreviations are as in Table 2. Given are
851 the estimated model coefficients (Coef), their error (error) and the p-value (p, significant values
852 highlighted in bold). A positive model coefficient indicates a positive relationship. For species and
853 environment this relationship goes from benthic to limnetic. A positive model coefficient thus means
854 that limnetic fish have a higher value than benthic fish (indicating a lower efficiency) in the response
855 variable and vice versa. For year a positive model coefficient thus means that fish in the second year
856 were less efficient. Environmentally induced effects could not be measured for Thun, which is
857 indicated by the term na.

		SP	ENV	L	PC2	Y
Time to food depletion Lucerne	Coef/error	0.41/0.2	0.7/0.17	0.01/0.005	-	-
	p	0.048	<0.001	0.065	-	-
Time to first attack Lucerne	Coef/error	0.42/0.37	0.71/0.33	-0.02/0.01	-	0.93/0.57
	p	0.26	0.03	0.08	-	0.11
N_{unsuccessful} Lucerne	Coef/error	0.2/0.25	0.05/0.21	-	9.78/6.12	-
	p	0.42	0.81	0.01	0.11	0.076
Time to food depletion Thun	Coef/error	4.55/1.95	na	-	-	-
	p	0.03	na	-	-	-
Time to first attack Thun	Coef/error	-	na	-	-	-
	p	0.56	na	-	-	-
N_{unsuccessful} Thun	Coef/error	0.03/0.36	na	-0.02/0.01	-	-39.95/13.48
	p	0.92	na	0.07	>0.01	0.06