

Supramolecular Structures in Lipid Digestion and Implications for Functional Food Delivery

Stefan Salentinig^{a,b}

^aDepartment of Chemistry, University of Fribourg, Chemin du Musée 9, 1700 Fribourg, Switzerland

^bLaboratory for Biointerfaces, Department Materials meet Life, Empa, Swiss Federal Laboratories for Materials Science and Technology, Lerchenfeldstrasse 5, 9014, St. Gallen, Switzerland.

Corresponding author email: stefan.salentinig@unifr.ch

This document is the accepted manuscript version of the following article:

Salentinig, S. (2019). Supramolecular structures in lipid digestion and implications for functional food delivery. *Current Opinion in Colloid and Interface Science*.
<https://doi.org/10.1016/j.cocis.2019.02.002>

This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

Abstract

The daily diet is important for our survival, health, and wellbeing. Functional food materials, which tailor the digestion process, can help maintaining and even improving human health and lifestyle. Knowledge on how food products, particularly food emulsions such as milk, interact with the digestive system, where they transform into supramolecular structures, can have a direct impact on the rational design of such advanced materials for functional food delivery applications. These materials have the potential to be personalized to digestive conditions and dietary nutrient requirements of the consumer or patient. They could help maintaining the uptake of co-delivered nutrients and drugs even under compromised digestion conditions such as a fat maldigestion, a low bile-salt concentration, or a limited lipase action. Such conditions are found, for instance, in preterm infants or patients with digestive disorders such as chronic pancreatitis or pancreatic insufficiency. Tailored nanostructure formation and transformation in these materials may further trigger the digestion rate, and thus have an impact on the related feeling of satiety, which may help curing eating disorders and reduce the societal challenges of obesity and related diseases.

In this contribution, the specific focus is set on discussing the equilibrium and dynamic colloidal properties of food emulsion droplets during digestion and their implications for designing nature-inspired functional food materials. These investigations provide a perspective towards the design of personalized food colloids.

1. Introduction

Designer-made supramolecular food materials may form the basis for personalized, health promoting diets of the future.¹ Delivering a complete diet with a content of hydrophobic, amphiphilic and hydrophilic nutrients, which is personalized to the needs of the consumers, could be beneficial for clinical, infant, and sport nutrition. A well-balanced nutrient content is important as, for instance, the overconsumption of lipid food components such as dietary triglycerides is related to major health disorders including heart disease, obesity, cancer, and diabetes.² In the ideal functional food material, the digestion-induced structural transformations and the produced types of colloidal structures tailor both the digestion rate and nutrient absorption by specific interactions with digestive enzymes, and the absorptive cells in the gastro-intestinal tract. Slowing down the digestion rate of colloidal food materials could support maintaining the feeling of satiety for longer and promoting the absorption of nutrients into the circulatory system of the body.³ Ultimately, the designed supramolecular structures not only deliver health promoting bioactive molecules but also protect them from chemical and physical degradation during the digestion process and storage. This is important to secure their delivery through the harsh conditions of the stomach into the small intestine, and ultimately into the circulatory system of the body.⁴⁻⁶ Examples for health promoting molecules that are sensitive to degradation during the digestion processes include proteins, peptides, polyunsaturated fatty acids (PUFAs), vitamins, and drugs.

There is growing interest in the bioavailability improvement of poorly water-soluble drugs after co-administration with lipids, as most of the recently discovered drug molecules are hydrophobic.⁷⁻¹⁰ To highlight the benefit of lipid delivery systems, particularly colloidal lipid formulations, an exceptional case is the reported increase in the bioavailability of vitamin E up to around 400% upon its administration with a lipid liquid formulation as compared to soft gelatine capsules.¹¹ In another example, the bioavailability of the drug danazol, a medication used for the treatment of endometriosis, was increased up to 900% *in vivo* in rats as well as in *in vitro* investigations, when provided in lipid-based formulations.¹² However, the underlying mechanisms at play that lead to this increase are still mostly unknown.

This review summarises the recent achievements that may guide the design of nature-inspired supramolecular functional food structures as a step towards the development of nanocarriers for the personalized delivery of nutrients. The comprehensive design of functional food materials requires a fundamental understanding of the mechanisms at play leading to structure formation, structural transformations, and morphological alterations during digestion. It is also important to gain insight into the role of interactions among the digestion-induced nanostructured materials and the digestive enzymes and cells in the gastro-intestinal tract in facilitating the

transfer of the nutrients in the body. This is only possible through recent advances in experimental techniques and sample manipulations that provide access to the mesoscopic length scale with a high spatio-temporal resolution such as time-resolved online small angle X-ray and neutron scattering (SAXS, SANS) for the real-time monitoring of dynamic structural alterations within seconds. The further combination of these techniques with microfluidics can increase the time-resolution to fractions of seconds for the *in situ* study fast-digesting samples and molecular self-assembly processes in general, as demonstrated in a recent microfluidics-SAXS study on vesicles formation.¹³ In real space, electron microscopy techniques such as cryogenic transmission electron microscopy (cryo-TEM/ cryo-SEM) are essential for gaining further insight into the morphological alterations and investigating possible coexistence of different self-assembled aggregates in the digested samples. At the same time, coupling of these methods with biochemical assays and advanced *in vitro* digestion models (e.g. relevant *in vitro* cell-culture models) is crucial to bridge the colloidal structure to the biological activity. The investigations on the structural, morphological, and size alterations of food emulsions during the digestion process, as illustrated in **Figure 1**, are important for shedding light on this possible interplay between the colloidal length-scale and the expected performance of digested emulsion droplets. Hence, thorough understanding of the mechanism behind the evolvement of self-assembled structures and the translation from basic to personalized food delivery applications is highly dependent on multidisciplinary research efforts involving scientists from different backgrounds: physics, chemistry, biology, engineering, food science, and medicine.




Figure 1: Illustration of the composition-nanostructure-activity relationship during digestion. The fundamental understanding of the link between the emulsion composition, the related digestion generated lipid-based nanostructures in the gastro-intestinal tract, and their interactions with components such as enzymes and epithelial cells in the digestive tract are crucial for the rational design of functional food materials.

1. Nature's own complete and personalized diet milk

The digestive behavior of nature's own personalized and functional food system, milk, which has been optimized over millions of years by evolution, is the key starting point in the design of advanced food materials and benchmark for nature-inspired functional foods. Milk is a complete diet that forms the basis for the survival and development of the offspring. Its lipid food components are emulsified in water in form of small micron-sized droplets, stabilized by bio-surfactants such as phospholipids and amphiphilic proteins.¹⁴⁻¹⁵ In addition to poorly water-soluble bioactive components, and metabolic messages such as human milk oligosaccharides that can modulate the gut development, milk contains components such as microRNAs that presumably regulate the further development of the offspring.¹⁶⁻¹⁸ Extracellular vesicles in milk act as nanocarriers for these labile nutrients and protects them from degradation in the digestive tract, and targets their delivery through a surface protein-mediated membrane binding.¹⁹⁻²⁰ This is of interest here as it highlights the presence of lipid-based nanocarriers in food designed by nature. Polyunsaturated fatty acids and their related di- and monoglyceride forms, which are mostly released from the breast milk triglycerides during digestion in the gastro-intestinal tract, are key elements in the development of the brain and central nervous system as well as normal vision in the infants.^{16, 21} The digestion process is a highly complex multi-stage bioprocess that secures survival by facilitating the effective delivery of the required nutrients.^{15, 22-27}

During their digestion, food emulsions such as milk undergo a range of molecular and colloidal transformations that are triggered by enzymatic reactions and stimuli such as metabolites, ions, and pH that secure the bioavailability of the nutrients and survival of the offspring.²⁸⁻³¹ Inspired by this dynamic self-assembly structure formation in milk, a detailed understanding of the fate of food emulsions during digestion can guide the design of advanced food systems that are personalized to the needs of the consumers, beyond 'simply' blueprint milk. This is also important as milk is targeted to the specific species and an increasing number of patients are diagnosed with allergies against specific milk proteins or lactose in bovine milk, which is also used as basis in many infant formulas.³²⁻³³

2. Physiology of the human digestive tract

The knowledge of how food molecules interact during their digestion is of major importance for controlling the feeling of hunger and the delivery of essential lipid food components such as lipids, vitamins, drugs or peptides with functional foods. The lipolysis of lipids is catalyzed by enzymes called lipases that hydrolyze the ester

bonds between glycerol and fatty acids, mainly in the stomach and small intestine.³⁴ Lipases are fascinating enzymes that are soluble in water but practically insoluble in their substrate, oil.³⁵ When the lipases such as pancreatic lipase in the small intestine, adsorb onto the oil-water interface of the emulsified oil droplet in normal food emulsions, a flap protecting its hydrophobic site is displaced in a process called interfacial activation, and the substrate, therefore, can reach the active site of the lipase for initiating the enzymatic reaction.³⁶⁻³⁸

The major part of lipid digestion that enhances the solubilization of lipophilic molecules occurs in the stomach and the small intestine by gastric and pancreatic lipases.^{7, 39} In the stomach, the pH (typically the local pH in the range of ~ 2.0 - 3.0), the osmolality, and the viscosity of the digested lipid foods are adjusted.^{29, 40} For food emulsions, gastric lipase in the stomach is on average responsible for the hydrolysis of up to 25% of the ester bonds between the fatty acid and glycerol moieties of administered triglycerides, generating mainly diglycerides and free fatty acids (FFAs) that accumulate on the emulsion droplet surfaces and promote the action of the pancreatic lipase in the following small intestine.^{23, 35} The major part of lipid digestion then takes place in the small intestine by lipases including the pancreatic lipase and phospholipases together with various cofactors (colipase, bile salts and calcium) at typical pH in the range of 6.0-8.0.²⁹ Gall bladder contractions lead to secretion of biliary juice containing mainly bile salts and phospholipids that help stabilize the emulsion droplets and further modify their oil-water interfaces for enhancing the adsorption of the pancreatic lipase-colipase complex.⁴¹⁻⁴⁴ It was proposed that the interactions among bile salts and colipase facilitate anchoring of a less interfacial-active pancreatic lipase presumably in the form of ternary complex that is mainly stabilized by van der Waals interactions.⁴⁴⁻⁴⁵ Adsorbed onto the emulsion droplet surfaces, pancreatic lipase stereospecifically hydrolyzes the two outer ester bonds of the triglycerides, denoted as *sn*-1 and *sn*-2 resulting in the generation of the *sn*-2 monoglyceride and the corresponding FFAs.³⁰ In this process, it was reported that the triglyceride composition affects the size distribution, the surface charge, and the microstructure of the digested emulsion droplets.⁴⁶ This indicates the presence of a link between the used oil type in food emulsions and the bioaccessibility of digestion-induced hydrophobic nutrients. Once incorporated into self-assembled structures in the presence of bile salts, the digestion products together with poorly water-soluble food components are transferred through the aqueous system of the small intestine. The generation of supramolecular structures owing to the presence amphiphilic lipids (e.g. monoglycerides and diglycerides in combination with FFAs) and biosurfactants may also play a crucial role in facilitating the delivery of health-promoting nutrients through the intestinal cell membranes and into to circulatory system of the body.

3. Self-assembly of lipid digestion products – the phase diagram approach under equilibrium conditions

Dietary fatty acids and monoglycerides are composed of hydrophobic moieties (e.g. saturated, mono- and unsaturated hydrocarbon chains (fatty acids)) and hydrophilic headgroups (e.g. carboxylic or glycerol functional groups), **Figure 2** presents the triolein digestion products sn2 monoolein (mono-unsaturated monoglyceride) and oleic acid. Depending on pH for oleic acid, and the lipid composition of systems based on monoolein, these surface-active molecules display inverse lyotropic non-lamellar liquid crystalline phases including the inverted type hexagonal (H_2) phase, the bicontinuous cubic phases ($Pn3m$ and $Im3m$ types), and the discontinuous micellar cubic $Fd3m$ phase, and inverse micelles (L_2 phase) on exposure to excess water.⁴⁷⁻⁴⁹ In a recent study, the polyunsaturated eicosapentaenoic acid monoglyceride was reported to self-assemble into H_2 structure in excess water.⁵⁰⁻⁵¹ These self-assembled, fluid-like lipid nanostructures are held together by non-covalent, intermolecular forces such as van der Waals, hydrophobic, hydrogen bonding, and electrostatic interactions between the amphiphilic molecules and the molecular shapes of oleic acid and monoolein play an important role in modulating their structural features.⁵² Dispersions of these inverse (oil-continuous) structures in excess water to form cubosomes, hexosomes, and related dispersed systems, provide attractive drug and functional food nanocarriers owing to their capability to solubilize not only hydrophobic but also hydrophilic and amphiphilic bioactive molecules.⁵³⁻⁵⁹ The encapsulation of degradation-sensitive molecules such as PUFAs, proteins, and peptides into these structures can protect them from degradation in the biological milieu, which may be crucial for facilitating their transport through the low-pH and protease-rich gastric fluid into the small intestine for absorption.^{56, 60} The solution conditions such as pH and ionic strength, that are significantly changing during digestion, are important to be taken into account as they affect the intermolecular forces and thereby modify the dynamic self-assembled structural features.^{48, 54, 61-62} For instance, in pH-tunable systems, the protonation and deprotonation of the fatty acids embedded in the water-lipid interface modify the geometric packing of the amphiphilic molecules leading to significant alterations in the self-assembled structure. Such pH responsive nanocarriers may act as sustained release systems, targeting the nutrient delivery to certain pH regions in the digestive tract. In fact, it is hypothesized that one of the driving forces for the absorption of lipid food components from the self-assembled structures into the enterocytes is the pH gradient between the bulk intestinal fluid and the local reduced pH environment of the unstirred water layer that may induce colloidal transformations in close proximity to the intestinal wall.^{23, 27, 62}

In addition to the role of the generated amphiphilic digestion products in the formation of self-assembled structures, bile salts together with amphiphilic lipids are also known to self-assemble into supramolecular structures that can solubilize hydrophobic molecules and facilitate their transport to the absorptive cells of the small intestine.^{43, 63-64} On their own, bile salts form small micelles with aggregation numbers between 4-6 for sodium cholate and 7-12 for sodium deoxycholate.⁶⁵ Bile salts also participate critically to the self-assembly of the amphiphilic digestion products, mainly monoglycerides and fatty acids, in water through localization into the lipid-water interface of their structures.⁶⁶⁻⁶⁷ For instance, the addition of bile salts to an aqueous solution of hexosomes at a monoolein/oleic acid weight ratio of 8/2 at pH = 6.8 led to a swelling of their internal H₂ structure followed by transformation to cubosomes at lipid/bile salt ratio of 17, and eventually to vesicles and direct micelles at a lipid/bile salt ratio of 8, within the reported bile salt concentrations found in the small intestine.⁶⁶ These results suggest that the addition of bile salts within the pancreatic juice during digestion in the small intestine favors the formation of more hydrophilic interfaces and ultimately direct micelles. It is worth noting that contrary to the structure-forming lipid digestion products such as sn-2 monoglycerides and fatty acids, bile salts are not up-taken by the enterocytes in the small intestine, but later in the ileum.⁶⁸ Hence, significant changes in the self-assembled structures, potentially even a back-tuning to inverse liquid crystalline phases may occur also at the enterocytes-membrane upon bile salt 'removal', based on the composition of the lipid digestion products and the reduced local pH environment.




Figure 2: The triolein digestion products sn-2 monoolein and oleic acid. Both molecules are amphiphilic, containing a hydrophobic tail and hydrophilic headgroup covalently linked in one molecule. Self-assembled structures including inverse micellar phases (L₂ and Fd3m), inverse hexagonal (H₂)-, bicontinuous cubic (Pn3m type) structure and vesicles are schematically presented in the figure with water in blue-, and oil in yellow- and

orange color. The structures arise from intermolecular forces and electrostatic interactions among the amphiphilic molecules in excess water. Related to electrostatic interactions, the 'apparent' pK_a of oleic acid in self-assembled structures was found between about 6 and 7, that is within the physiological pH range found in the small intestine.⁴⁸ Thus, if the solution pH is changed, the electrostatic forces, and consequently the colloidal structures are modified.

4. Dynamic self-assembly during emulsion digestion studied *in situ*

Lipid digestion is a dynamic process with constant modifications in the lipid composition over time including triglycerides, diglycerides, monoglycerides and fatty acids. This section discusses the dynamic assembly of these products during digestion *in situ*. These research studies require suitable *in vitro* digestion models coupled with highly contemporary small-angle scattering approaches, and electron microscopy techniques to unravel the dynamic nanostructure formation and the structural transition mechanisms with high resolution in time and space.^{66, 69-71}

Structural changes on the surface of the digesting olive-oil droplets were first demonstrated with an *in vitro* intestinal digestion model coupled with light microscopy.⁷² However, a detailed structural assignment was not possible with this technique. The dynamic formation and assignment of highly ordered lyotropic liquid crystalline structures in model triolein-in-water emulsion droplets was then demonstrated using a suitable *in vitro* intestinal digestion model coupled with online SAXS and cryo-TEM.⁶⁶ Recently, these structural transformations were also observed in bovine and human breast milks and the triglyceride-based food product mayonnaise, in which oleic acid is also a dominating lipid component.^{31, 69} Considering that the self-assembled structures are very sensitive to modifications in their environment such as composition, salt concentration, and pH, it is a surprising result to find them in such digested complex food systems in the presence of multiple hydrophilic, hydrophobic and amphiphilic components including proteins, carbohydrates, lipids, DNA, and even cells. The discovered biogenerated structures and their transformations with time of digestion are in-line with the expectations from the composition- and pH-dependent self-assembly of oleic acid and monoolein under equilibrium conditions, as discussed above. With time of digestion at pH 6.5, transitions from normal emulsion droplets through a variety of colloidal objects with differently internally ordered nanostructures, including the L_2 phase, the $Fd3m$ structure, the H_2 phase, and the bicontinuous cubic $Pn3m$ and $Im3m$ phases, to vesicles and normal micelles were observed in all systems in absence and at low bile-salt concentrations (**Figure 3**). **Figure**

4 shows the time-resolved SAXS data for mayonnaise and cow milk during digestion with pancreatin extract from the porcine pancreas without added bile salts. The sequence of structures is similar between the different food systems. The up to 4 times slower rate of structure formation in mayonnaise compared to the homogenized cow milk may be mainly caused by the larger emulsion droplet size in the mayonnaise and variation in composition. SAXS data during the digestion of triolein emulsions at elevated bile-salt levels and elevated pH of 7.5 indicated their transformation to normal micelles and vesicles without intermediate non-lamellar liquid crystalline structures.^{70, 66} However, as SAXS is a statistical method providing an average over all nanoparticles in the sample volume, the presence of a minor fraction of liquid crystalline nanoparticles as intermediate structures cannot be excluded even at high bile salt- or pH levels. These transformations from normal emulsions to fatty acid and monoglyceride-based inverse liquid crystalline structures and further to vesicles and direct (normal) micelles also imply that water molecules start to diffuse into the droplets' interiors and create water pockets, which are eventually transformed into a network of hydrophilic domains in the internal oil-continuous liquid crystalline structures of the formed colloidal objects, see illustration in Figure 3. The lipase-driven transfer of water into the triglyceride (oil) phase at the macroscopic oil-water interface was recently demonstrated experimentally with ellipsometry.⁷³ The location and activity of the lipase may be triggered by the formation of nanostructures with the extensive water-domains inside the oil droplets during digestion. These highly ordered nanostructures were observed during the digestion of long-chain triglycerides, medium chain triglycerides (MCT) were found to be digested much faster and direct transformations to vesicles and micelles were reported.⁶⁶ The bioaccessibility was found to strongly depend on the emulsion oil type, with medium chain triglycerides resulting in low – and long chain triglycerides resulting in high bioavailability of the hydrophobic vitamin D.⁷⁴ Further, clinical nutrition studies with formulas containing medium chain triglycerides, where liquid crystalline structures are not present during digestion, showed a decrease in the uptake of fat-soluble vitamins in patients.⁷⁵

Towards this end, the question arises if designer-made nanostructured particles with liquid crystalline interiors can be engineered to induce specific phase transformations that tailor the digestion process and the delivery of loaded nutrients in defined and specific locations in the digestive tract, for instance the induction of certain phase transitions through responses to specific enzymes. Recently, particles with liquid crystalline *Fd3m* digestion structure (micellar cubosomes) were formulated with lipase-specific phase transition patterns.⁷⁶ In a system containing nanoparticles with a 1/1 mixture of phospholipid and diglycerides, the cubic *Fd3m* structure was transformed into more hydrophobic inverse micelles (L_2 phase) with phospholipase and into more

hydrophilic vesicles with pancreatic lipase. Also pH has a significant effect on the dynamic structural transformations as it was demonstrated during the digestion of monoolein-based cubosomes by using *thermomyces lanuginosus* lipase.⁷⁷ The inner cubic *Im3m* structure transformed to H_2 phase, cubic *Fd3m* phase, and L_2 phase at pH between 6.5 and 7.5. Contrary, at increased pH above 8.0, transformations from cubosomes with an internal cubic *Im3m* phase to sponge-like structure (L_3 phase) and multilamellar vesicles (MLVs) were reported.⁷⁷ These differences in structures were discussed in relation to the protonation of the produced oleic acid that may trigger a more negative curvature of the oil-water interface further supporting the role of the pH as one of the key parameters for nanostructure formation and structural transformations in the digestion process of emulsified lipid systems such as food emulsions.^{31, 48, 62, 78} In this context, further important parameters include the ionic strength, specifically also the calcium concentration during digestion as calcium ions can form insoluble complexes with fatty acids (soaps) that precipitate from the solution and thus impact on digestion rate and extent of digestion.⁷⁹ Hence, these approaches pave the path towards the design of enzyme- and pH-triggered biointerfaces with potential for sustained nutrient release and tailored interactions with the cells in the digestive tract. However, the important impact of these structures on lipid absorption, and its implications for the rational design of new healthy functional foods is still poorly understood.^{23, 80}

It is worth highlighting at this point that the current *in vitro* digestion models applied for the study of nanostructure formation so far in food emulsions and other related lipid dispersions neglect the dynamic changes in composition through the absorption of the digestion products into the absorptive cells of the intestine, occurring during the *in vivo* digestion. Overall, as the *in vivo* conditions are highly complex, a comprehensive understanding of the role of individual process parameters is only possible with well-defined *in vitro* models at reduced complexity. Such parameters include emulsion properties, digestive juice composition, pH, selected lipases, and interactions with cells. In this research area, the next major step towards a more thorough understanding of the digestion-triggered nanostructure formation and its implications on nutrient uptake requires coupling the current online SAXS *in vitro* digestion model with gastric pre-digestion and physiologically relevant cell culture models. These cell-culture models should approximate the structure of the intestinal epithelium such as Caco-2 cells co-cultured with mucus secreting cells.⁸¹⁻⁸⁵




Figure 3: (a) Cryo-TEM image of store-bought homogenized milk showing milk emulsion droplets in water (1*) in combination with casein micelles (2*). (b) Shows the appearance of internally nanostructured particles, containing a bicontinuous cubic phase (1), that coexist with vesicles (2) and direct micelles (3) at the final state of milk digestion in porcine pancreatin solution without additional bile salts. The inset shows the fast Fourier transform of the structured region in the cryo-TEM with the diffraction pattern that is characteristic for the bicontinuous cubic structure. The graphical illustration sketches the nanostructure formation inside the milk emulsion droplet during digestion. The internal oil structure of the normal emulsion is dynamically transformed via inverse micellar phases to, inverse hexagonal and bicontinuous cubic structure during digestion. Figure adapted from Ref.^{69, 71}




Figure 4: *In situ* SAXS data recorded during the digestion of mayonnaise (a) and store-bought homogenized cow milk (b). The formation of highly geometrically ordered liquid crystalline structures was observed in both food emulsions. However, the Bragg reflections of the lamellar phase may most likely be attributed to a difference in the lipid type and composition in mayonnaise as compared to milk. For instance, milk contains a significant amount of saturated fatty acids that may form coexisting multilamellar structures. Figure adapted from Ref.^{31, 71}

5. Implications of supramolecular structures for healthy food

Enzyme and pH responsive supramolecular food materials are interesting concepts for tailoring the delivery of specific nutrients as well as the digestion profile to the needs of the needs of the consumer. For instance, switching from protected mode to active mode as the developed nanocarrier migrates from the stomach into the small intestine could be used to transport labile molecules through the harsh and degradative environment of the stomach to the small intestine for uptake into the circulatory system. Related to this, pH responsive nanocarriers of antimicrobial peptides have been designed with the triolein digestion products glycerol monooleate and oleic acid as structure forming molecules.^{54, 56} In this system, cubosomes did not interact with bacteria cells as their encapsulation within the curved bilayers of the internal cubic phase in cubosomes hinders their access to the bacterial membrane. However, in form of micelles and vesicles at pH around 7.0, the peptides are exposed at the outer interface and the nanostructures are highly antibacterial active.

These lipid-based supramolecular materials may help maintaining the transport and absorption of lipid food components including hydrophobic vitamins, carotenoids and drugs under compromised digestion conditions such as in preterm infants and patients with exocrine pancreatic insufficiency or chronic pancreatitis.⁸⁶ Here, pancreatic lipase loaded functional food structures could be designed to transport the pH-sensitive enzyme through the stomach into the small intestine, where it can expel its activity. A concept for pancreatic lipase delivery based on pH responsive hydrogel beads has been demonstrated recently.⁵ The major advantage of the lipid-based nanocarriers is the potential for co-administering further hydrophilic and hydrophobic nutrients and the lipid building blocks of the nanocarriers that are of high demand for the body under these conditions.

Tailoring digestion rates through structure formation and transformation with designer-made food materials may provide also be an interesting approach to trigger the feeling of satiety and hunger for improved diets.⁸⁷ Clinical studies showed that the consumption of milk significantly increased satiety and decreased energy intake in a

second follow-up meal.⁸⁸ The underlying mechanisms are not yet understood but the formation of lipid structures in combination with proteins may play a major role in this process. A detailed understanding of the nanostructure formation and transformation in milk, and its implications on the activity and interaction with the lipases during digestion can guide the development of functional food materials. This may open an alternative route to current approaches that mainly focus on engineering the surfaces of emulsion droplets, for instance with hydrogels, that inhibit or delay lipase access *in vitro*.^{87, 89}

6. Conclusions

The self-assembly of the triglyceride digestion products of food emulsions such as milk leads to a rich variety of nanostructures including highly geometrically ordered liquid crystalline structures, depending on composition, temperature, and environmental conditions during digestion such as pH, bile salt levels, and ionic strength. These supramolecular structures possess unique properties including extensive lipid-water interfacial areas and ultra-low surface tensions allowing the spontaneous nanostructure formation. Hence, the formation of these structures *in situ* during digestion of these emulsions may not only help solubilizing poorly water-soluble nutrients but also tailor the interactions with cell membranes in the intestine, and secure lipid digestion under compromised conditions such as low bile salt levels. Towards this end, a detailed understanding of the fate of the nature's personalized diet milk during digestion with focus on composition, nanostructure generation, and interactions with the epithelial cells of the digestive tract may provide crucial inspirations for advanced functional food materials.

Similar to milk, designer made lipid-based nanocarriers may be optimized for the transport and absorption of lipid food components into the epithelial cells in the small intestine under compromised digestion conditions. This would allow the tailored delivery of degradation sensitive nutrients such as vitamins, polyunsaturated fatty acids, and peptides through the stomach to the epithelial cells of the small intestine for absorption. In this context, the adaption of *in situ* scattering techniques for the time-resolved study of nanostructural transformations opens a new research area on the colloidal aspects of food digestion and the design of novel functional food delivery systems.

In the future, ideal supramolecular food nanocarriers are expected to be personalized by considering the digestive conditions of the consumer that include composition, pH profile, and lipase activity. Combined with a detailed understanding of the self-assembly of the food components during digestion, as well as their specific

interactions with lipases and the intestinal cells, this will allow tailoring the digestion profile and the formation of most efficient structures for the delivery of essential nutrients and drugs to the epithelial cells of the small intestine.

References (papers of special interest (•) or outstanding interest (••))

1. J. E. Norton; Y. Gonzalez Espinosa; R. L. Watson; F. Spyropoulos; I. T. Norton, Functional food microstructures for macronutrient release and delivery, *Food. Funct.* 6 (2015) 663-678.
2. B. G. Nordestgaard; A. Varbo, Triglycerides and cardiovascular disease, *Lancet* 384 (2014) 626-635.
- 3. M. N. Corstens; C. C. Berton-Carabin; R. de Vries; F. J. Troost; A. A. Masclee; K. Schroen, Food-grade micro-encapsulation systems that may induce satiety via delayed lipolysis: A review, *Crit. Rev. Food Sci. Nutr.* 57 (2017) 2218-2244. This article reviews strategies to delay lipolysis to induce satiety. It discusses the design of oil-water interfaces for controlled lipase access as a promising route.
4. N. P. Aditya; Y. G. Espinosa; I. T. Norton, Encapsulation systems for the delivery of hydrophilic nutraceuticals: Food application, *Biotechnol. Adv.* 35 (2017) 450-457.
- 5. Z. Zhang; F. Chen; R. Zhang; Z. Deng; D. J. McClements, Encapsulation of pancreatic lipase in hydrogel beads with self-regulating internal pH microenvironments: Retention of lipase activity after exposure to gastric conditions, *J. Agric. Food Chem.* 64 (2016) 9616-9623. This study demonstrates the design of alginate-based polymer beads to deliver pancreatic lipase intact through the degradative low-pH environment of the stomach. The pH inside the buffer-loaded beads remained almost neutral even in the gastric environment, and lipase activity was retained after exposure to these conditions.
6. D. J. McClements, Encapsulation, protection, and delivery of bioactive proteins and peptides using nanoparticle and microparticle systems: A review, *Adv. Colloid Interface Sci.* 253 (2018) 1-22.
- 7. M. Koziolok; F. Carriere; C. J. H. Porter, Lipids in the stomach - implications for the evaluation of food effects on oral drug absorption, *Pharm. Res.* 35 (2018) 55. This article reviews the impact of lipids on drug solubilisation under gastric conditions and discusses its implications for the following intestinal digestion and drug absorption.
8. C. W. Pouton, Formulation of poorly water-soluble drugs for oral administration: Physicochemical and physiological issues and the lipid formulation classification system, *Eur. J. Pharm. Sci.* 29 (2006) 278-287.
9. C. J. Porter; N. L. Trevaskis; W. N. Charman, Lipids and lipid-based formulations: Optimizing the oral delivery of lipophilic drugs, *Nat. Rev. Drug Discov.* 6 (2007) 231-248.
10. A. Mullard, 2017 FDA drug approvals, *Nat. Rev. Drug Discov.* 17 (2018) 81-85.
11. T. Julianto; K. H. Yuen; A. M. Noor, Improved bioavailability of vitamin E with a self emulsifying formulation, *Int. J. Pharm.* 200 (2000) 53-57.
12. A. Larsen; R. Holm; M. L. Pedersen; A. Mullertz, Lipid-based formulations for danazol containing a digestible surfactant, labrafil m2125cs: In vivo bioavailability and dynamic in vitro lipolysis, *Pharm. Res.* 25 (2008) 2769-2777.
- 13. A. Ghazal; M. Gontsarik; J. P. Kutter; J. P. Lafleur; D. Ahmadvand; A. Labrador; S. Salentinig; A. Yagmur, Microfluidic platform for the continuous production and characterization of multilamellar vesicles: A synchrotron small-angle x-ray scattering (saxs) study, *J. Phys. Chem. Lett.* 8 (2017) 73-79. This article gives insights into the formation of multilamellar vesicles using a novel microfluidic platform that allows *in situ* online synchrotron SAXS studies with high spatio-temporal resolution.
14. E. Dickinson, Biopolymer-based particles as stabilizing agents for emulsions and foams, *Food Hydrocoll.* 68 (2017) 219-231.

15. D. J. McClements, Enhanced delivery of lipophilic bioactives using emulsions: A review of major factors affecting vitamin, nutraceutical, and lipid bioaccessibility, *Food Funct.* 9 (2018) 22-41.
16. L. A. Milligan, Do bigger brains mean better milk? In: *Building babies: Primate development in proximate and ultimate perspective*, Clancy, K. B. H.; Hinde, K.; Rutherford, J. N., Eds. Springer New York: New York, NY, 2013; pp 209-231.
17. C. Admyre; S. M. Johansson; K. R. Qazi; J. J. Filen; R. Lahesmaa; M. Norman; E. P. Neve; A. Scheynius; S. Gabrielsson, Exosomes with immune modulatory features are present in human breast milk, *J. Immunol.* 179 (2007) 1969-1978.
18. T. Gura, Nature's first functional food, *Science* 345 (2014) 747-749.
- 19. J. Zemleni; A. Aguilar-Lozano; M. Sadri; S. Sukreet; S. Manca; D. Wu; F. Zhou; E. Mutai, Biological activities of extracellular vesicles and their cargos from bovine and human milk in humans and implications for infants, *J. Nutr.* 147 (2017) 3-10. This article gives an overview of biological activities of extracellular vesicles and their role in protecting their cargo from degradation. It also discusses the delivery of RNAs in bovine-milk exosomes to circulating immune cells in humans.
- 20. S. Kahn; Y. Liao; X. Du; W. Xu; J. Li; B. Lonnerdal, Exosomal micrornas in milk from mothers delivering preterm infants survive in vitro digestion and are taken up by human intestinal cells, *Mol. Nutr. Food Res.* 62 (2018) e1701050. This study demonstrates the role of exosomes from breast milk in protecting microRNA from degradation during simulated gastric/pancreatic digestion. It further discusses the biology and the molecular basis by which exosome miRNAs may uniquely affect the development of preterm infants.
21. W. S. Harris; M. L. Baack, Beyond building better brains: Bridging the docosahexaenoic acid (dha) gap of prematurity, *J. Perinatol.* 35 (2015) 1-7.
22. D. J. McClements; Y. Li, Structured emulsion-based delivery systems: Controlling the digestion and release of lipophilic food components, *Adv. Colloid Interface Sci.* 159 (2010) 213-228.
23. C. J. H. Porter; N. L. Trevaskis; W. N. Charman, Lipids and lipid-based formulations: Optimizing the oral delivery of lipophilic drugs, *Nat. Rev. Drug Discov.* 6 (2007) 231-248.
- 24. Y. Yang; H. Xiao; D. J. McClements, Impact of lipid phase on the bioavailability of vitamin E in emulsion-based delivery systems: Relative importance of bioaccessibility, absorption, and transformation, *J. Agric. Food Chem.* 65 (2017) 3946-3955. This article shows that long-chain triglyceride emulsions are more suitable for vitamin E delivery than medium chain triglyceride emulsions. Higher vitamin E bioaccessibility and higher rates of esterase-catalysed transformation of the α -tocopherol acetate to α -tocopherol were observed.
25. Y. Yang; D. J. McClements, Vitamin e bioaccessibility: Influence of carrier oil type on digestion and release of emulsified α -tocopherol acetate, *Food Chem.* 141 (2013) 473-481.
26. L. Salvia-Trujillo; S. H. Verkempinck; L. Sun; A. M. Van Loey; T. Grauwet; M. E. Hendrickx, Lipid digestion, micelle formation and carotenoid bioaccessibility kinetics: Influence of emulsion droplet size, *Food Chem.* 229 (2017) 653-662.
27. H. D. Williams; N. L. Trevaskis; Y. Y. Yeap; M. U. Anby; C. W. Pouton; C. J. Porter, Lipid-based formulations and drug supersaturation: Harnessing the unique benefits of the lipid digestion/absorption pathway, *Pharm. Res.* 30 (2013) 2976-2992.
28. H. Singh; A. Ye; D. Horne, Structuring food emulsions in the gastrointestinal tract to modify lipid digestion, *Prog. Lipid Res.* 48 (2009) 92-100.
29. M. Golding; T. J. Wooster, The influence of emulsion structure and stability on lipid digestion, *Curr. Opin. Colloid Interface Sci.* 15 (2010) 90-101.
30. S. Salentinig; N. R. Yepuri; A. Hawley; B. J. Boyd; E. Gilbert; T. A. Darwish, Selective deuteration for molecular insights into the digestion of medium chain triglycerides, *Chem. Phys. Lipids* 190 (2015) 43-50.

- 31. S. Salentinig; H. Amenitsch; A. Yagmur, In situ monitoring of nanostructure formation during the digestion of mayonnaise, *ACS Omega* 2 (2017) 1441-1446. This study researches for the first time the formation of lyotropic liquid crystalline phases during the *in vitro* digestion of mayonnaise at low bile-salt concentrations. It demonstrates the effect of the gradual pH increase in the intestine on the digestion generated nanostructures.
- 32. A. Golkar; J. M. Milani; T. Vasiljevic, Altering allergenicity of cow's milk by food processing for applications in infant formula, *Crit. Rev. Food Sci. Nutr.* (2018) 1-14.
- 33. P. H. F. da Silva; V. C. D. Oliveira; L. M. Perin, Chapter 14 - cow's milk protein allergy and lactose intolerance. In: Raw milk, Nero, L. A.; De Carvalho, A. F., Eds. Academic Press: 2019; pp 295-309.
- 34. J. C. Bakala N'Goma; S. Amara; K. Dridi; V. Jannin; F. Carriere, Understanding the lipid-digestion processes in the GI tract before designing lipid-based drug-delivery systems, *Ther. Deliv.* 3 (2012) 105-124.
- 35. P. Reis; K. Holmberg; H. Watzke; M. E. Leser; R. Miller, Lipases at interfaces: A review, *Adv. Colloid Interface Sci.* 147-48 (2009) 237-250.
- 36. R. Verger, 'Interfacial activation' of lipases: Facts and artifacts, *Trends Biotechnol.* 15 (1997) 32-38.
- 37. F. K. Winkler; A. D'Arcy; W. Hunziker, Structure of human pancreatic lipase, *Nature* 343 (1990) 771-774.
- 38. C. Cheng; T. Jiang; Y. Wu; L. Cui; S. Qin; B. He, Elucidation of lid open and orientation of lipase activated in interfacial activation by amphiphilic environment, *Int. J. Biol. Macromol.* 119 (2018) 1211-1217. This study focuses on the effect of the oil-water interface on lipase activation using molecular dynamic simulations. It demonstrates that the hydrophobic regions of lipase, located at the "lid" and "flap" regions, tailor the orientation of the active center towards the oil-water interface to achieve enzyme activation.
- 39. F. Carriere, Impact of gastrointestinal lipolysis on oral lipid-based formulations and bioavailability of lipophilic drugs, *Biochimie* 125 (2016) 297-305. This article discusses *in vitro* models to study oral lipid formulations for drug delivery. It highlights the importance of gastric lipolysis that is mostly ignored in current models, to understand the fate of lipid based formulations during digestion.
- 40. N. Scheuble; A. Iles; R. C. R. Wootton; E. J. Windhab; P. Fischer; K. S. Elvira, Microfluidic technique for the simultaneous quantification of emulsion instabilities and lipid digestion kinetics, *Anal. Chem.* 89 (2017) 9116-9123. This study demonstrates the application of microfluidic approaches to quantify the coalescence of oil-in-water emulsion droplets during lipid digestion on a single droplet level. Using this methodology it was shown that protein stabilised emulsion coalesced under human gastric conditions which delayed the overall lipid digestion kinetics.
- 41. M. Wickham; M. Garrood; J. Leney; P. D. Wilson; A. Fillery-Travis, Modification of a phospholipid stabilized emulsion interface by bile salt: Effect on pancreatic lipase activity, *J. Lipid Res.* 39 (1998) 623-632.
- 42. A. F. Hofmann, Bile acids: The good, the bad, and the ugly, *News Physiol. Sci.* 14 (1999) 24-29.
- 43. A. Sarkar; A. Q. Ye; H. Singh, On the role of bile salts in the digestion of emulsified lipids, *Food Hydrocoll.* 60 (2016) 77-84.
- 44. R. Parker; N. M. Rigby; M. J. Ridout; A. P. Gunning; P. J. Wilde, The adsorption-desorption behaviour and structure function relationships of bile salts, *Soft Matter* 10 (2014) 6457-6466.
- 45. N. Haque; N. Prakash Prabhu, Binding orientation and interaction of bile salt in its ternary complex with pancreatic lipase-colipase system, *Biochem. Biophys. Res. Commun.* 499 (2018) 907-912.
- 46. R. Zhang; Z. Zhang; H. Zhang; E. A. Decker; D. J. McClements, Influence of lipid type on gastrointestinal fate of oil-in-water emulsions: In vitro digestion study, *Food Res. Int.* 75 (2015) 71-78.

47. S. Mele; O. Soderman; H. Ljusberg-Wahren; K. Thuresson; M. Monduzzi; T. Nylander, Phase behavior in the biologically important oleic acid/sodium oleate/water system, *Chem. Phys. Lipids* 211 (2018) 30-36.
48. S. Salentinig; L. Sagalowicz; O. Glatter, Self-assembled structures and pka value of oleic acid in systems of biological relevance, *Langmuir* 26 (2010) 11670-11679.
49. H. Qiu; M. Caffrey, The phase diagram of the monoolein/water system: Metastability and equilibrium aspects, *Biomaterials* 21 (2000) 223-234.
50. X. Shao; G. Bor; S. Al-Hosayni; S. Salentinig; A. Yaghmur, Structural characterization of self-assemblies of new omega-3 lipids: Docosahexaenoic acid and docosapentaenoic acid monoglycerides, *Phys. Chem. Chem. Phys.* 20 (2018) 23928-23941.
51. A. Yaghmur; S. Al-Hosayni; H. Amenitsch; S. Salentinig, Structural investigation of bulk and dispersed inverse lyotropic hexagonal liquid crystalline phases of eicosapentaenoic acid monoglyceride, *Langmuir* 33 (2017) 14045-14057.
52. J. N. Israelachvili, *Intermolecular and surface forces*, third ed., Academic press, Oxford, 2011.
53. A. Angelova; V. M. Garamus; B. Angelov; Z. Tian; Y. Li; A. Zou, Advances in structural design of lipid-based nanoparticle carriers for delivery of macromolecular drugs, phytochemicals and anti-tumor agents, *Adv. Colloid Interface Sci.* 249 (2017) 331-345.
54. M. Gontsarik; M. Mohammadtaheri; A. Yaghmur; S. Salentinig, pH-triggered nanostructural transformations in antimicrobial peptide/oleic acid self-assemblies, *Biomater. Sci.* 6 (2018) 803-812.
55. S. Salentinig; K. J. Tangso; A. Hawley; B. J. Boyd, pH-driven colloidal transformations based on the vasoactive drug nicergoline, *Langmuir* 30 (2014) 14776-14781.
56. M. Gontsarik; M. T. Buhmann; A. Yaghmur; Q. Ren; K. Maniura-Weber; S. Salentinig, Antimicrobial peptide-driven colloidal transformations in liquid-crystalline nanocarriers, *J. Phys. Chem. Lett.* 7 (2016) 3482-3486.
57. L. Sagalowicz; C. Moccand; T. Davidek; R. Ghanbari; I. Martiel; R. Negrini; R. Mezzenga; M. E. Leser; I. Blank; M. Michel, Lipid self-assembled structures for reactivity control in food, *Philos. Trans. A Math. Phys. Eng. Sci.* 374 (2016).
58. L. Sagalowicz; M. Michel; I. Blank; O. Schafer; M. E. Leser, Self-assembly in food — a concept for structure formation inspired by nature, *Curr. Opin. Colloid Interface Sci.* 28 (2017) 87-95.
59. A. Chemelli; M. Maurer; R. Geier; O. Glatter, Optimized loading and sustained release of hydrophilic proteins from internally nanostructured particles, *Langmuir* 28 (2012) 16788-16797.
- 60. L. Boge; A. Umerska; N. Matougui; H. Bysell; L. Ringstad; M. Davoudi; J. Eriksson; K. Edwards; M. Andersson, Cubosomes post-loaded with antimicrobial peptides: Characterization, bactericidal effect and proteolytic stability, *Int. J. Pharm.* 526 (2017) 400-412. This article shows that cubosomes, loaded with antimicrobial peptides that are sensitive to degradation in the biological environment, can protect their sensitive load from proteolytic degradation.
61. R. Negrini; R. Mezzenga, pH-responsive lyotropic liquid crystals for controlled drug delivery, *Langmuir* 27 (2011) 5296-5303.
62. S. Salentinig; S. Phan; T. A. Darwish; N. Kirby; B. J. Boyd; E. P. Gilbert, pH-responsive micelles based on caprylic acid, *Langmuir* 30 (2014) 7296-7303.
63. O. Hernell; J. E. Staggers; M. C. Carey, Physical-chemical behavior of dietary and biliary lipids during intestinal digestion and absorption. 2. Phase analysis and aggregation states of luminal lipids during duodenal fat digestion in healthy adult human beings, *Biochemistry* 29 (1990) 2041-2056.

64. P. Schurtenberger; N. Mazer; W. Kanzig, Micelle to vesicle transition in aqueous-solutions of bile-salt and lecithin, *J. of Phys. Chem.* 89 (1985) 1042-1049.
65. D. Madenci; S. U. Egelhaaf, Self-assembly in aqueous bile salt solutions, *Curr. Opin. Colloid Interface Sci.* 15 (2010) 109-115.
66. S. Salentinig; L. Sagalowicz; M. E. Leser; C. Tedeschi; O. Glatter, Transitions in the internal structure of lipid droplets during fat digestion, *Soft Matter* 7 (2011) 650-661.
67. A. Sadeghpour; M. Rappolt; S. Misra; C. V. Kulkarni, Bile salts caught in the act: From emulsification to nanostructural reorganization of lipid self-assemblies, *Langmuir* 34 (2018) 13626-13637.
68. P. A. Dawson; S. J. Karpen, Intestinal transport and metabolism of bile acids, *J. Lipid Res.* 56 (2015) 1085-1099.
69. S. Salentinig; S. Phan; A. Hawley; B. J. Boyd, Self-assembly structure formation during the digestion of human breast milk, *Angew. Chem. Int. Ed.* 54 (2015) 1600-1603.
70. S. Marze; C. Gaillard; P. Roblin, In vitro digestion of emulsions: High spatiotemporal resolution using synchrotron SAXS, *Soft Matter* 11 (2015) 5365-5373.
71. S. Salentinig; S. Phan; J. Khan; A. Hawley; B. J. Boyd, Formation of highly organized nanostructures during the digestion of milk, *ACS Nano* 7 (2013) 10904-10911.
72. J. Patton; M. Carey, Watching fat digestion, *Science* 204 (1979) 145-148.
73. A. Stamm; A. Svendsen; J. Skjold-Jorgensen; T. Vissing; I. Berts; T. Nylander, The triolein/aqueous interface and lipase activity studied by spectroscopic ellipsometry and coarse grained simulations, *Chem. Phys. Lipids* 211 (2018) 37-43.
- 74. B. Ozturk; S. Argin; M. Ozilgen; D. J. McClements, Nanoemulsion delivery systems for oil-soluble vitamins: Influence of carrier oil type on lipid digestion and vitamin D3 bioaccessibility, *Food Chem.* 187 (2015) 499-506. This article highlights the importance of oil-type for the *in vitro* bioaccessibility of vitamin D3. Emulsified long chain triglycerides were very effective at increasing vitamin bioaccessibility, compared to medium chain triglycerides.
75. S. S. Christodoulides; E. G. Neal; G. Fitzsimmons; H. M. Chaffe; Y. M. Jeanes; H. Aitkenhead; J. H. Cross, The effect of the classical and medium chain triglyceride ketogenic diet on vitamin and mineral levels, *J. Hum. Nutr. Diet* 25 (2012) 16-26.
- 76. M. Wadsater; J. Barauskas; F. Tiberg; T. Nylander, The lipolytic degradation of highly structured cubic micellar nanoparticles of soy phosphatidylcholine and glycerol dioleate by phospholipase a2 and triacylglycerol lipase, *Chem. Phys. Lipids* 211 (2018) 86-92. This study describes the design of micellar cubosomes with lipase-specific phase transition patterns. Using a 1/1 mixture of phospholipid and diglycerides, the structures were transformed into more hydrophobic inverse micelles (L2 phase) or into more hydrophilic vesicles depending on the type of lipase.
77. J. Barauskas; H. Anderberg; A. Svendsen; T. Nylander, *Thermomyces lanuginosus* lipase-catalyzed hydrolysis of the lipid cubic liquid crystalline nanoparticles, *Colloids Surf. B Biointerfaces* 137 (2016) 50-59.
78. S. Phan; S. Salentinig; E. Gilbert; T. A. Darwish; A. Hawley; R. Nixon-Luke; G. Bryant; B. J. Boyd, Disposition and crystallization of saturated fatty acid in mixed micelles of relevance to lipid digestion, *J. Colloid Interface Sci.* 449 (2015) 160-166.
79. M. Hu; Y. Li; E. A. Decker; D. J. McClements, Role of calcium and calcium-binding agents on the lipase digestibility of emulsified lipids using an in vitro digestion model, *Food Hydrocoll.* 24 (2010) 719-725.
- 80. D. J. McClements, The future of food colloids: Next-generation nanoparticle delivery systems, *Curr. Opin. Colloid Interface Sci.* 28 (2017) 7-14. This article explores the role of colloid science in food science

discussing implications on nutrient delivery, food safety, packaging and quality. It further highlights the importance of multidisciplinary approaches in this complex field of research.

81. P. Artursson; K. Palm; K. Luthman, Caco-2 monolayers in experimental and theoretical predictions of drug transport, *Adv. Drug Deliv. Rev.* 46 (2001) 27-43.
82. B. M. Folmer; D. Barron; E. Hughes; L. Miguet; B. Sanchez; O. Heudi; M. Rouvet; L. Sagalowicz; P. Callier; M. Michel; G. Williamson, Monocomponent hexa- and dodecaethylene glycol succinyl-tocopherol esters: Self-assembly structures, cellular uptake and sensitivity to enzyme hydrolysis, *Biochem. Pharmacol.* 78 (2009) 1464-1474.
83. C. Hilgendorf; H. Spahn-Langguth; C. G. Regardh; E. Lipka; G. L. Amidon; P. Langguth, Caco-2 versus caco-2/ht29-mtx co-cultured cell lines: Permeabilities via diffusion, inside- and outside-directed carrier-mediated transport, *J. Pharm. Sci.* 89 (2000) 63-75.
84. G. J. Mahler; M. L. Shuler; R. P. Glahn, Characterization of caco-2 and ht29-mtx cocultures in an in vitro digestion/cell culture model used to predict iron bioavailability, *J. Nutr. Biochem.* 20 (2009) 494-502.
85. L. M. Ensign; R. Cone; J. Hanes, Oral drug delivery with polymeric nanoparticles: The gastrointestinal mucus barriers, *Adv. Drug Deliv. Rev.* 64 (2012) 557-570.
86. M. E. Lowe, Medical treatment of chronic pancreatitis. In: *The pancreas* Beger, H. G.; Warshaw, A. L.; Hruban, R. H.; Büchler, M. W.; Lerch, M. M.; Neoptolemos, J. P.; Shimosegawa, T.; Whitcomb, D. C.; Groß, C., Eds. John Wiley & Sons Ltd. : Oxford, UK, 2018; pp 426-428.
87. Q. Guo; A. Ye; N. Bellissimo; H. Singh; D. Rousseau, Modulating fat digestion through food structure design, *Prog. Lipid Res.* 68 (2017) 109-118.
- 88. S. Onvani; F. Haghighatdoost; P. J. Surkan; L. Azadbakht, Dairy products, satiety and food intake: A meta-analysis of clinical trials, *Clin. Nutr.* 36 (2017) 389-398. Through analysis of clinical trials, this study analyses the effect of dairy intake on satiety. It shows that the consumption of 500mL of dairy products significantly increased satiety in consumers.
89. D. Ma; Z. C. Tu; H. Wang; Z. Zhang; D. J. McClements, Microgel-in-microgel biopolymer delivery systems: Controlled digestion of encapsulated lipid droplets under simulated gastrointestinal conditions, *J. Agric. Food Chem.* 66 (2018) 3930-3938.