Clear language for ecosystem management in the Anthropocene: a reply to Bridgewater & Hemming

Tina Heger^{1,2,3}, Maud Bernard-Verdier^{4,2}, Arthur Gessler^{5,6,2}, Alex D. Greenwood^{7,8,2}, Hans-Peter Grossart^{9,10,2}, Monika Hilker^{4,2}, Silvia Keinath^{11,2}, Ingo Kowarik^{12,2}, Elisabeth Marquard^{4,13,2}, Johannes Müller^{11,2}, Stephanie Niemeier^{11,2}, Gabriela Onandia^{6,2}, Jana S. Petermann^{14,2}, Matthias C. Rillig^{4,2}, Mark-Oliver Rödel^{11,2}, Wolf-Christian Saul^{15,4,10,2}, Conrad Schittko^{1,2}, Klement Tockner^{16,2}, Jasmin Joshi^{1,2,17} & Jonathan M. Jeschke^{4,10,2}

¹ University of Potsdam, Biodiversity Research/Systematic Botany, Potsdam, Germany

² Berlin-Brandenburg Institute of Advanced Biodiversity Research (BBIB), Berlin, Germany

³ Technical University of Munich, Restoration Ecology, Freising, Germany

⁴ Freie Universität Berlin, Institute of Biology, Berlin, Germany

⁵ Swiss Federal Research Institute WSL, Forest Dynamics, Birmensdorf, Switzerland

⁶ Leibniz Centre for Agricultural Landscape Research (ZALF), Müncheberg, Germany

⁷ Leibniz Institute for Zoo and Wildlife Research, Berlin, Germany

⁸ Freie Universität Berlin, Department of Veterinary Medicine, Berlin, Germany

⁹ University of Potsdam, Institute of Biochemistry and Biology, Potsdam, Germany

¹⁰ Leibniz-Institute of Freshwater Ecology and Inland Fisheries (IGB), Berlin, Germany

¹¹ Museum für Naturkunde – Leibniz Institute for Evolution and Biodiversity Science, Berlin, Germany

¹² Technische Universität Berlin, Department of Ecology, Ecosystem Science/Plant Ecology, Berlin, Germany

¹³ UFZ – Helmholtz Centre for Environmental Research GmbH, Department Conservation Biology, Leipzig, Germany

¹⁴ University of Salzburg, Department of Biosciences, Salzburg, Austria

¹⁵ Stellenbosch University, Centre for Invasion Biology (CIB), Department of Botany and Zoology & Department of Mathematical Sciences, Matieland, South Africa

¹⁶ Austrian Science Funds – FWF, Vienna, Austria

¹⁷ Institute for Landscape and Open Space, HSR Hochschule für Technik, Rapperswil, Switzerland

In a comment on our recent article (Heger et al. 2019), Bridgewater and Hemming (2020) call for putting the umbrella concept “ecological novelty” into a policy context. They suggest linking ecological novelty to the “nature-based solution” paradigm and comment that “*Global policy discussions around the ‘New Deal for Nature’ seem to only embrace traditional conservation paradigms, leaving ecological novelty in a policy vacuum*” (Bridgewater and Hemming 2020). We agree with Bridgewater and Hemming (2020) that political frameworks need new ways of expressing at which forms of “nature” conservation actions should aim, and we also share their opinion that the umbrella concept of “ecological novelty” can be helpful here. For instance, in the recently published ‘Zero Draft’ for a Post-2020 Global Biodiversity Framework (CBD 2020), “*recovery of natural ecosystems*” is formulated as one of the main aims for the CBD member states for the following 20 years; however, what the term “natural ecosystems” means in the Anthropocene remains an open question.

As we detail in Heger et al. (2019), “ecological novelty” can be used to describe and quantify the degree to which an environment differs from reference conditions. We believe that a scientific assessment of the ecological novelty of current as well as of projected future states and functions of study systems can provide a solid basis for management decisions in the Anthropocene. There is an array of options for how to implement nature-based solutions, and it is essential to make informed decisions matching local contexts. For example, reforestation has been suggested as a powerful means to sequester CO₂. Depending on how this measure is implemented, the outcome may be a highly novel system optimized for fast growth and high sequestration rates (thus increasing these ecosystem services), but using trees that have not been present in the area before with the potential to negatively affect native biodiversity; or a slow-growing forest that is similar to a historic reference – lending

additional benefit for biodiversity conservation, but possibly being less efficient concerning fast sequestration of CO₂.

The “ecological novelty” framework can help to enhance clear communication and allow for openly discussing pros and cons of such measures. Different aspects of novelty that already occupy major sectors of environmental policy, such as biological invasions, genetically modified organisms or habitat degradation, can be brought together under one umbrella concept to provide an integrated analysis of ecosystems. Based on an assessment of ecological novelty of the study system compared to reference conditions, it can be discussed whether management measures should be taken (arrow 1 in Fig. 1), for example in order to reduce novelty, and planned interventions can be evaluated in the light of their projected outcomes (arrow 2 in Fig. 1).

Figure 1: The umbrella concept “ecological novelty” allows to describe current states of ecosystems as well as projected outcomes of management measures and other human interventions. The concept could thus be used to clarify what is meant with “nature”, and which state of nature is tolerated or aimed for with policy regulations and management measures.

Future research should aim at developing methods for quantifying the novelty of a system and for predicting the consequences of novelty for ecosystem functioning, evolution, and biodiversity conservation. These scientific descriptions and assessments can then form the basis for a productive search for local consensus on whether or not ecological novelty can be tolerated, prior to action as well as in outcomes of interventions.

Implementing the concept of ecological novelty in environmental policy could thus allow for more explicit discussions about the aims of nature conservation in the Anthropocene. Using a common concept and speaking the same language will be useful in such discussions, particularly because opinions usually are highly context-dependent. For example, the course of discussions about management aims will be different if the focal system is a city (e.g., Planchuelo et al. 2019) versus a nature reserve.

Since the Rio-Declaration (UNCED 1992), international environmental policy does no longer aim for conservation of pristine nature only. This is reasonable, as in the face of the ongoing biodiversity crisis there is an urgent need to discuss and implement nature conservation actions as broadly and diversely as possible. We hope the concept of ecological novelty can enhance such discussions and help reach local consensus on conservation actions.

References Cited

Bridgewater P, Hemming K. 2020. Ecological novelty is inevitable, can be positive, but need policy context; a comment on Heger and colleagues (2019). Bioscience.

CBD. 2020. Zero Draft of the Post-2020 Global Biodiversity Framework.

<https://www.cbd.int/doc/c/efb0/1f84/a892b98d2982a829962b6371/wg2020-02-03-en.pdf>.

Heger T, et al. 2019. Towards an Integrative, Eco-Evolutionary Understanding of Ecological Novelty: Studying and Communicating Interlinked Effects of Global Change. *Bioscience* 69:888-899.

Planchuelo G, Lippe M, Kowarik I. 2019. Untangling the role of urban ecosystems as habitats for endangered plant species. *Landsc Urban Plann* 189. DOI: 10.1016/j.landurbplan.2019.05.007

UNCED. 1992. The Rio Declaration.

https://www.un.org/en/development/desa/population/migration/generalassembly/docs/globalcompact/A_CONF.151_26_Vol.I_Declaration.pdf in Development UNCoEa, editor.