**Characterization of nuclear microsatellite loci in the calcareous fen
specialist *Scorpidium cossonii* (Calliergonaceae)¹**

Yuppayao Kophimai^{2,4}, Carolina Cornejo², Markus Peintinger³, Ariel Bergamini² and
Christoph Scheidegger²

²WSL Swiss Federal Research Institute, Zürcherstrasse 111, CH-8903 Birmensdorf,
Switzerland

³Institute of Evolutionary Biology and Environmental Studies, University of Zurich,
Winterthurerstrasse 190, CH-8057 Zürich, Switzerland

Email addresses: YK: yuppayao.kophimai@wsl.ch

CC: carolina.cornejo@wsl.ch

MP: peinti@t-online.de

AB: ariel.bergamini@wsl.ch

CS: christoph.scheidegger@wsl.ch

Number of words: 961

¹Manuscript received_____; revision accepted_____.

⁴Author for correspondence: yuppayao.kophimai@wsl.ch

The authors thank the Genetic Diversity Centre, ETH Zurich, for technical assistance and S.
Werth for comments on a previous version of the manuscript. This study was supported by a
scholarship to YK (Higher Educational Strategic Scholarships for Frontier Research
Network) and grants from the Federal Office for the Environment (FOEN) to CS (MfM-U
Biodiversity) and AB (06.0126.PZ / I244-0978).

ABSTRACT

- *Premise of the study:* Nuclear microsatellite markers were developed in the fen specialist moss, *Scorpidium cossonii* (Schimp.) Hedenäs, to study genetic diversity and genetic structure of this species in relation to land-use types, habitat fragmentation and habitat conservation measures.
- *Methods and Results:* Fourteen polymorphic microsatellite markers were isolated and characterized using an enrichment method. All primers were tested on 140 gametophytes collected from four populations in Switzerland. The primers amplified di- and trinucleotide repeats with three to 26 alleles per locus. The primers worked also in the two other species of the genus: All and twelve primers successfully amplified in *S. revolvens* and *S. scorpioides*, respectively.
- *Conclusions:* In future studies, our primers will provide valuable information on genetic diversity, genetic structure as well as on historical and recent gene flow in *S. cossonii* and they enable related research in other *Scorpidium* species.

Key words: bryophytes; calcareous fens; cross-species amplification; microsatellites; *Scorpidium cossonii*

INTRODUCTION

Scorpidium cossonii (Schimp.) Hedenäs is a locally abundant pleurocarpous moss species (Hedenäs, 1989) which is characteristic for calcareous fens. In Central Europe, calcareous fens belong to the most species rich habitats. But despite legal protection, these fens are highly endangered by land-use changes, disturbances of the hydrology, nutrient enrichment and habitat fragmentation (Lienert et al., 2002; Peintinger et al., 2003; Bergamini et al., 2009). *S. cossonii* is a haploid and dioicous moss species (Hedenäs, 1989). Sporophytes are rarely produced and local population maintenance is presumably mainly by clonal growth. Its vegetative spread, however, seems to be quite restricted, at least in the short term (Hajkova et al., 2009). Nevertheless, the species is widely distributed in northern arctic to temperate regions and at high elevations in the Andes (Hedenäs, 2009).

In this study, we present the first set of polymorphic nuclear microsatellite markers for *S. cossonii*, a species assumed to be suffering from isolation and habitat deterioration. These markers will subsequently be applied to study the effects of habitat fragmentation and land-use types on the genetic diversity at different spatial scales as well as the genetic differentiation between populations of this fen specialist species.

METHODS AND RESULTS

We pooled DNA of *S. cossonii* specimens sampled from six different populations in Northern Switzerland. Ecogenics GmbH (Zurich, Switzerland) developed a genetic library from size selected genomic DNA ligated into SNX forward/SNX reverse linker (Hamilton et al., 1999) and enriched by magnetic bead selection with biotin-labelled (CT)₁₃, (GT)₁₃, (AAC)₁₀ and (ATC)₁₀ oligonucleotide repeats (Gautschi et al., 2000a,b). Of 528 recombinant

colonies screened, 341 gave a positive signal after hybridization. Plasmids from 48 positive clones were sequenced and primers were designed for 20 microsatellite inserts, of which 14 were tested for polymorphism.

The microsatellite loci were tested for variability on 140 gametophytes of *S. cossonii* collected from four populations in the pre-Alps of north-eastern Switzerland. Total genomic DNA was extracted from stem tissue using the DNeasy 96 plant kit (Qiagen, Hombrechtikon, Switzerland). All forward primers were labelled with one fluorescent dye at the 5' end region and three PCR multiplexes were set up (Table 1). PCR reactions were performed in a volume of 5 µl containing about 1-5 ng genomic DNA, varying primer concentrations (Table 1) and 1x multiplex PCR master mix (Qiagen). PCR cycles were started with 15 min at 95°C initial denaturation, followed by 30 cycles of 30 s at 94 °C, 90 s at annealing temperature (Table1), 1 min at 72 °C, ending with a final extension step of 60 min at 60°C. PCR products were run on a 3130xl Genetic Analyser (Applied Biosystems, Foster City, California, USA) using LIZ-500 as internal size standard and alleles were sized with GENEMAPPER 3.7 (Applied Biosystems). The variability of each microsatellite locus was measured by counting the number of alleles and calculating gene diversity using Arlequin version 3.11 (Excoffier et al., 2005).

Among the 14 microsatellite motifs, nine were dinucleotide repeats; three were trinucleotide repeats and two were combinations between two different trinucleotide repeats (Table 1). The microsatellite loci produced three to 26 alleles per locus and mean gene diversities over four populations varied from 0.13 to 0.68 (Table 2). Three individuals of each of two closely related species (Hedenäs et al., 2009), *S. revolvens* (Sw. ex anon.) Rubers and *S. scorpioides* (Hedw.) Limpr. were tested for cross-species amplifications. After PCR

optimization for the annealing temperature, all 14 primers successfully amplified in *S. revolvens* and 12 primers (except Sc2 and Sc22) in *S. scorpioides*.

CONCLUSIONS

The polymorphic microsatellite loci reported here are a promising tool for investigations of current genetic diversity, genetic structure and historical as well as recent gene flow in *S. cossonii*. This research will considerably improve our basic understanding of the genetic structure of fragmented bryophyte populations and it will serve as a base for the development of conservation strategies in this highly specialised fen species. Furthermore, because these loci are applicable to other *Scorpidium* species, they can be widely used and a number of possible studies using these microsatellites are in progress.

LITERATURE CITED

- BERGAMINI, A., M. PEINTINGER, S. FAKHERAN, H. MORADI, B. SCHMID, AND J. JOSHI. 2009. Loss of habitat specialists despite conservation management in fen remnants 1995-2006. *Perspectives in Plant Ecology Evolution and Systematics* 11: 65–79.
- EXCOFFIER, L., G. LAVAL, AND S. SCHNEIDER. 2005. Arlequin ver. 3.0: An integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online* 1: 47–50.
- GAUTSCHI, B., A. WIDMER, AND J. KOELLA. 2000a. Isolation and characterization of microsatellite loci in the dice snake (*Natrix tessellata*). *Molecular Ecology* 9: 2191–2193.
- GAUTSCHI, B., I. TENZER, J. P. MÜLLER, AND B. SCHMID. 2000b. Isolation and characterization of microsatellite loci in the bearded vulture (*Gypaetus barbatus*) and cross-amplification in three Old World vulture species. *Molecular Ecology* 9: 2193–2195.

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
 - 61
 - 62
 - 63
 - 64
 - 65
- HAIKOVA, P., M. HAJEK, AND K. KINTROVA. 2009. How can we effectively restore species richness and natural composition of a *Molinia*-invaded fen? *Journal of Applied Ecology* 46: 417–425.
- HAMILTON, M. B., E. L. PINCUS, A. DI FIORE, AND R. C. FLEISCHER. 1999. Universal linker and ligation procedures for construction of genomic DNA libraries enriched for microsatellites. *Biotechniques* 27: 500–507.
- HEDENÄS L. 1989. The genera *Scorpidium* and *Hamatocaulis*, gen. nov., in northern Europe. *Lindbergia* 15: 8–36.
- HEDENÄS L. 2009. Relationships among arctic and non-arctic haplotypes of the moss species *Scorpidium cossonii* and *Scorpidium scorpioides* (Calliergonaceae). *Plant Systematics and Evolution* 277: 217–231.
- LIENERT, J., M. FISCHER, J. SCHNELLER, AND M. DIEMER. 2002. Isozyme variability of the wetland specialist *Swertia perennis* (Gentianaceae) in relation to habitat size, isolation, and plant fitness. *American Journal of Botany* 89: 801–811.
- PEINTINGER, M., A. BERGAMINI, AND B. SCHMID. 2003. Species-area relationships and nestedness of four taxonomic groups in fragmented wetlands. *Basic and Applied Ecology* 4: 385–394.

TABLE 1. Characteristics of 14 nuclear microsatellite loci in *S. cossonii*. For each locus, the forward and reverse primer sequences, repeat motif, labelled fluorescent dye, size of the original fragment, PCR multiplex group, primer concentration, annealing temperature when run in multiplex (Ta) and GenBank accession number are shown.

Locus	Primer sequence forward/reverse	Repeat motif	Fluorescent dye	Size (bp)	Multiplex	Primer conc. (μM)	Ta (°C)	GenBank accession no.
Sc01	AGCCAAAACGTATGGAAACC GATAGGAAGGCAGTGCAACC	(AG) ₂₄	FAM	178	1	0.14 0.14	59	JF700218
Sc02	AAATGGAGTATTGGATGAGATGG CTAATGCAAGGCCATGAGTG	(AC) ₁₃	NED	159	1	0.21 0.21	59	JF700219
Sc03	CTCAAGTCACGGCTGCTTC CAAAGGGACTTCAATTCCAAC	(GT) ₁₆	NED	149	2	0.08 0.08	59	JF700220
Sc04	TGTGAACGTGGAAGCTTGTG GCCCCAAGTAGTTGCTTCTC	(GT) ₁₇	PET	137	1	0.07 0.07	59	JF700221
Sc07	AGGATGAGAGTTCCTGGTG TGGCATTCAAACACACATCC	(CA) ₁₃	FAM	205	3	0.08 0.08	60	JF700222
Sc09	ACGTCAAGTGACCCACAC TTGGGTAGTTTGGCGTTAGG	(AC) ₂₂	VIC	156	1	0.07 0.07	59	JF700223
Sc13	ACAGCAGCACATGTCCAGAG GAGGTAATCCAGCAGCCATC	(CAA) ₉	PET	143	2	0.08 0.08	59	JF700224
Sc16	AATCAAGGCCCCATCTAAGC TCACAGATCGCAAACAAACC	(CAT) ₁₃ (CAA) ₄	FAM	208	2	0.08 0.08	59	JF700225
Sc17	GTGTGAACCTGCACGAGATG CATGGCCGTAAGAGTGGAAC	(GTT) ₈ (GCT) ₆	VIC	191	2	0.08 0.08	59	JF700226
Sc18	CTTCAGAAAGCCGATCAAGC TTGCGGAAGATCTCCTCTTG	(CAA) ₈	NED	127	3	0.08 0.08	60	JF700227
Sc19	GCGCTTGATAAGCCTGAGC TGCCACTAACGAAGGGACTG	(GTT) ₇	PET	135	3	0.08 0.08	60	JF700228
Sc20	CGCTCTAATTGAGGCATTG TGTTTCGATTTCGACAAG	(AC) ₂₀	VIC	172	3	0.16 0.16	60	JF700229
Sc21	TGGAGGCTTGTCAGAATTG CGACACTGCAAACCAACTAGAC	(TG) ₁₃	FAM	93	3	0.08 0.08	60	JF700230
Sc22	GGCATCCCTACCTATTGACCG GGGGGCTTTCACCAGGAATTCA	(CA) ₂₂	FAM	159	2	0.16 0.16	59	JF700231

TABLE 2. Results of the initial primer screening in four populations of *S. cossonii*. Shown for each primer are the number of alleles observed (N_A) and gene diversity (He). Sample sizes per population and geographical coordinates (Swiss grid) are given in parentheses.

	Population 1 (N=35) (725.83 E, 234.95 N)		Population 2 (N=35) (703.45 E, 222.48 N)		Population 3 (N=35) (712.21 E, 220.75 N)		Population 4 (N=35) (707.16 E, 220.40 N)		Total	Mean
Locus	N_A	He	N_A	He	N_A	He	N_A	He	N_A	He
Sc01	9	0.82	11	0.89	7	0.31	6	0.70	26	0.68
Sc02	4	0.66	6	0.58	4	0.31	7	0.67	13	0.55
Sc03	3	0.51	3	0.54	3	0.26	7	0.70	8	0.50
Sc04	4	0.66	4	0.57	6	0.35	3	0.54	9	0.53
Sc07	3	0.64	4	0.39	2	0.05	5	0.62	8	0.42
Sc09	4	0.66	7	0.65	5	0.31	7	0.70	13	0.58
Sc13	2	0.44	1	0.00	2	0.05	2	0.05	3	0.13
Sc16	3	0.62	2	0.48	3	0.25	2	0.05	4	0.35
Sc17	2	0.11	4	0.43	2	0.05	3	0.55	5	0.28
Sc18	3	0.68	3	0.54	3	0.21	4	0.58	5	0.50
Sc19	3	0.65	3	0.54	3	0.25	3	0.58	3	0.50
Sc20	2	0.11	2	0.39	2	0.11	2	0.49	3	0.27
Sc21	3	0.60	3	0.51	3	0.11	2	0.20	3	0.35
Sc22	4	0.75	4	0.56	4	0.21	7	0.70	9	0.55
Mean	3.50	0.57	4.07	0.50	3.50	0.20	4.29	0.51	8.00	0.44
SD	1.74	0.21	2.53	0.19	1.56	0.11	2.13	0.23	6.26	0.24